

## References

- [1] A. Baker, *Linear forms in the logarithms of algebraic numbers II*, *Mathematika* 14 (1967), pp. 102–107.
- [2] P. Erdős, *A theorem of Sylvester and Schur*, *J. London Math. Soc.* 9 (1934), pp. 282–288.
- [3] — *On consecutive integers*, *Nieuw Arch. Voor Wisk.* 3 (1955), pp. 124–128.
- [4] M. N. Huxley, *On the difference between consecutive primes*, *Invent. Math.* 15 (1972), pp. 164–170.
- [5] K. Ramachandra, *A note on numbers with a large prime factor II*, *J. Indian Math. Soc.* 34 (1970), pp. 39–48.
- [6] — *A note on numbers with a large prime factor III*, *Acta Arith.* 19 (1971), pp. 49–62.
- [7] — *Application of Baker's theory to two problems considered by Erdős and Selfridge*, *J. Indian Math. Soc.* (to appear).
- [8] — *Largest prime factor of the product of  $k$ -consecutive integers*, *Reports of an International Conference on Number Theory (Moscow, 1971)*, (to appear).
- [9] — *Lectures on Transcendental Numbers*, *Ramanujan Institute, Madras 1969*.
- [10] T. N. Shorey, *Linear forms in the logarithms of algebraic numbers with small coefficients I*, *J. Indian Math. Soc.* (to appear).
- [11] R. Tijdeman, *On the maximal distance of numbers with a large prime factor*, *J. London Math. Soc.* 5 (1972), pp. 313–320.
- [12] — *An auxiliary result in the theory of transcendental numbers*, *Journal of Number Theory* (to appear).

TATA INSTITUTE OF FUNDAMENTAL RESEARCH  
Bombay, India

Received on 29. 6. 1972

(299)

ACTA ARITHMETICA  
XXIV (1973)

## ERRATA

Page, line	For	Read
35 <sup>16</sup>	$M - \sqrt{C} + 1h$	$M - \sqrt{C} h$
39 <sub>8</sub>	$s > 4nk$	$s > 4nk$
43 <sub>10</sub>	$a \bmod p^0$	$a \bmod p^l$
50 <sub>6</sub>	$r_1 + r_2$ $\prod_{m=r_1+r}$	$r_1 + r_2$ $\prod_{m=r_1+1}$