

in (9.23) each summand by $(x-1)/2$ it follows

$$(9.24) \quad \sum_r \text{Min} \left(\frac{x-1}{2}, r' \right) = (\delta+2t) \frac{x-1}{2} = \frac{s-1}{2} (x-1).$$

33. Using now (9.8) and summing over r we obtain from (9.22) and (9.24) the same expression as in (9.21). It follows finally

$$(9.25) \quad sg(x) = \left(s - \frac{1+\delta}{2} \right) x - t \quad (1-1/s \leq x \leq 1+1/s).$$

Since obviously

$$sg(x) - sx/2 = t(x-1),$$

we see that $g(x)$ in $\langle 1-1/s, 1+1/s \rangle$ is always different from $x/2$ save for $x=1$. The assertion of Section 25 is proved.

References

- [1] L. Fejér, *Gesammelte Arbeiten*, Bd. II, Birkhäuser, Basel 1970, p. 843.
- [2] E. Hecke, *Über analytische Funktionen und die Verteilung der Zahlen mod. eins*, Hamb. Math. Abh. 1 (1922), pp. 54-76; see also: E. Hecke, *Mathematische Werke*, Vandenhoeck und Ruprecht, Göttingen 1959, p. 329.
- [3] A. M. Ostrowski, *Sur le formule de Moivre-Laplace*, C. R. Acad. Sci. Paris 223 (1946), pp. 1090-1092.
- [4] — *On the remainder term of the de Moivre-Laplace formula*, to appear in *Aequationes Math.*
- [5] — *Math. Miscellen LX, Notiz zur Theorie der Diophantischen Approximationen*, J.-B.d.D.M.V. 36 (1927), pp. 178-180.
- [6] G. Pólya und G. Szegő, *Aufgaben und Lehrsätze*, Bd. I, Springer, Berlin 1925, pp. 72, 237.
- [7] J. V. Uspensky, *Introduction to mathematical probability*, McGraw-Hill, New York 1937, p. 129.

Received on 13. 5. 1977

(941)

On two definitions of the integral of a p -adic function

by

KURT MAHLER (Canberra)

In memory of Paul Turán

In his basic paper on functions of a p -adic variable Dieudonné [1], introduced a special kind of integral (primitive) of a continuous function. A completely different definition of such an integral was more recently given by M. van der Put (see A. C. M. van Rooij and W. H. Schikhof [2]). The aim of this note is to show that these two definitions lead to the same result. This is rather surprising because there is a large set of non-constant p -adic functions of derivative 0.

Since it simplifies the discussion, we shall study the two kinds of integrals for the class of functions $f: J \rightarrow Q_p$ where p is any positive rational prime, Q_p is the field of p -adic numbers, and $J = \{0, 1, 2, \dots\}$ is the set of all non-negative rational integers. The set J is not closed, and its p -adic closure is the set $I = \{x \in Q_p; |x|_p \leq 1\}$ of all p -adic integers which is compact.

1. Let $f: J \rightarrow Q_p$ be an arbitrary function on J . The two integrals of f are defined by the following constructions.

Write $x \in J$ in the canonic form as

$$x = x_0 + x_1 p + x_2 p^2 + \dots$$

where x_0, x_1, x_2, \dots are digits $0, 1, \dots, p-1$. At most finitely many of these digits are distinct from 0; so, if $x \neq 0$, let $x_s \neq 0$ be the non-vanishing digit of largest suffix s . Firstly put

$$q(0) = 0, \quad q(x) = x_s p^s \quad \text{for } x \neq 0.$$

Secondly write

$$x^{(n)} = x_0 + x_1 p + \dots + x_{n-1} p^{n-1} \quad (n = 1, 2, 3, \dots)$$

so that

$$x^{(n+1)} = x^{(n)} \quad \text{for } n > s.$$

The Dieudonné integral of f is now defined by

$$D(x) = \sum_{n=1}^{\infty} (x^{(n+1)} - x^{(n)})f(x^{(n)}).$$

Since the terms of this series vanish for $n > s$, there is no problem of convergence. One can show that, whenever f is continuous at a point x_0 of J , then $D'(x_0) = f(x)$, as required for an integral.

2. Let m be any integer in J . With m we associate a positive integer M where

$$M = 1 \quad \text{if} \quad m = 0,$$

while for $m \geq 1$ the integer M is chosen such that

$$p^{M-1} \leq m \leq p^M - 1.$$

Denote by $S(m)$ the ball consisting of all $x \in J$ for which

$$|x - m|_p \leq p^{-M},$$

and by $X(x, m)$ the characteristic function of $S(m)$ defined by

$$X(x, m) = \begin{cases} 1 & \text{if } x \in S(m), \\ 0 & \text{otherwise.} \end{cases}$$

It can be proved that every function $f: J \rightarrow Q_p$ has a unique van der Put series

$$f(x) = \sum_{m=0}^{\infty} b_m X(x, m) \quad \text{for all } x \in J.$$

Here the coefficients b_m can be determined by the formulae

$$b_m = \begin{cases} f(m) & \text{if } m = 0, 1, \dots, p-1; \\ f(m) - f(m - q(m)) & \text{if } m \geq p. \end{cases}$$

Since

$$X(x, m) = 0 \quad \text{if } x < m,$$

the van der Put series for $f(x)$ breaks off after finitely many terms, and there is again no problem of convergence.

In the special case when $f(x)$ is the function x , we obtain the series

$$x = \sum_{m=0}^{\infty} q(m) X(x, m).$$

Once the van der Put series for $f(x)$ is known, its van der Put integral is defined by the development

$$P(x) = \sum_{m=0}^{\infty} b_m X(x, m)(x - m).$$

Also this integral satisfies the relation $P'(x_0) = f(x_0)$ at every point $x_0 \in J$ at which the function f is continuous, as it should be.

3. Without any restrictions on f we can now prove the following result.

THEOREM 1. For every function $f: J \rightarrow Q_p$,

$$D(x) = P(x) \quad \text{for all } x \in J.$$

Proof. The van der Put series for $f(x)$ shows that it suffices to prove this theorem only for all the characteristic functions

$$f(x) = X(x, m).$$

Denote therefore by $D(x, m)$ and $P(x, m)$ the Dieudonné and the van der Put integrals of $X(x, m)$; we must prove that

$$D(x, m) = P(x, m) \quad \text{for all } x \in J.$$

This will be done by evaluating these two integrals explicitly, and we shall begin with the more difficult function $D(x, m)$.

Let x be an arbitrary element of J so that also $x^{(n)} \in J$ for all $n \geq 1$. If $X(x^{(n)}, m) = 0$ for all $n \geq 1$, then $D(x, m) = 0$; we exclude this easy case. There is then a smallest integer $N \geq 1$ such that $x^{(N)} \in S(m)$. Then

$$|x^{(N)} - m|_p \leq p^{-M}$$

and therefore there is a rational integer x^* such that

$$x^{(N)} = m + p^M x^*.$$

Here

$$p^{M-1} \leq m \leq p^M - 1,$$

from which it follows that x^* cannot be negative because then

$$x^{(N)} \leq m - p^M \leq -1,$$

contrary to $x^{(N)} \in J$. Therefore either

$$(1) \quad x^{(N)} = m,$$

or

$$(2) \quad x^{(N)} \geq m + p^M \geq p^M.$$

Now

$$\begin{aligned} x^{(N)} &= x_0 + x_1 p + \dots + x_{N-1} p^{N-1} \\ &\leq (p-1) + (p-1)p + \dots + (p-1)p^{N-1} \leq p^N - 1. \end{aligned}$$

Hence, in the case (2),

$$p^M \leq x^{(N)} \leq p^N - 1$$



and therefore $N \geq M+1$. It would then follow that

$$x^{(N)} = x^{(M)} + x_M p^M + \dots + x_{N-1} p^{N-1},$$

and therefore

$$|x^{(N)} - x^{(M)}|_p \leq p^{-M},$$

whence also

$$|x^{(M)} - m|_p = |(x^{(M)} - x^{(N)}) + (x^{(N)} - m)|_p \leq p^{-M}.$$

Thus $x^{(M)} \in S(m)$, contrary to the minimum hypothesis for N .

Therefore the case (1) holds, and

$$(3) \quad x^{(N)} = m.$$

We assert that moreover

$$(4) \quad N = M.$$

For, if $N > M$, the proof just given leads to a contradiction; if, however, $N < M$, then

$$0 \leq x^{(N)} = m \leq p^N - 1 \leq p^{M-1} - 1 < p^{M-1},$$

and this likewise is false.

On account of (3) and (4) we can now prove that exactly

$$x^{(n)} \in S(m) \quad \text{for all } n \geq M.$$

For if $n \geq M+1$, we have again

$$x^{(n)} = x^{(M)} + x_M p^M + \dots + x_{n-1} p^{n-1}$$

and therefore

$$|x^{(n)} - x^{(M)}|_p = |x^{(n)} - m|_p \leq p^{-M},$$

as asserted.

The integral $D(x, m)$ can now be determined and is found to have the value

$$D(x, m) = \begin{cases} \sum_{n=M}^{\infty} (x^{(n+1)} - x^{(n)}) \times 1 = x - x^{(M)} = x - m & \text{if } x \in S(m), \\ 0 & \text{otherwise.} \end{cases}$$

For $x^{(n)}$ becomes equal to x as soon as n is sufficiently large.

Since by definition also $P(x, m) = X(x, m)(x - m)$, we have proved the theorem.

4. From any integral of the arbitrary function $f(x)$ we obtain others by adding any function the derivative of which vanishes identically. In the present p -adic case there are very many such almost-constants. For instance, as C. S. Weisman has proved, every function

$$\sum_{m=0}^{\infty} \beta_m X(x, m),$$

where

$$\lim_{m \rightarrow \infty} m |\beta_m|_p = 0,$$

has everywhere the derivative 0.

Since there is then such a great choice of possible integrals of $f(x)$, the question may be asked whether the special integral $D(x) = P(x)$ has any distinguishing properties.

I obtained one such property. Write $f(x)$ and $P(x)$ as interpolation series

$$f(x) = \sum_{n=0}^{\infty} a_n \binom{x}{n} \quad \text{and} \quad D(x) = \sum_{n=0}^{\infty} A_n \binom{x}{n}.$$

Then the coefficients A_n of the integral can be expressed as linear forms

$$(5) \quad A_n = \sum_{m=0}^{n-1} c_{mn} a_m \quad (n \geq 1)$$

where the coefficients c_{mn} are rational integers. This is quite different from the position for functions of a real variable where, e.g.

$$\int \binom{x}{2} dx = \binom{x}{3} + \frac{1}{2} \binom{x}{2} - \frac{1}{12} \binom{x}{1} + \text{constant}$$

with fractional rational coefficients. In the p -adic case the Diendonné-van der Put integral of $\binom{x}{2}$ is a rather more complicated infinite interpolation series

$$\sum_{n=1}^{\infty} c_{2n} \binom{x}{n}.$$

I shall establish and study the formulae (5) elsewhere.

References

- [1] J. Diendonné, *Sur les fonctions continues p-adiques*, Bull. Sci. Math. (2) 68 (1944), pp. 79-95.
- [2] A. C. M. van Rooij and W. H. Schikhof, *Non-archimedean analysis*, Nieuw Archief v. Wiskunde (2), 19 (1971), pp. 120-160.
- [3] C. S. Weisman, *On p-adic differentiability*, J. Number Theory 9 (1977), pp. 79-86.

MATHEMATICS DEPARTMENT
RESEARCH SCHOOL OF PHYSICAL SCIENCES
AUSTRALIAN NATIONAL UNIVERSITY
Canberra, ACT 2600, Australia