

## Literaturverzeichnis

- [1] N. C. Ankeny and C. A. Rogers, *A conjecture of Chowla*, Ann. of Math. 53 (1951), pp. 541–550.  
 [2] H. Flanders, *Generalisation of a theorem of Ankeny and Rogers*, *ibid.* 57 (1953), pp. 392–400.  
 [3] I. Gerst, *On the theory of  $n$ -th power residues and a conjecture of Kronecker*, Acta Arith. 17 (1970), pp. 121–139.  
 [4] A. Schinzel, *On power residues and exponential congruences*, *ibid.* 27 (1975), pp. 397–420.  
 [5] — *A refinement of a theorem of Gerst on power residues*, *ibid.* 17 (1970), pp. 161–168.  
 [6] — *Abelian binomials, power residues and exponential congruences*, *ibid.* 32 (1977), pp. 245–274.  
 [7] V. Schulze, *Potenzreste*, *ibid.* 33 (1977), pp. 379–404.  
 [8] E. Trost, *Zur Theorie der Potenzreste*, Nieuw. Arch. Wiskunde 18 (1934), pp. 58–61.

FREIE UNIVERSITÄT BERLIN  
 II. MATH. INST. FB 19  
 Königin-Luise-Str. 24/26  
 D-1000 Berlin 33

Eingegangen am 4. 2. 1980  
 und in revidierter Form am 28. 11. 1980

(1235)

## Some asymptotic formulas on generalized divisor functions, III

by

P. ERDÖS and A. SÁRKÖZY (Budapest)

1. Throughout this paper, we use the following notation:

$c_1, c_2, \dots, X_0, X_1, \dots$  denote positive absolute constants. We denote the number of elements of the finite set  $S$  by  $|S|$ . We write  $e^x = \exp(x)$ . We denote the least prime factor of  $n$  by  $p(n)$ , while the greatest prime factor of  $n$  is denoted by  $P(n)$ . We write  $p^a \parallel n$  if  $p^a | n$  but  $p^{a+1} \nmid n$ .  $\omega(n)$  denotes the number of all the prime factors of  $n$  so that  $\omega(n) = \sum_{p^a \parallel n} a$  and we write

$$\omega(n, x, y) = \sum_{\substack{p^a \parallel n \\ x < p \leq y}} a.$$

The divisor function is denoted by  $d(n)$ :

$$d(n) = \sum_{d|n} 1.$$

Let  $A$  be a finite or infinite sequence of positive integers  $a_1 < a_2 < \dots$ . Then we write

$$N_A(x) = \sum_{\substack{a \in A \\ a \leq x}} 1, \quad f_A(x) = \sum_{\substack{a \in A \\ a | x}} \frac{1}{a}, \quad d_A(n) = \sum_{\substack{a \in A \\ a | n}} 1$$

(in other words,  $d_A(n)$  denotes the number of divisors amongst the  $a_i$ 's) and

$$D_A(x) = \max_{1 \leq n \leq x} d_A(n).$$

The aim of this series is to investigate the function  $D_A(x)$ . (See [1] and [2]; see also Hall [4].) Clearly,

$$\sum_{1 \leq n \leq x} d_A(n) = x f_A(x) + O(x).$$

Thus if  $f_A(x)$  is large then we have  $D_A(x)/f_A(x) \gg 1$ . In Part II of this

series (see [2]), we showed that  $f_A(x) \rightarrow +\infty$  implies that

$$\limsup_{x \rightarrow +\infty} D_A(x)/f_A(x) = +\infty,$$

in fact, we have

$$\limsup_{x \rightarrow +\infty} D_A(x) \exp(c_1(\log f_A(x))^2) = +\infty.$$

We proved this by showing that if  $f_A(x)$  is large (in fact, it is sufficient to assume that  $N_A(x)$  is large) then there exists an integer  $y$  such that  $x \leq y \leq \exp((\log x)^2)$  and  $D_A(y)$  is large. In this paper, our aim is to prove that if we have more information about  $f_A(x)$  then  $D_A(x)/f_A(x)$  must be large for the same  $x$ . In fact we prove that

**THEOREM 1.** For all  $\omega > 0$  and for  $x > X_0(\omega)$ ,

$$(1) \quad f_A(x) > (\log \log x)^{20}$$

implies that

$$(2) \quad D_A(x) > \omega f_A(x).$$

(Note that by Theorem 1 in [1], the lower bound  $(\log \log x)^{20}$  in (1) cannot be replaced by  $\log \log x$ .)

Sections 2 and 3 are devoted to the proof of this theorem while in Sections 4 and 5 we discuss some other related results.

2. In order to prove Theorem 1, we need some lemmas.

**LEMMA 1.** There exists an absolute constant  $c_2$  such that for all  $u \geq 0$  and  $y \geq 1$  we have

$$(3) \quad \sum_{\substack{u < n \leq uy \\ \pi(n) > y}} \frac{1}{n} < c_2.$$

*Proof.* Lemma 1 can be proved easily by using Brun's sieve. In fact, (3) is trivial for  $u \leq 1$  (since in this case, the left-hand side is equal to 0), while for  $u > 1$ , (3) is a consequence of [7], p. 53, Theorem 4.10.

**LEMMA 2.** Let us write

$$(4) \quad Q(x) = x - (1+x) \log(1+x).$$

Then for  $1 \leq y, 2y < z \leq v, 0 \leq \alpha \leq 1$  we have

$$(5) \quad \sum_{\substack{n \leq v \\ \omega(n, y, z) \leq (1-\alpha) \\ \sum_{y < p \leq z} 1/p}} 1 < c_3 v \exp\left(Q(-\alpha) \log \frac{\log z}{\log y}\right).$$

*Proof.* Let  $1 \leq v, 0 \leq \alpha \leq 1$ , and let  $E$  be an arbitrary nonempty set of prime numbers not exceeding  $v$ . Put  $E(v) = \sum_{p \in E} 1/p$ . K. K. Norton

proved (see [5], Theorem (5.9); see also Halász [3]) that

$$\sum_{\substack{n \leq v \\ p^\beta | n, p \in E}} 1 < c_4 v \exp(Q(-\alpha) E(v)).$$

By using this theorem with  $E = \{p: y < p \leq z\}$  (note that  $E \neq \emptyset$  by  $2y < z$ ), and with respect to the well-known formula

$$(6) \quad \sum_{p \leq x} 1/p = \log \log x + c_5 + o(1),$$

we obtain (5).

**LEMMA 3.** For  $1 \leq y, 2y < z \leq v, 0 < \alpha \leq \beta < 1$  we have

$$(7) \quad \sum_{\substack{n \leq v \\ \omega(n, y, z) \geq (1+\alpha) \\ \sum_{y < p \leq z} 1/p}} 1 < c_6(\beta) \alpha^{-1} v \left(\sum_{y < p \leq z} \frac{1}{p}\right)^{-1/2} \exp\left(Q(\alpha) \log \frac{\log z}{\log y}\right)$$

(where  $Q(x)$  is defined by (4)).

*Proof.* Let  $0 < \alpha \leq \beta < 1$ , and let  $E$  be an arbitrary nonempty set of prime numbers not exceeding  $v$ . Put  $E(v) = \sum_{p \in E} 1/p$ . K. K. Norton proved (see [5], Theorem (5.12); see also Halász [3]) that

$$\sum_{\substack{n \leq v \\ \sum_{p \in E} 1/p \geq (1+\alpha) E(v)}} 1 < c_7(\beta) \alpha^{-1} v (E(v))^{-1/2} \exp(Q(\alpha) E(v)).$$

By using this theorem with  $E = \{p: y < p \leq z\}$  (again,  $E \neq \emptyset$  by  $2y < z$ ), and with respect to (6), we obtain (7).

3. In this section, we complete the proof of Theorem 1. Define the positive integer  $R$  by

$$(8) \quad x^{1-1/2^{R-1}} \leq x e^{-f_A(x)/3} < x^{1-1/2^R},$$

i.e.,

$$2^{R-1} < \frac{3 \log x}{f_A(x)} \leq 2^R, \quad R-1 < \frac{1}{\log 2} \log \frac{3 \log x}{f_A(x)} \leq R.$$

Then for large  $x$ , we have

$$(9) \quad R < \frac{1}{\log 2} \log \frac{3 \log x}{f_A(x)} + 1 < 2 \log \log x.$$

For  $i = 0, 1, \dots, R$ , let

$$x_i = x^{1-1/2^i},$$

and for  $i = 1, 2, \dots, R$ , put

$$A_i = A \cap [x_{i-1}, x_i];$$

Then by (1) and (8), for large  $x$  we have

$$(10) \quad \sum_{i=1}^R \left( \sum_{a \in A_i} \frac{1}{a} \right) = \sum_{\substack{a \in A \\ a < x_R}} \frac{1}{a} = \sum_{\substack{a \in A \\ a < x}} \frac{1}{a} - \sum_{\substack{a \in A \\ x_R \leq a < x}} \frac{1}{a}$$

$$\geq f_A(x) - \sum_{x_R \leq n < x} \frac{1}{n} \geq f_A(x) - \sum_{x e^{-f_A(x)/3} \leq n < x} \frac{1}{n}$$

$$= f_A(x) - (1 + o(1)) \log e^{f_A(x)/3} > f_A(x) - \frac{f_A(x)}{2} = \frac{f_A(x)}{2}.$$

Obviously, there exists an integer  $j$  such that  $1 \leq j \leq R$  and

$$\sum_{a \in A_j} \frac{1}{a} \geq \frac{1}{R} \sum_{i=1}^R \left( \sum_{a \in A_i} \frac{1}{a} \right)$$

hence with respect to (9) and (10),

$$(11) \quad f_{A_j}(x) = \sum_{a \in A_j} \frac{1}{a} > \frac{1}{R} \frac{f_A(x)}{2} > \frac{f_A(x)}{4 \log \log x}.$$

Let us fix an integer  $j$  ( $1 \leq j \leq R$ ) satisfying (11), and write  $A_j$  in the form

$$(12) \quad A_j = A'_j \cup A''_j$$

where  $A'_j$  consists of the integers  $a$  such that  $a \in A_j$  and there exists an integer  $d$  satisfying

$$(13) \quad (\log x)^3 < d \leq x^{1/2^j + 1} \omega f_A(x)$$

and  $d|a$ , while  $A''_j$  consists of the integers  $a$  such that  $a \in A_j$  and  $d \nmid a$  for all  $d$  satisfying (13). (For  $x^{1/2^j + 1} \omega f_A(x) \leq (\log x)^3$ , we have  $A'_j = \emptyset$ .) We have to distinguish two cases.

Case 1. Assume first that

$$f_{A'_j}(x) = \sum_{a \in A'_j} \frac{1}{a} > \frac{1}{2} f_{A_j}(x).$$

Then by (11), we have

$$(14) \quad f_{A'_j}(x) = \sum_{a \in A'_j} \frac{1}{a} > \frac{1}{2} f_{A_j}(x) > \frac{f_A(x)}{8 \log \log x}.$$

For  $a \in A'_j$ , write  $a$  in the form

$$a = d^*(a) b(a),$$

where  $d^*(a)$  denotes the least integer  $d$  such that  $d$  satisfies (13) and  $d|a$ . Then for  $a \in A'_j$  we have  $b(a) \leq a < x_j = x^{1-1/2^j}$  and  $(\log x)^3 < d^*(a)$  so that

$$(15) \quad \sum_{a \in A'_j} \frac{1}{a} = \sum_{a \in A'_j} \frac{1}{d^*(a) b(a)} = \sum_{b \leq x^{1-1/2^j}} \frac{1}{b} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} \frac{1}{d^*(a)}$$

$$< \sum_{b \leq x^{1-1/2^j}} \frac{1}{b} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} \frac{1}{(\log x)^3} = \frac{1}{(\log x)^3} \sum_{b \leq x^{1-1/2^j}} \frac{1}{b} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} 1$$

$$\leq \frac{1}{(\log x)^3} \left( \max_{1 \leq b \leq x^{1-1/2^j}} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} 1 \right) \sum_{b < x} \frac{1}{b}$$

$$< \frac{1}{(\log x)^3} \left( \max_{1 \leq b \leq x^{1-1/2^j}} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} 1 \right) 2 \log x = \frac{2}{(\log x)^2} \left( \max_{1 \leq b \leq x^{1-1/2^j}} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} 1 \right).$$

If  $x$  is large enough (in terms of  $\omega$ ) then (14) and (15) yield that

$$\max_{1 \leq b \leq x^{1-1/2^j}} \sum_{\substack{a \in A'_j \\ b^*(a)=b}} 1 > \frac{(\log x)^2}{2} \sum_{a \in A'_j} \frac{1}{a} > \frac{(\log x)^2}{16 \log \log x} f_{A'}(x) > \omega f_A(x) + 1$$

so that there exists an integer  $b_0$  for which

$$(16) \quad 1 \leq b_0 \leq x^{1-1/2^j}$$

and

$$(17) \quad \sum_{\substack{a \in A'_j \\ b^*(a)=b_0}} 1 > \omega f_A(x) + 1.$$

Put  $s = [\omega f_A(x)] + 1$ . Then by (17), there exist distinct integers  $a_1, a_2, \dots, a_s$  such that  $a_i$  can be written in the form

$$a_i = b_0 d^*(a_i) = b_0 d_i$$

where

$$(18) \quad ((\log x)^3 < d_i \leq x^{1/2^j + 1} \omega f_A(x)).$$

Let

$$u = b_0 d_1 d_2 \dots d_s.$$

Then by (16) and (18), we have

$$(19) \quad u = b_0 d_1 d_2 \dots d_s \leq x^{1-1/2^j} (x^{1/2^j + 1} \omega f_A(x))^s < x^{1-1/2^j} (x^{1/2^j + 1} \omega f_A(x))^{2 \omega f_A(x)} = x,$$

and obviously,  $a_i = b_0 d_i / u$  and  $a_i = b_0 d_i \in A$  so that

$$(20) \quad d_A(u) \geq s = [\omega f_A(x)] + 1 > \omega f_A(x).$$

(19) and (20) yield (2) and this completes the proof of Theorem 1 in this case.

Case 2. Assume now that

$$(21) \quad f_{A'_j}(x) = \sum_{a \in A'_j} \frac{1}{a} \leq \frac{1}{2} f_{A_j}(x) = \frac{1}{2} \sum_{a \in A_j} \frac{1}{a}.$$

Then (12) and (21) yield that

$$(22) \quad f_{A''_j}(x) = \sum_{a \in A''_j} \frac{1}{a} \geq \sum_{a \in A_j} \frac{1}{a} - \sum_{a \in A'_j} \frac{1}{a} \\ \geq f_{A_j}(x) - \frac{1}{2} f_{A_j}(x) = \frac{1}{2} f_{A_j}(x) > \frac{f_A(x)}{8 \log \log x}.$$

For  $u \geq 1$ , let

$$g(u) = \left( \frac{3 \log f_A(x)}{u} \right)^u.$$

Then for  $1 \leq u < \frac{3}{e} \log f_A(x)$ , the function  $g(u)$  is increasing since

$$g'(u) = g(u) \log \frac{3 \log f_A(x)}{eu} > 0$$

and for large  $x$ , we have

$$g(1) = 3 \log f_A(x) < \frac{f_A(x)}{(\log \log x)^2}$$

and

$$g\left(\frac{3}{e} \log f_A(x)\right) = (f_A(x))^{3/e} > \frac{f_A(x)}{(\log \log x)^2}.$$

Thus there exists a uniquely determined real number  $t$  such that

$$(23) \quad 1 < t < \frac{3}{e} \log f_A(x)$$

and

$$(24) \quad g(t) = \left( \frac{3 \log f_A(x)}{t} \right)^t = \frac{f_A(x)}{(\log \log x)^2}.$$

We need a lower bound for this number  $t$ . By (1), we have

$$(25) \quad g\left(\frac{1}{2} \log f_A(x)\right) = (6^{1/2})^{\log f_A(x)} = (f_A(x))^{(1/2) \log 6} \\ < (f_A(x))^{9/10} = \frac{f_A(x)}{(f_A(x))^{1/10}} < \frac{f_A(x)}{(\log \log x)^2}.$$

(23), (24) and (25) imply that

$$(26) \quad \frac{1}{2} \log f_A(x) < t$$

(since  $g(u)$  is increasing for  $1 < u < \frac{3}{e} \log f_A(x)$ ).

Let us write

$$z_j = \max\{x^{1/2^j+1} \omega f_A(x), (\log x)^3\}.$$

Let  $A_j^*$  denote the set of the integers  $a$  such that  $a \in A'_j$  and

$$\omega(a, z_j, x^{1/2^j}) > t.$$

Now we are going to give an upper estimate for

$$\sum_{\substack{a \in A''_j \\ a \notin A_j^*}} \frac{1}{a} = \sum_{\substack{a \in A''_j \\ \omega(a, z_j, x^{1/2^j}) \leq t}} \frac{1}{a}.$$

If  $a \in A''_j$  and  $\omega(a, z_j, x^{1/2^j}) \leq t$  then by the definition of  $A''_j$ , we have

$$x^{1-1/2^j-1} = x_{j-1} \leq a < x_j = x^{1-1/2^j}$$

and  $a$  can be written in the form

$$a = u p_1^{\alpha_1} \dots p_m^{\alpha_m} v$$

where  $P(u) \leq (\log x)^3$ ,  $z_j < p_1 < \dots < p_m \leq x^{1/2^j}$ ,  $m \leq \omega(a, z_j, x^{1/2^j}) \leq t$  and  $p(v) > x^{1/2^j}$ .

Thus by Lemma 1, we have

$$(27) \quad \sum_{\substack{a \in A''_j \\ a \notin A_j^*}} \frac{1}{a} \leq \sum_{P(u) \leq (\log x)^3} \frac{1}{u} \left\{ \sum_{0 \leq m \leq t} \sum_{z_j < p_1 < \dots < p_m \leq x^{1/2^j}} \frac{1}{\prod_{i=0}^m p_i^{\alpha_i}} \times \right. \\ \left. \times \left( \sum_{\substack{p(v) > x^{1/2^j} \\ x_{j-1} < u p_1^{\alpha_1} \dots p_m^{\alpha_m} v < x_j = x_{j-1} x^{1/2^j}}} \frac{1}{v} \right) \right\}$$

$$\begin{aligned} &< c_2 \left( \sum_{P(u) \leq (\log x)^3} \frac{1}{u} \right) \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} \left( \sum_{z_j < p \leq x^{1/2^j}} \sum_{a=1}^{+\infty} \frac{1}{p^a} \right)^m \right) \\ &= c_2 \left( \prod_{p \leq (\log x)^3} \sum_{a=0}^{+\infty} \frac{1}{p^a} \right) \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} \left( \sum_{z_j < p \leq x^{1/2^j}} \sum_{a=1}^{+\infty} \frac{1}{p^a} \right)^m \right) \\ &= c_2 \left( \prod_{p \leq (\log x)^3} \frac{1}{1-1/p} \right) \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} \left( \sum_{z_j < p \leq x^{1/2^j}} \sum_{a=1}^{+\infty} \frac{1}{p^a} \right)^m \right). \end{aligned}$$

It is well-known that

$$\prod_{p \leq y} \frac{1}{1-1/p} < c_8 \log y$$

and

$$\sum_{p \leq y} \sum_{a=1}^{+\infty} \frac{1}{p^a} = \log \log y + c_9 + o(1).$$

Thus with respect to (1), (23), (24) and (26), and by using the Stirling-formula, we obtain from (27) that for  $x > X_1(\omega)$ ,

$$\begin{aligned} (28) \quad \sum_{\substack{a \in A_j^* \\ a \notin A_j}} \frac{1}{a} &< c_{10} \log((\log x)^3) \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} (\log \log x^{1/2^j} - \log \log z_j + c_{11})^m \right) \\ &= c_{10} \log \log x \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} \left( \log \frac{\log x^{1/2^j}}{\log z_j} + c_{11} \right)^m \right) \\ &\leq c_{10} \log \log x \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} \left( \log \frac{\log x^{1/2^j}}{\log x^{1/2^j+1} \omega f_A(x)} + c_{11} \right)^m \right) \\ &= c_{10} \log \log x \left( 1 + \sum_{m=1}^{[t]} \frac{1}{m!} (\log 2 \omega f_A(x) + c_{11})^m \right) \\ &< c_{12} t \log \log x \frac{(\log \omega f_A(x) + c_{13})^t}{t!} \\ &< c_{14} t^{1/2} \log \log x \left( \frac{e(\log \omega f_A(x) + c_{13})}{t} \right)^t \\ &< \frac{1}{16} \log \log x \left( \frac{3 \log f_A(x)}{t} \right)^t = \frac{1}{16} \log \log x \frac{f_A(x)}{(\log \log x)^3} \\ &= \frac{1}{16} \frac{f_A(x)}{\log \log x}. \end{aligned}$$

(22) and (28) yield that

$$\begin{aligned} (29) \quad f_{A_j^*}(x) &= \sum_{a \in A_j^*} \frac{1}{a} = \sum_{a \in A_j} \frac{1}{a} - \sum_{\substack{a \in A_j \\ a \notin A_j^*}} \frac{1}{a} \\ &> \frac{f_A(x)}{8 \log \log x} - \frac{f_A(x)}{16 \log \log x} = \frac{f_A(x)}{16 \log \log x}. \end{aligned}$$

Let  $S$  denote the set of the integers  $n$  such that  $n \leq x$  and  $n$  can be written in the form

$$(30) \quad n = au \quad \text{where} \quad a \in A_j^* \quad \text{and} \quad \omega(u, z_j, x^{1/2^j}) > \frac{18}{19} \log f_A(x).$$

For fixed  $n \in S$ , let  $\varphi(n)$  denote the number of representations of  $n$  in the form (30).

Then we have

$$\begin{aligned} (31) \quad \sum_{n \leq x} \varphi(n) &= \sum_{a \in A_j^*} \sum_{\substack{au \leq x \\ \omega(u, z_j, x^{1/2^j}) > \frac{18}{19} \log f_A(x)}} 1 \\ &= \sum_{a \in A_j^*} \left( \sum_{u \leq x/a} 1 - \sum_{\substack{u \leq x/a \\ \omega(u, z_j, x^{1/2^j}) \leq \frac{18}{19} \log f_A(x)}} 1 \right). \end{aligned}$$

In order to estimate the last sum, we use Lemma 2 with  $z_j, x^{1/2^j}, x/a$  and  $1/400$  in place of  $y, z, v$  and  $\alpha$ , respectively. Then  $1 \leq y$  and  $2y < z$  hold trivially by the definition of  $z_j$  (and by  $j \leq R$ ), and also  $z \leq v$  holds by

$$z = x^{1/2^j} = \frac{x}{x^{1-1/2^j}} = \frac{x}{x_j} < \frac{x}{a} = v$$

(since we have  $a \in A_j^*$  and thus  $a < x_j$ ). Thus Lemma 2 can be applied, and we obtain with respect to (1) and the definition of  $z_j$  that for large  $x$

$$\begin{aligned} (32) \quad \sum_{\substack{n \leq x/a \\ \omega(u, z_j, x^{1/2^j}) \leq \frac{399}{400} \\ z_j < p \leq x^{1/2^j} p}} 1 &< c_3 \frac{x}{a} \exp \left( Q \left( -\frac{1}{400} \right) \log \frac{\log x^{1/2^j}}{\log z_j} \right) \\ &< c_3 \frac{x}{a} \exp \left( -3 \cdot 10^{-6} \log \frac{\log x^{1/2^j}}{\log x^{1/2^j+1} \omega f_A(x)} \right) \\ &= c_3 \frac{x}{a} \exp \left( -3 \cdot 10^{-6} \log 2 \omega f_A(x) \right) < \frac{1}{3} \frac{x}{a}. \end{aligned}$$

Furthermore, by (6) we have

$$(33) \quad \frac{399}{400} \sum_{z_j < p \leq x^{1/2^j}} \frac{1}{p} > \frac{399}{400} \left( \log \frac{\log x^{1/2^j}}{\log z_j} - c_{15} \right).$$

By (8), we have

$$(34) \quad \frac{f_A(x)}{6} \leq \log x^{1/2^R} \leq \log x^{1/2^j}.$$

We obtain from (1) and (34) that

$$(35) \quad \begin{aligned} \log z_j &= \log \max \{ (\log x)^3, x^{1/2^j + 1 \omega f_A(x)} \} \\ &= \max \left\{ 3 \log \log x, \frac{\log x^{1/2^j}}{2 \omega f_A(x)} \right\} \\ &= \max \left\{ \log x^{1/2^j} \frac{3 \log \log x}{\log x^{1/2^j}}, \frac{\log x^{1/2^j}}{2 \omega f_A(x)} \right\} \\ &\leq \max \left\{ \log x^{1/2^j} \frac{3 \log \log x}{f_A(x)/6}, \frac{\log x^{1/2^j}}{2 \omega f_A(x)} \right\} \\ &\leq \max \left\{ \log x^{1/2^j} \frac{3 (f_A(x))^{1/20}}{f_A(x)/6}, \frac{\log x^{1/2^j}}{2 \omega f_A(x)} \right\} \\ &= \frac{18 \log x^{1/2^j}}{(f_A(x))^{19/20}}. \end{aligned}$$

(33) and (35) yield for large  $x$  that

$$(36) \quad \begin{aligned} \frac{399}{400} \sum_{z_j < p \leq x^{1/2^j}} \frac{1}{p} &> \frac{399}{400} \left( \log \frac{\log x^{1/2^j}}{\log z_j} - c_{15} \right) \\ &\geq \frac{399}{400} \left( \log \frac{(f_A(x))^{19/20}}{18} - c_{15} \right) > \frac{18}{19} \log f_A(x) \end{aligned}$$

hence

$$(37) \quad \sum_{\substack{u \leq x/a \\ \omega(u, z_j, x^{1/2^j}) < \frac{18}{19} \log f_A(x)}} 1 \leq \sum_{\substack{u \leq x/a \\ \omega(u, z_j, x^{1/2^j}) \leq \frac{399}{400} \sum_{z_j < p \leq x^{1/2^j} p}} 1.$$

(32) and (37) yield that

$$\sum_{\substack{u \leq x/a \\ \omega(u, z_j, x^{1/2^j}) > \frac{18}{19} \log f_A(x)}} 1 < \frac{1}{3} \frac{x}{a}.$$

Thus we obtain from (29) and (31) that

$$(38) \quad \begin{aligned} \sum_{n \leq x} \varphi(n) &\geq \sum_{a \in A_j^*} \left( \left[ \frac{x}{a} \right] - \frac{1}{3} \frac{x}{a} \right) > \sum_{a \in A_j^*} \frac{x}{2a} \\ &= \frac{x}{2} f_{A_j^*}(x) > \frac{x f_A(x)}{32 \log \log x}. \end{aligned}$$

Now we are going to give an upper estimate for  $\sum_{n \leq x} \varphi(n)$ . Obviously, for  $n \leq x$  we have  $\varphi(n) \leq d_A(n) \leq D_A(x)$  hence

$$(39) \quad \sum_{n \leq x} \varphi(n) = \sum_{n \in S} \varphi(n) \leq \sum_{n \in S} D_A(x) = |S| D_A(x).$$

Thus in order to obtain an upper bound for  $\sum_{n \leq x} \varphi(n)$ , we have to estimate  $|S|$ .

If  $n \in S$  then by (26), (30) and the definition of the set  $A_j^*$ , we have  $\omega(n, z_j, x^{1/2^j}) = \omega(au, z_j, x^{1/2^j}) = \omega(a, z_j, x^{1/2^j}) + \omega(u, z_j, x^{1/2^j})$

$$> t + \frac{18}{19} \log f_A(x) > \frac{1}{2} \log f_A(x) + \frac{18}{19} \log f_A(x) = \frac{55}{38} \log f_A(x)$$

hence

$$(40) \quad |S| \leq \sum_{\substack{n \leq x \\ \omega(n, z_j, x^{1/2^j}) > \frac{55}{38} \log f_A(x)}} 1.$$

In order to estimate this sum, we use Lemma 3 with  $z_j, x^{1/2^j}, x, 17/37$  and  $9/10$  in place of  $y, z, v, a$  and  $\beta$ , respectively. ( $1 \leq y, 2y < z \leq v$  and  $0 < a \leq \beta < 1$  hold trivially with respect to the definition of  $z_j$ .) We obtain with respect to (8) and the definition of  $z_j$  that for  $x > X_2(\omega)$ ,

$$(41) \quad \begin{aligned} &\sum_{\substack{n \leq x \\ \omega(n, z_j, x^{1/2^j}) > \frac{54}{37} \sum_{z_j < p \leq x^{1/2^j} p} \frac{1}{p}}} 1 \\ &< c_6 \omega \left( \sum_{z_j < p \leq x^{1/2^j} p} \frac{1}{p} \right)^{-1/2} \exp \left( Q \left( \frac{17}{37} \right) \log \frac{\log x^{1/2^j}}{\log z_j} \right) \\ &< c_6 x \exp \left( - \frac{91}{1000} \log \frac{\log x^{1/2^j}}{\log x^{1/2^j + 1 \omega f_A(x)}} \right) \\ &< c_6 x \exp \left( - \frac{91}{1000} \log \omega f_A(x) \right) < x \exp \left( - \frac{9}{100} \log f_A(x) \right) = x (f_A(x))^{-9/100}. \end{aligned}$$

Furthermore, with respect to (6) and the definition of  $z_j$  we have

$$\begin{aligned} \frac{54}{37} \sum_{z_j < p \leq x^{1/2^j}} \frac{1}{p} &\leq \frac{54}{37} \sum_{x^{1/2^{j+1}} \omega_{f_A}(x) < p \leq x^{1/2^j}} \frac{1}{p} \\ &= \frac{54}{37} \left( \sum_{p \leq x^{1/2^j}} \frac{1}{p} - \sum_{p \leq x^{1/2^{j+1}} \omega_{f_A}(x)} \frac{1}{p} \right) \\ &< \frac{54}{37} (\log \log x^{1/2^j} - \log \log x^{1/2^{j+1}} \omega_{f_A}(x) + c_{16}) \\ &= \frac{54}{37} (\log 2 \omega_{f_A}(x) + c_{16}) < \frac{55}{38} \log f_A(x) \end{aligned}$$

and thus

$$(42) \quad \sum_{\substack{n \leq x \\ \omega(n, z_j, x^{1/2^j}) > \frac{55}{38} \log f_A(x)}} 1 \leq \sum_{\substack{n \leq x \\ \omega(n, z_j, x^{1/2^j}) > \frac{54}{37} \sum_{z_j < p \leq x^{1/2^j}} \frac{1}{p}}} 1.$$

(40), (41) and (42) yield that

$$(43) \quad |S| < x (f_A(x))^{-9/100}.$$

Finally, we obtain from (38), (39) and (43) that

$$\frac{x f_A(x)}{32 \log \log x} < \sum_{n \leq x} \varphi(n) \leq |S| D_A(x) < x (f_A(x))^{-9/100} D_A(x)$$

hence with respect to (1),

$$\begin{aligned} D_A(x) &> f_A(x) \frac{(f_A(x))^{9/100}}{32 \log \log x} > f_A(x) \frac{(f_A(x))^{9/100}}{32 (f_A(x))^{1/20}} \\ &= f_A(x) \cdot \frac{1}{32} (f_A(x))^{1/25} > \omega f_A(x) \end{aligned}$$

for  $x > X_3(\omega)$ . Thus (2) holds also in Case 2 and this completes the proof of Theorem 1.

4. By using the same method, we can show that Theorem 1 is true also with  $(\log \log x)^{2+\varepsilon}$  in place of  $(\log \log x)^{20}$  on the right-hand side of (1). In fact, in order to prove this, the only non-trivial modifications are that  $t$  must be defined as  $t = \eta \log f_A(x)$  where  $\eta = \eta(\varepsilon) (> 0)$  is sufficiently small in terms of  $\varepsilon$ , and in (30), the condition  $\omega(u, z_j, x^{1/2^j}) > \frac{18}{19} \log f_A(x)$  must be replaced by  $\omega(u, z_j, x^{1/2^j}) > K \log f_A(x)$  where  $K = K(\varepsilon)$  is sufficiently large in terms of  $\varepsilon$ . Furthermore, then Lemmas 2

and 3 must be replaced by lower and upper estimates for

$$\sum_{\substack{n \leq x \\ \omega(n, y, z) \geq L}} \frac{1}{p}$$

where  $L$  is arbitrary large but fixed. Such estimates could be deduced by the methods used by K. K. Norton in [6]. (Norton's estimates cannot be used in the original form since the error terms in his lower and upper estimates depend implicitly on the set  $E$  of the prime numbers whose multiples we investigate. Thus in our case, these results would yield lower and upper bounds depending implicitly on  $\{p: y < p \leq z\}$ , i.e., on  $y$  and  $z$ , instead of the explicit estimates needed by us.)

On the other hand, we guess that also the exponent  $2+\varepsilon$  could be improved, and, perhaps, Theorem 1 is true also with  $(\log \log x)^{1+\varepsilon}$  or even  $c_{17}(\omega) \log \log x$  on the right-hand side of (1). This is the reason of that that we preferred to work out the slightly weaker estimate given in Theorem 1 whose proof is much simpler.

5. One may expect that if we know that  $f_A(y)$  is large for all  $y \leq x$  then Theorem 1 can be sharpened in the sense that the lower bound given for  $f_A(x)$  in (1) (for fixed  $x$ ) can be replaced by a much smaller lower bound for  $f_A(y)$  (for all  $y$ ). In fact, we show in this section that

**THEOREM 2.** For all  $\omega > 0$ , there exists a real number  $X_4 = X_4(\omega)$  such that if  $x > X_4$  and writing  $y = \exp\left(\frac{\log x}{(\log \log x)^{21}}\right)$ , we have

$$(44) \quad f_A(y) > 22 \log \log \log y,$$

then

$$(45) \quad D_A(x) > \omega f_A(x).$$

Furthermore, we show that Theorem 2 is best possible except the value of the constant factor on the right of (44):

**THEOREM 3.** There exist positive constants  $c_{18}, c_{19}, X_5$  and an infinite sequence  $A$  such that

$$(46) \quad f_A(x) > c_{18} \log \log \log x \quad \text{for all } x > X_5$$

and

$$(47) \quad \liminf_{x \rightarrow +\infty} \frac{D_A(x)}{f_A(x)} < c_{19}.$$

In order to prove Theorem 2, we need the following lemma:

**LEMMA 4.** If  $x > 1, t \geq 1$  and  $A$  is an arbitrary sequence of positive integers such that

$$(48) \quad D_A(x) \leq t$$

then we have

$$N_A(x^{1/(t+1)}) \leq t.$$

Proof of Lemma 4. Assume indirectly that

$$N_A(x^{1/(t+1)}) > t,$$

i.e.,

$$N_A(x^{1/(t+1)}) \geq [t]+1.$$

Then there exist integers  $a_1, a_2, \dots, a_{[t]+1}$  such that  $a_1 \in A, a_2 \in A, \dots, a_{[t]+1} \in A$  and

$$(49) \quad a_1 < a_2 < \dots < a_{[t]+1} \leq x^{1/(t+1)}.$$

Put  $u = a_1 a_2 \dots a_{[t]+1}$ . Then  $a_i | u$  for  $1 \leq i \leq [t]+1$  and thus

$$(50) \quad d_A(u) \geq [t]+1 > t.$$

On the other hand, by (49) we have

$$(51) \quad u = a_1 a_2 \dots a_{[t]+1} \leq (x^{1/(t+1)})^{[t]+1} \leq (x^{1/(t+1)})^{t+1} = x.$$

(50) and (51) imply that

$$D_A(x) > t$$

in contradiction with (48) which completes the proof of Lemma 4.

Proof of Theorem 2. We have to distinguish two cases.

Case 1. Let

$$f_A(x) > (\log \log x)^{20}.$$

Then for  $x > X_6(\omega)$ , (45) holds by Theorem 1.

Case 2. Let

$$(52) \quad f_A(x) \leq (\log \log x)^{20}.$$

Assume indirectly that

$$(53) \quad D_A(x) \leq \omega f_A(x).$$

Then by using Lemma 4 with  $t = \omega f_A(x)$ , we obtain that

$$(54) \quad N_A(x^{1/(t+1)}) = N_A(x^{1/(\omega f_A(x)+1)}) \leq t = \omega f_A(x).$$

Put  $M = N_A(x^{1/(\omega f_A(x)+1)})$  and let  $a_1 < a_2 < \dots < a_M$  denote the  $a$ 's not exceeding  $x^{1/(\omega f_A(x)+1)}$ . Then by (52) and (54), we have

$$(55) \quad \begin{aligned} f_A(x^{1/(\omega f_A(x)+1)}) &= \sum_{i=1}^M \frac{1}{a_i} \leq \sum_{i=1}^M \frac{1}{i} < \log M + c_{20} \\ &\leq \log \omega f_A(x) + c_{20} \leq \log \omega (\log \log x)^{20} + c_{20} \\ &< 21 \log \log \log x. \end{aligned}$$

On the other hand, by (51) we have

$$\begin{aligned} x^{1/(\omega f_A(x)+1)} &= \exp\left(\frac{\log x}{\omega f_A(x)+1}\right) \geq \exp\left(\frac{\log x}{\omega (\log \log x)^{20}+1}\right) \\ &\geq \exp\left(\frac{\log x}{(\log \log x)^{21}}\right) = y. \end{aligned}$$

Thus (44) yields that

$$\begin{aligned} f_A(x^{1/(\omega f_A(x)+1)}) &\geq f_A(y) > 22 \log \log \log y \\ &= 22 \log \log \frac{\log x}{(\log \log x)^{21}} > 21 \log \log \log x \end{aligned}$$

in contradiction with (55) which completes the proof of Theorem 2.

Proof of Theorem 3. In the proof of Theorem 1 in [1], for  $x \geq X_7$  we constructed a sequence  $B(x)$  such that

$$(56) \quad f_{B(x)}(x) > c_{21} \log \log x$$

and

$$(57) \quad D_{B(x)}(x) < 2 \log \log x.$$

Let us define the infinite sequence  $x_1 < x_2 < \dots$  by the following recursion: let

$$x_1 = X_7 \quad \text{and} \quad x_k = \exp(\exp(\exp(x_{k-1}))).$$

For  $x > 1$ , let

$$E(x) = \{n: \sqrt{x} < n \leq x\}.$$

Finally, let

$$A = \bigcup_{k=1}^{+\infty} B(x_k) \cup E(\log \log x_k).$$

We are going to show that this sequence  $A$  satisfies both (46) and (47).

First we prove (46). Assume that  $x > X_7$ . Then there exists a uniquely determined positive integer  $k$  ( $\geq 2$ ) such that  $x_{k-1} < x \leq x_k$ . Then either

$$(58) \quad x_{k-1} < x \leq \exp(x_{k-1}) = \log \log x_k$$

or

$$(59) \quad \exp(x_{k-1}) = \log \log x_k < x \leq x_k$$

holds. If (58) holds, then by (56) we have

$$f_A(x) \geq f_A(x_{k-1}) \geq f_{B(x_{k-1})}(x_{k-1}) > c_{21} \log \log x_{k-1} \geq c_{21} \log \log \log x$$



while if (59) holds then

$$f_A(x) \geq E(\log \log x_k) = \sum_{(\log \log x_k)^{1/2} < n \leq \log \log x_k} \frac{1}{n} \\ > \frac{1}{3} \log \log \log x_k \geq \frac{1}{3} \log \log \log x.$$

Thus in fact, (46) holds in both cases.

In order to prove that also (47) holds, it is sufficient to show that for  $k = 1, 2, \dots$ , we have

$$(60) \quad \frac{D_A(x_k)}{f_A(x_k)} < c_{22}.$$

If  $u \leq x_k$  then by (57) we have

$$d_A(u) = \sum_{\substack{a|u \\ a \in A}} 1 = \sum_{\substack{a \leq \log \log x_k \\ a|u, a \in A}} 1 + \sum_{\substack{\log \log x_k < a \\ a|u, a \in A}} 1 \\ = \sum_{\substack{a \leq \log \log x_k \\ a|u, a \in A}} 1 + \sum_{\substack{a|u \\ a \in B(x_k)}} 1 = \sum_{\substack{a \leq \log \log x_k \\ a|u, a \in A}} 1 + d_{B(x_k)}(u) \\ \leq \log \log x_k + D_{B(x_k)}(u) < 3 \log \log x_k$$

hence

$$(61) \quad D_A(x_k) < 3 \log \log x_k.$$

Furthermore, by (56), we have

$$(62) \quad f_A(x_k) = \sum_{\substack{a \leq x_k \\ a \in A}} \frac{1}{a} \geq \sum_{a \in B(x_k)} \frac{1}{a} = f_{B(x_k)}(x_k) > c_{21} \log \log x_k.$$

(61) and (62) yield (60) and the proof of Theorem 3 is completed.

#### References

- [1] P. Erdős and A. Sárközy, *Some asymptotic formulas on generalized divisor functions*, I, Studies in Pure Mathematics to the Memory of Paul Turán, Akadémiai Kiadó, Budapest, to appear.
- [2] P. Erdős and A. Sárközy, *Some asymptotic formulas on generalized divisor functions*, II, J. Number Theory, to appear.
- [3] G. Halász, *Remarks to my paper "On the distribution of additive and the mean values of multiplicative arithmetic functions"*, Acta Math. Acad. Sci. Hungar. 23 (1972), pp. 425-432.

- [4] R. R. Hall, *On a conjecture of Erdős and Sárközy*, Bull. London Math. Soc. 12 (1980), pp. 21-24.
- [5] K. K. Norton, *On the number of restricted prime factors of an integer*, I, Illinois J. Math. 20 (1976), pp. 681-705.
- [6] K. K. Norton, *On the number of restricted prime factors of an integer*, III, to appear.
- [7] K. Prachar, *Primzahlverteilung*, Springer Verlag, Berlin 1957.

Received on 9. 1. 1981

(1238)