

## Natural transformations of higher order cotangent bundle functors

by JAN KUREK (Lublin)

**Abstract.** We determine all natural transformations of the  $r$ th order cotangent bundle functor  $T^{r*}$  into  $T^{s*}$  in the following cases:  $r = s$ ,  $r < s$ ,  $r > s$ . We deduce that all natural transformations of  $T^{r*}$  into itself form an  $r$ -parameter family linearly generated by the  $p$ th power transformations with  $p = 1, \dots, r$ .

Using general methods developed in [2]–[5], we deduce that all natural transformations of the  $r$ th order cotangent bundle functor  $T^{r*}$  into itself form an  $r$ -parameter family generated by the  $p$ th power transformations  $A_p^{r,r}$  with  $p = 1, \dots, r$ .

Then we deduce that all natural transformations of  $T^{r*}$  into  $T^{(r+q)*}$  form an  $r$ -parameter family generated by the generalized  $p$ th power transformations  $A_p^{r,r+q}$  with  $p = q + 1, \dots, q + r$ .

Moreover, we deduce that all natural transformations of  $T^{r*}$  into  $T^{(r-q)*}$  form an  $(r - q)$ -parameter family generated by the generalized  $p$ th power transformations  $A_p^{r,r-q}$  with  $p = 1, \dots, r - q$ .

The author is grateful to Professor I. Kolář for suggesting the problem and for valuable remarks and useful discussions.

1. Let  $M$  be a smooth  $n$ -dimensional manifold. Let  $T^{r*}M = J^r(M, \mathbb{R})_0$  be the space of all  $r$ -jets  $j_x^r f$  of smooth functions  $f : M \rightarrow \mathbb{R}$  with source at  $x \in M$  and target at  $0 \in \mathbb{R}$ . The fibre bundle  $\pi_M : T^{r*}M \rightarrow M$  with source  $r$ -jet projection  $\pi_M : j_x^r f \mapsto x$  has a canonical structure of a vector bundle with

$$(1.1) \quad j_x^r f + j_x^r g = j_x^r (f + g), \quad k \cdot j_x^r f = j_x^r (k \cdot f)$$

for  $x \in M$  and  $k \in \mathbb{R}$  [1].

---

1991 *Mathematics Subject Classification*: Primary 58A20.

*Key words and phrases*: higher order cotangent bundle, jet,  $p$ th power transformation.

The vector bundle  $\pi_M : T^{r*}M \rightarrow M$  is called the  $r$ -th cotangent bundle over  $M$ .

Every local diffeomorphism  $\varphi : M \rightarrow N$  is extended to a vector bundle morphism  $T^{r*}\varphi : T^{r*}M \rightarrow T^{r*}N$ ,  $j_x^r f \mapsto j_{\varphi(x)}^r (f \circ \varphi^{-1})$ , where  $\varphi^{-1}$  is locally defined. Hence, the  $r$ th cotangent bundle functor  $T^{r*}$  is defined on the category  $\mathcal{M}f_n$  of smooth  $n$ -dimensional manifolds with local diffeomorphisms as morphisms and has values in the category  $\mathcal{VB}$  of vector bundles.

If  $(x^i)$  are local coordinates on  $M$ , then we have the induced fibre coordinates  $(u_i, u_{i_1 i_2}, \dots, u_{i_1 \dots i_r})$  on  $T^{r*}M$  (symmetric in all indices):

$$(1.2) \quad \begin{aligned} u_i(j_x^r f) &= \left. \frac{\partial f}{\partial x^i} \right|_{(x)}, & u_{i_1 i_2}(j_x^r f) &= \left. \frac{\partial^2 f}{\partial x^{i_1} \partial x^{i_2}} \right|_{(x)}, \dots, \\ u_{i_1 \dots i_r}(j_x^r f) &= \left. \frac{\partial^r f}{\partial x^{i_1} \dots \partial x^{i_r}} \right|_{(x)}. \end{aligned}$$

Since the functor  $T^{r*}$  takes values in the category  $\mathcal{VB}$  of vector bundles, we may define natural transformations  $A_p^{r,r}$  of  $T^{r*}$  into itself for  $p = 1, \dots, r$ .

DEFINITION 1. The natural transformation  $A_p^{r,r}$  of the  $r$ th cotangent bundle functor  $T^{r*}$  into itself defined by

$$(1.3) \quad A_p^{r,r} : j_x^r f \mapsto j_x^r (f)^p,$$

where  $(f)^p$  denotes the  $p$ th power of  $f$ , is called the  $p$ -th power transformation.

DEFINITION 2. The natural transformation  $A_p^{r,s}$  of  $T^{r*}$  into  $T^{s*}$  defined by

$$(1.4) \quad A_p^{r,s} : j_x^r f \mapsto j_x^s (f)^p$$

is called the generalized  $p$ -th power transformation.

We note that in the case  $s = r + q$  this definition is correct only for  $p = q + 1, \dots, q + r$ .

DEFINITION 3. The natural transformation  $P^{r,r-q}$  of  $T^{r*}$  into  $T^{(r-q)*}$  defined by

$$(1.5) \quad P^{r,r-q} : j_x^r f \mapsto j_x^{r-q} f$$

is called a projection.

Note that

$$(1.6) \quad A_p^{r,r-q} = A_p^{r-q,r-q} \circ P^{r,r-q} \quad \text{for } p = 1, \dots, r - q.$$

**2.** In this part we determine, by induction on  $r$ , all natural transformations of  $T^{r*}$  into itself.

THEOREM 1. All natural transformations  $A : T^{r*} \rightarrow T^{r*}$  form the  $r$ -parameter family

$$(2.1) \quad A = k_1 A_1^{r,r} + k_2 A_2^{r,r} + \dots + k_r A_r^{r,r}$$

with any real parameters  $k_1, k_2, \dots, k_r \in \mathbb{R}$ .

PROOF. The functor  $T^{r*}$  is defined on the category  $\mathcal{M}f_n$  of  $n$ -dimensional smooth manifolds with local diffeomorphisms as morphisms and is of order  $r$ . Thus, its standard fibre  $S = (T^{r*}\mathbb{R}^n)_0$  is a  $G_n^r$ -space, where  $G_n^r$  is the group of all invertible  $r$ -jets from  $\mathbb{R}^n$  into  $\mathbb{R}^n$  with source and target at 0.

According to general standard methods [2]–[5], the natural transformations  $A : T^{r*} \rightarrow T^{r*}$  are in bijection with the  $G_n^r$ -equivariant maps  $f^{r,r} : (T^{r*}\mathbb{R}^n)_0 \rightarrow (T^{r*}\mathbb{R}^n)_0$  of the standard fibres.

Let  $\tilde{a} = a^{-1}$  denote the inverse element in  $G_n^r$  and let

$$(2.2) \quad t_{(i_1 \dots i_r)} = \frac{1}{r!} \sum_{\sigma \in S_r} t_{i_{\sigma(1)} \dots i_{\sigma(r)}}$$

denote the symmetrization of a tensor with components  $t_{i_1 \dots i_r}$ .

By (1.2) the action of an element  $(a_{j_1}^i, a_{j_1 j_2}^i, \dots, a_{j_1 \dots j_r}^i) \in G_n^r$  on  $(u_{i_1}, u_{i_1 i_2}, \dots, u_{i_1 \dots i_r}) \in (T^{r*}\mathbb{R}^n)_0$  is given by

$$(2.3) \quad \begin{aligned} \bar{u}_{i_1} &= u_{j_1} \tilde{a}_{i_1}^{j_1}, \quad \bar{u}_{i_1 i_2} = u_{j_1 j_2} \tilde{a}_{i_1}^{j_1} \tilde{a}_{i_2}^{j_2} + u_{j_1} \tilde{a}_{i_1 i_2}^{j_1}, \dots, \\ \bar{u}_{i_1 \dots i_r} &= u_{j_1 \dots j_r} \tilde{a}_{i_1}^{j_1} \dots \tilde{a}_{i_r}^{j_r} + u_{j_1 \dots j_{r-1}} \frac{r!}{(r-2)!} \tilde{a}_{(i_1}^{j_1} \dots \tilde{a}_{i_{r-2}}^{j_{r-2}} \tilde{a}_{i_{r-1} i_r}^{j_{r-1}}) + \dots \\ &+ u_{j_1 \dots j_{r-s}} \left[ \frac{r!}{(r-s-1)!(s+1)!} \tilde{a}_{(i_1}^{j_1} \dots \tilde{a}_{i_{r-s-1}}^{j_{r-s-1}} \tilde{a}_{i_{r-s} \dots i_r}^{j_{r-s}}) + \dots \right] \\ &+ u_{j_1 j_2} \left[ \frac{r!}{(r-1)!} \tilde{a}_{(i_1}^{j_1} \tilde{a}_{i_2 \dots i_r}^{j_2} + \dots \right] + u_{j_1} \tilde{a}_{i_1 \dots i_r}^{j_1}. \end{aligned}$$

I. Consider the case  $r = 2$ . The equivariance of a  $G_n^2$ -equivariant map  $f^{2,2} = (f_i, f_{ij})$  of  $(T^{2*}\mathbb{R}^n)_0$  into itself with respect to homotheties in  $G_n^2$ :  $\tilde{a}_j^i = k\delta_j^i$ ,  $\tilde{a}_{j_1 j_2}^i = 0$ , gives the homogeneity conditions

$$(2.4) \quad kf_i(u_i, u_{ij}) = f_i(ku_i, k^2 u_{ij}), \quad k^2 f_{ij}(u_i, u_{ij}) = f_{ij}(ku_i, k^2 u_{ij}).$$

By the homogeneous function theorem [2]–[5], we deduce that, first, the  $f_i$  are linear in  $u_i$  and independent of  $u_{ij}$ , and secondly, the  $f_{ij}$  are linear in  $u_{ij}$  and quadratic in  $u_i$ . Using the invariant tensor theorem for  $G_n^1$  [2]–[5], we obtain  $f^{2,2}$  in the form

$$(2.5) \quad f_i = k_1 u_i, \quad f_{ij} = k_2 u_i u_j + k_3 u_{ij}$$

with any real parameters  $k_1, k_2, k_3 \in \mathbb{R}$ .

The equivariance of  $f^{2,2}$  of the form (2.5) with respect to the kernel of the projection  $G_n^2 \rightarrow G_n^1 : \tilde{a}_j^i = \delta_j^i$  and  $\tilde{a}_{jk}^i$  arbitrary, gives

$$(2.6) \quad k_3 = k_1.$$

This proves our theorem for  $r = 2$ .

II. Suppose that the theorem holds for  $r - 1$  and the  $G_n^{r-1}$ -equivariant maps  $f^{r-1, r-1}$  of  $(T^{(r-1)*}\mathbb{R}^n)_0$  into itself define the  $(r-1)$ -parameter family  $\bar{A} = k_1 A_1^{r-1, r-1} + \dots + k_{r-1} A_{r-1}^{r-1, r-1}$  with any real parameters  $k_1, \dots, k_{r-1} \in \mathbb{R}$ .

Our aim is to obtain the general form of any  $G_n^r$ -equivariant map of  $(T^{r*}\mathbb{R}^n)_0$  into itself.

Let  $(u_1, u_2, \dots, u_r) := (u_{i_1}, u_{i_1 i_2}, \dots, u_{i_1 \dots i_r})$  denote the fibre coordinates on  $T^{r*}M$ . We assume that a  $G_n^r$ -equivariant map  $f^{r, r}$  is of the general form  $f^{r, r} = (f_1, \dots, f_{r-1}, f_r)$  and the given map  $f^{r-1, r-1}$  defines the first  $r - 1$  components  $(f_1, \dots, f_{r-1})$  of  $f^{r, r}$ .

Considering the equivariance of  $f^{r, r}$  with respect to the homotheties  $\tilde{a}_j^i = k\delta_j^i$ ,  $\tilde{a}_{j_1 j_2}^i = 0, \dots, \tilde{a}_{j_1 \dots j_r}^i = 0$  in  $G_n^r$ , for the  $r$ th component  $f_r$  we obtain the homogeneity condition

$$(2.7) \quad k^r f_r(u_1, u_2, \dots, u_r) = f_r(ku_1, k^2 u_2, \dots, k^r u_r).$$

By the homogeneous function theorem [2]–[5],  $f_r$  is of the general form

$$(2.8) \quad f_{i_1 \dots i_r} = h_r u_{i_1} \cdot \dots \cdot u_{i_r} + h_{r-1} u_{(i_1 \dots i_{r-2} i_{r-1} i_r)} + \dots + h_{2,1} u_{(i_1 i_2 \dots i_r)} + h_{2,2} u_{(i_1 i_2 i_3 \dots i_r)} + \dots + h_1 u_{i_1 \dots i_r}$$

with any real parameters  $h_1, h_{2,1}, h_{2,2}, \dots, h_{r-1}, h_r \in \mathbb{R}$ . The equivariance of  $f^{r, r}$  with respect to the kernel of the projection  $G_n^r \rightarrow G_n^{r-1} : \tilde{a}_j^i = \delta_j^i$ ,  $\tilde{a}_{j_1 j_2}^i = 0, \dots, \tilde{a}_{j_1 \dots j_{r-1}}^i = 0$  and  $\tilde{a}_{j_1 \dots j_r}^i$  arbitrary, gives

$$(2.9) \quad h_1 = k_1.$$

Thus, we obtain the 1st power transformation  $A_1^{r, r}$ .

Now, considering the equivariance of  $A - k_1 A_1^{r, r}$  with respect to the kernel of the projection  $G_n^{r-1} \rightarrow G_n^1 : \tilde{a}_j^i = \delta_j^i$  and  $\tilde{a}_{j_1 j_2}^i, \dots, \tilde{a}_{j_1 \dots j_{r-1}}^i$  arbitrary, we obtain

$$(2.10) \quad h_{2,1} = \frac{r!}{1!(r-1)!} k_2, \quad h_{2,2} = \frac{r!}{2!(r-2)!} k_2, \quad \dots$$

Thus, we obtain the 2nd power transformation  $A_2^{r, r}$ .

Then, considering the equivariance of  $A - k_1 A_1^{r, r} - k_2 A_2^{r, r}$  with respect to the kernel of the projection  $G_n^{r-2} \rightarrow G_n^1$ , we obtain the general form of the 3rd power transformation  $A_3^{r, r}$ . Continuing this procedure gives the next power transformations  $A_3^{r, r}, \dots, A_{r-2}^{r, r}$ . The equivariance of  $A - k_1 A_1^{r, r} - k_2 A_2^{r, r} - \dots - k_{r-2} A_{r-2}^{r, r}$  with respect to the kernel of the projection  $G_n^2 \rightarrow G_n^1$ :

$\tilde{a}_j^i = \delta_j^i$  and  $\tilde{a}_{jk}^i$  arbitrary, leads to the next relation

$$(2.11) \quad h_{r-1} = \frac{r!}{(r-2)!2!} k_{r-1}.$$

Thus, we obtain the  $(r-1)$ th power transformation  $A_{r-1}^{r,r}$ .

Finally, the  $G_n^r$ -equivariant map

$$(2.12) \quad A - k_1 A_1^{r,r} - k_2 A_2^{r,r} - \dots - k_{r-1} A_{r-1}^{r,r} = h_r A_r^{r,r}$$

is defined by the  $r$ th power transformation with any real parameter  $h_r \in \mathbb{R}$ . If we put  $h_r = k_r$ , this proves our theorem.

**3.** In this part we determine all natural transformations  $T^{r*} \rightarrow T^{s*}$  in two cases:  $r < s$  and  $r > s$ .

**THEOREM 2.** *All natural transformations  $A : T^{r*} \rightarrow T^{(r+q)*}$  form the  $r$ -parameter family*

$$(3.1) \quad A = k_{q+1} A_{q+1}^{r,r+q} + k_{q+2} A_{q+2}^{r,r+q} + \dots + k_{q+r} A_{q+r}^{r,r+q}$$

with any real parameters  $k_{q+1}, k_{q+2}, \dots, k_{q+r} \in \mathbb{R}$ .

**Proof.** We apply induction on  $q$ .

I. Consider the case  $q = 1$ . According to general standard methods [2]–[5], the natural transformations  $A : T^{r*} \rightarrow T^{(r+1)*}$  are in bijection with the  $G_n^{r+1}$ -equivariant maps of the standard fibres  $f^{r,r+1} : (T^{r*}\mathbb{R}^n)_0 \rightarrow (T^{(r+1)*}\mathbb{R}^n)_0$ .

Considering the equivariance of  $f^{r,r+1} = (f_1, \dots, f_r, f_{r+1})$  with respect to homotheties:  $\tilde{a}_j^i = k\delta_j^i$ ,  $\tilde{a}_{j_1 j_2}^i = 0$ ,  $\dots$ ,  $\tilde{a}_{j_1 \dots j_{r+1}}^i = 0$ , we obtain the homogeneity conditions

$$(3.2) \quad \begin{aligned} k f_1(u_1, u_2, \dots, u_r) &= f_1(ku_1, k^2 u_2, \dots, k^r u_r), \dots, \\ k^r f_r(u_1, u_2, \dots, u_r) &= f_r(ku_1, k^2 u_2, \dots, k^r u_r), \\ k^{r+1} f_{r+1}(u_1, u_2, \dots, u_r) &= f_{r+1}(ku_1, k^2 u_2, \dots, k^r u_r). \end{aligned}$$

Additionally, using the equivariance of  $f^{r,r} = (f_1, \dots, f_r)$  with respect to the kernel of the projection  $G_n^r \rightarrow G_n^1$ , we obtain, by Theorem 1, the  $r$ -parameter family of the form (2.1):  $\bar{A} = k_1 A_1^{r,r} + k_2 A_2^{r,r} + \dots + k_r A_r^{r,r}$  with any real parameters  $k_1, k_2, \dots, k_r \in \mathbb{R}$ .

Moreover, by the homogeneous function theorem and the invariant tensor theorem [2]–[5], we deduce that the  $(r+1)$ th component  $f_{r+1}$  is of the general form

$$(3.3) \quad \begin{aligned} f_{i_1 \dots i_{r+1}} &= l_{r+1} u_{i_1} u_{i_2} \dots u_{i_{r+1}} + l_r u_{(i_1 \dots i_{r-1} i_{r+1})} \\ &+ \dots + l_{2,1} u_{(i_1 i_2 \dots i_{r+1})} + l_{2,2} u_{(i_1 i_2 i_3 \dots i_{r+1})} + \dots \end{aligned}$$

with any real parameters  $l_{2,1}, l_{2,2}, \dots, l_r, l_{r+1} \in \mathbb{R}$ .

The equivariance of  $f^{r,r+1}$  with respect to the kernel of the projections  $G_n^{r+1} \rightarrow G_n^r$  and  $G_n^{r+1} \rightarrow G_n^1$  gives the relations

$$(3.4) \quad k_1 = 0,$$

$$(3.5) \quad l_{2,1} = \frac{(r+1)!}{r!1!} k_2, \quad l_{2,2} = \frac{(r+1)!}{(r-1)!2!} k_2, \dots, l_r = \frac{(r+1)!}{(r-1)!2!} k_r.$$

If we put  $l_{r+1} = k_{r+1}$ , this gives the  $r$ -parameter family  $f^{r,r+1} = (f^{r,r}, f_{r+1})$  of the form

$$(3.6) \quad A = k_2 A_2^{r,r+1} + \dots + k_{r+1} A_{r+1}^{r,r+1}$$

with any real parameters  $k_2, \dots, k_{r+1} \in \mathbb{R}$ .

II. Suppose that the theorem holds for  $q-1$  and the  $G_n^{r+q-1}$ -equivariant maps  $f^{r,r+q-1} : (T^{r*} \mathbb{R}^n)_0 \rightarrow (T^{(r+q-1)*} \mathbb{R}^n)_0$  define the  $r$ -parameter family

$$(3.7) \quad \bar{A} = k_q A_q^{r,r+q-1} + k_{q+1} A_{q+1}^{r,r+q-1} + \dots + k_{q+r-1} A_{q+r-1}^{r,r+q-1}$$

with any real parameters  $k_q, k_{q+1}, \dots, k_{q+r-1} \in \mathbb{R}$ .

Consider a  $G_n^{r+q}$ -equivariant map  $f^{r,r+q} : (T^{r*} \mathbb{R}^n)_0 \rightarrow (T^{(r+q)*} \mathbb{R}^n)_0$  of the form  $f^{r,r+q} = (f^{r,r+q-1}, f_{r+q})$ .

The equivariance of  $f^{r,r+q}$  with respect to the homotheties in  $G_n^{r+q}$ :  $\tilde{a}_j^i = k \delta_j^i$ ,  $\tilde{a}_{j_1 j_2}^i = 0, \dots, \tilde{a}_{j_1 \dots j_{r+q}}^i = 0$ , gives for the  $(r+q)$ th component  $f_{r+q}$  the homogeneity condition

$$(3.8) \quad k^{r+q} f_{r+q}(u_1, u_2, \dots, u_r) = f_{r+q}(k u_1, k^2 u_2, \dots, k^r u_r).$$

By the homogeneous function theorem and the invariant tensor theorem [2]–[5],  $f_{r+q}$  is of the form

$$(3.9) \quad f_{i_1 \dots i_{r+q}} = l_{r+q} u_{i_1} \dots u_{i_{r+q}} + l_{r+q-1} u_{(i_1 \dots u_{i_{r+q-2}} u_{i_{r+q-1} i_{r+q}})} \\ + \dots + l_{q+1} u_{(i_1 \dots u_{i_q} u_{i_{q+1} \dots i_{q+r}})}$$

with any real parameters  $l_{q+1}, \dots, l_{q+r-1}, l_{q+r} \in \mathbb{R}$ .

The equivariance of  $f^{r,r+q}$  with respect to the kernel of the projections  $G_n^{r+q} \rightarrow G_n^r$  and  $G_n^{r+q} \rightarrow G_n^1$  gives the relations

$$(3.10) \quad k_q = 0, \dots,$$

$$(3.11) \quad l_{q+1} = \frac{(q+r)!}{r!q!} k_{q+1}, \dots, l_{q+r-1} = \frac{(q+r)!}{(q+r-2)!2!} k_{q+r-1}.$$

If we put  $l_{q+r} = k_{q+r}$ , this proves our theorem.

**THEOREM 3.** *All natural transformations  $A : T^{r*} \rightarrow T^{(r-q)*}$  form the  $(r-q)$ -parameter family*

$$(3.12) \quad A = k_1 A_1^{r,r-q} + k_2 A_2^{r,r-q} + \dots + k_{r-q} A_{r-q}^{r,r-q}$$

with any real parameters  $k_1, k_2, \dots, k_{r-q} \in \mathbb{R}$ .

**Proof.** Applying the same general procedure, we conclude that each  $A : T^{r*} \rightarrow T^{(r-q)*}$  is the composition of the projection  $P^{r,r-q} : T^{r*} \rightarrow T^{(r-q)*}$

and a transformation  $\bar{A}$  of  $T^{(r-q)*}$  into itself:  $A = \bar{A} \circ P^{r,r-q}$ . This proves our theorem.

### References

- [1] J. Gancarzewicz, *Differential Geometry*, PWN, Warszawa 1987 (in Polish).
- [2] I. Kolář, *Some natural operators in differential geometry*, in: Proc. Conf. Diff. Geom. and its Applications, Brno 1986, Kluwer, Dordrecht 1987, 91–110.
- [3] I. Kolář and G. Vosmanská, *Natural transformations of higher order tangent bundles and jet spaces*, Časopis Pěst. Mat. 114 (2) (1989), 181–186.
- [4] I. Kolář, P. Michor and J. Slovák, *Natural Operations in Differential Geometry*, to appear.
- [5] J. Kurek, *On natural operators on sectorform fields*, Časopis Pěst. Mat. 115 (3) (1990), 232–239.

INSTITUTE OF MATHEMATICS  
MARIA CURIE-SKŁODOWSKA UNIVERSITY  
PL. M. CURIE-SKŁODOWSKIEJ 1  
20-031 LUBLIN, POLAND

*Reçu par la Rédaction le 20.5.1991*  
*Révisé le 10.2.1992 et 29.6.1992*