

LOCAL BEHAVIOUR OF OPERATORS

VLADIMÍR MÜLLER

*Institute of Mathematics, Academy of Sciences of the Czech Republic
Žitná 25, 115 67 Praha 1, Czech Republic*

Introduction. Let T be a bounded operator in a Banach space X and let $x \in X$. Denote by \mathcal{P} the set of all complex polynomials. We are going to study the following problem:

A. What can we say about the set $\{p(T)x : p \in \mathcal{P}\}$?

A weaker version of this problem is:

B. What can we say about the set $\{T^k x : k = 0, 1, \dots\}$?

The sets $\{T^k x : k = 0, 1, \dots\}$ for operators in Hilbert spaces were called “orbits” by Rolewicz [15] and intensively studied by Beauzamy [2].

Questions of type A or B appear naturally in many problems of operator theory.

EXAMPLES. 1. The local spectral radius $r(T, x)$ of an operator T at a point $x \in X$ can be defined by $r(T, x) = \limsup_{k \rightarrow \infty} \|T^k x\|^{1/k}$, i.e. it is a quantity defined in terms of B. The local spectral radius plays an important role in local spectral theory.

2. The local capacity (see below) can be considered as an analogue to the local spectral radius for the set of all polynomials.

3. The invariant subspace problem can also be easily reformulated by using the sets $\{p(T)x : p \in \mathcal{P}\}$: An operator T in X has no non-trivial invariant subspace if and only if $\{p(T)x : p \in \mathcal{P}\}$ is dense for all $x \in X$. Many positive results (e.g. results based on the Scott Brown technique) consist in finding $x \in X$ such that $\|p(T)x\| \geq 1$ for all polynomials p with $p(0) = 1$.

1991 *Mathematics Subject Classification*: 47A05, 47A11, 47A13.

The paper is in final form and no version of it will be published elsewhere.

The present paper is a survey of results obtained in [8]–[14]. The results show that for every Banach space X and every bounded linear operator T on X there exists $x \in X$ such that $\|p(T)x\|$ is big enough for all polynomials p .

I. Essential approximate point spectrum. Denote by $B(X)$ the algebra of all bounded operators in a Banach space X . Denote by \mathbb{C} and \mathbb{N} the set of all complex numbers and positive integers, respectively.

Let T be a bounded operator in a Banach space X . If x is an eigenvalue of T , $Tx = \lambda x$ for some complex λ , then $p(T)x = p(\lambda)x$ for every polynomial p so that we have a complete information about the set $\{p(T)x : p \in \mathcal{P}\}$. Unfortunately, operators in infinite-dimensional Banach spaces usually have no eigenvalues. The proper tool appears to be the notion of the essential approximate point spectrum of T .

Denote by $\sigma_e(T)$ the *essential spectrum* of $T \in B(X)$, i.e. the spectrum of $\varrho(T)$ in the Calkin algebra $B(X)/K(X)$, where $K(X)$ is the ideal of all compact operators in X and $\varrho : B(X) \rightarrow B(X)/K(X)$ is the canonical projection. Denote further by $\sigma_{\pi_e}(T)$ the *essential approximate point spectrum* of T , i.e. $\sigma_{\pi_e}(T)$ is the set of all complex λ such that

$$\inf\{\|(T - \lambda)x\| : x \in M, \|x\| = 1\} = 0$$

for every subspace $M \subset X$ with $\text{codim } M < \infty$.

It is easy to see that $\lambda \notin \sigma_{\pi_e}(T)$ if and only if $\dim \text{Ker}(T - \lambda) < \infty$ and $T - \lambda$ has closed range, i.e. if $T - \lambda$ is upper semi-Fredholm.

The terminology is not unified, the essential approximate point spectrum was studied under various names (see e.g. [1], [3], [7]).

By [7], $\sigma_{\pi_e}(T)$ contains the topological boundary of the essential spectrum, in particular it is always a non-empty compact subset of $\sigma_e(T)$.

We start with studying the elements $\lambda \in \sigma_{\pi_e}(T)$ for operators in Hilbert spaces. We show that, for given $k \in \mathbb{N}$, there always exists x such that the powers $T^i x$ ($0 \leq i \leq k$) are smaller and smaller and almost orthogonal to each other.

PROPOSITION 1. *Let T be an operator in a Hilbert space H such that $0 \in \sigma_{\pi_e}(T)$ and $\text{codim } \overline{TH} < \infty$. Let $k \in \mathbb{N}$ and $\varepsilon > 0$. Then there exists $x \in X$ with $\|x\| = 1$ such that*

- (1) $\|T^{i+1}x\| \leq \varepsilon \|T^i x\|$ ($i = 0, 1, \dots, k-1$),
- (2) $\|p(T)x\| \geq (1 - \varepsilon)|p(0)|$ ($p \in \mathcal{P}$, $\deg p \leq k$),
- (3) $|\langle T^i x, T^j x \rangle| \leq \varepsilon \|T^i x\| \cdot \|T^j x\|$ ($0 \leq i, j \leq k$, $i \neq j$).

Remark. The condition $\text{codim } \overline{TH} < \infty$ is only technical and rather weak. If this condition is not satisfied then 0 is an eigenvalue of T^* . In particular, T has a non-trivial invariant subspace, so that this case is not interesting (at least from the point of view of the invariant subspace problem).

The following lemma is an important tool for various constructions in Banach spaces. It enables one to generalize constructions in Hilbert spaces which use

the orthogonal complement of a finite-dimensional subspace to general Banach spaces.

LEMMA 2. *Let E be a finite-dimensional subspace of a Banach space X and let $\varepsilon > 0$. Then there exists a subspace $Y \subset X$ with $\text{codim } Y < \infty$ such that*

$$\|e + y\| \geq (1 - \varepsilon) \max\{\|e\|, \frac{1}{2}\|y\|\}$$

for every $e \in E$ and $y \in Y$.

By using Lemma 2 we can get an analogue to Proposition 1 for operators in Banach spaces.

PROPOSITION 3. *Let T be an operator in a Banach space X such that $0 \in \sigma_{\pi_e}(T)$ and $\text{codim } \overline{T X} < \infty$. Let $k \in \mathbb{N}$ and $\varepsilon > 0$. Then there exists $x \in X$ with $\|x\| = 1$ such that*

- (1) $\|T^{i+1}x\| \leq \varepsilon\|T^i x\|$ ($i = 0, 1, \dots, k - 1$),
- (2) $\|p(T)x\| \geq \frac{1}{2}(1 - \varepsilon)|p(0)|$ ($p \in \mathcal{P}$, $\text{deg } p \leq k$).

By an inductive construction which uses the previous proposition we can construct a point $x \in X$ (actually, a dense subset of X) such that $\|p(T)x\|$ is big enough for all polynomials p (see [11]).

THEOREM 4. *Let $T \in B(X)$, $\lambda \in \sigma_{\pi_e}(T)$. Let $\{a_k\}_{k=0}^\infty$ be a sequence of positive numbers with $\lim_{k \rightarrow \infty} a_k = 0$. Then there exists $x \in X$ such that*

$$\|p(T)x\| \geq a_{\text{deg } p} \cdot |p(\lambda)|$$

for every polynomial p .

THEOREM 5. *Let $T \in B(X)$, $\lambda \in \sigma_{\pi_e}(T)$, $x \in X$, $\varepsilon > 0$. Let $\{a_k\}_{k=0}^\infty$ be a sequence of positive numbers with $\lim_{k \rightarrow \infty} a_k = 0$. Then there exists $y \in X$ and a positive constant $C = C(\varepsilon)$ such that $\|y - x\| \leq \varepsilon$ and*

$$\|p(T)y\| \geq C a_{\text{deg } p} \cdot |p(\lambda)|$$

for every polynomial p .

REMARK. Let $T \in B(X)$, $\lambda \in \sigma_{\pi_e}(T)$ and suppose that there exist $y \in X$ and a constant $c > 0$ such that $\|p(T)y\| \geq c|p(\lambda)|$ for every polynomial p . Then either $(T - \lambda)y = 0$ or $M = \overline{\{(T - \lambda)p(T)y : p \in \mathcal{P}\}}$ is a non-trivial invariant subspace. Indeed, $y \notin M$ as $\|y - (T - \lambda)p(T)y\| \geq c$ for every polynomial p .

As there are examples of operators in Banach spaces without non-trivial invariant subspaces, in general it is not possible to replace the sequence $\{a_k\}$ by a constant $c > 0$. Thus Theorems 4 and 5 are the best possible, at least for Banach spaces.

Denote by $r(T)$ and $r_e(T)$ the spectral radius and the essential spectral radius of an operator $T \in B(X)$, respectively.

Theorems 4 and 5 easily imply the corresponding results for powers $T^i x$ (cf. [9] or [2]).

COROLLARY 6. *Let $T \in B(X)$ and let $\{a_k\}_{k=0}^\infty$ be a sequence of positive numbers with $\lim_{k \rightarrow \infty} a_k = 0$. Then there exists $x \in X$ such that*

$$\|T^k x\| \geq a_k r(T)^k \quad (k = 0, 1, \dots).$$

Proof. Let $\lambda \in \sigma_{\pi_e}(T)$ with $|\lambda| = \max\{|z| : z \in \sigma(T)\} = r(T)$. Then either λ is an eigenvalue of T and $\|T^k x\| = r(T)^k$ for the corresponding eigenvector x or $\lambda \in \sigma_{\pi_e}(T)$ and we can apply Theorem 4.

COROLLARY 7. *Let $T \in B(X)$, $x \in X$, $\varepsilon > 0$ and let $\{a_k\}_{k=0}^\infty$ be a sequence of positive numbers with $\lim_{k \rightarrow \infty} a_k = 0$. Then there exists $y \in X$ and a positive constant $C = C(\varepsilon)$ such that $\|y - x\| \leq \varepsilon$ and*

$$\|T^k y\| \geq C a_k r(T)^k \quad (k = 0, 1, \dots).$$

COROLLARY 8 (see [17]). *Let $T \in B(X)$. Then the set $\{x \in X : r(T, x) = r(T)\}$ is dense in X .*

As another corollary we deduce that the infimum and the supremum in the spectral radius formula

$$r(T) = \inf_{k \in \mathbb{N}} \|T^k\|^{1/k} = \inf_{k \in \mathbb{N}} \sup_{\|x\|=1} \|T^k x\|^{1/k}$$

can be interchanged.

COROLLARY 9. *Let $T \in B(X)$. Then*

$$r(T) = \inf_{k \in \mathbb{N}} \sup_{\|x\|=1} \|T^k x\|^{1/k} = \sup_{\|x\|=1} \inf_{k \in \mathbb{N}} \|T^k x\|^{1/k}.$$

II. Capacity. In the previous section we estimated $\|p(T)x\|$ by means of $|p(\lambda)|$ where λ was a fixed element of $\sigma_{\pi_e}(T)$. In this section we are looking for an estimate in terms of $\max\{|p(\lambda)| : \lambda \in \sigma_{\pi_e}(T)\}$. As $\delta\sigma_e(T) \supset \sigma_{\pi_e}(T)$, by the spectral mapping theorem for σ_e we have

$$\max_{\lambda \in \sigma_{\pi_e}(T)} |p(\lambda)| = \max_{\lambda \in \sigma_e(T)} |p(\lambda)| = \max\{|z| : z \in \sigma_e(p(T))\} = r_e(p(T)).$$

An important tool for the results in this section is the following classical lemma of Fekete [4]:

LEMMA 10. *Let K be a non-empty compact subset of the complex plane and let $k \geq 1$. Then there exist points $u_0, u_1, \dots, u_k \in K$ such that*

$$\max\{|p(z)| : z \in K\} \leq (k+1) \cdot \max_{0 \leq i \leq k} |p(u_i)|$$

for every polynomial p with $\deg p \leq k$.

By using the previous lemma and the results of the previous section we can get (see [10])

PROPOSITION 11. Let $T \in B(X)$, $\varepsilon \geq 0$ and $k \geq 1$. Then there exists $x \in X$ with $\|x\| = 1$ and

$$\|p(T)x\| \geq \frac{1 - \varepsilon}{2(k + 1)^2} r_e(p(T))$$

for every polynomial p with $\deg p \leq k$.

THEOREM 12. Let $T \in B(X)$, $x \in X$ and $\varepsilon > 0$. Then there exists $y \in X$ and a positive constant $C = C(\varepsilon)$ such that $\|y - x\| \leq \varepsilon$ and

$$\|p(T)y\| \geq C(1 + \deg p)^{-(2+\varepsilon)} r_e(p(T))$$

for every polynomial p .

REMARK. In case of a Hilbert space operator one can get a better estimate

$$\|p(T)y\| \geq C(1 + \deg p)^{-(1+\varepsilon)} r_e(p(T)).$$

The notion of capacity of an operator was defined by Halmos in [5]. If $T \in B(X)$ then

$$\text{cap } T = \lim_{k \rightarrow \infty} (\text{cap}_k T)^{1/k} = \inf_k (\text{cap}_k T)^{1/k}$$

where

$$\text{cap}_k T = \inf \{ \|p(T)\| : p \in \mathcal{P}_k^1 \}$$

and \mathcal{P}_k^1 is the set of all monic (i.e. with leading coefficient equal to 1) polynomials of degree k .

This is a generalization of the classical notion of capacity of a compact subset K of the complex plane:

$$\text{cap } K = \lim_{k \rightarrow \infty} (\text{cap}_k K)^{1/k} = \inf_k (\text{cap}_k K)^{1/k}$$

where

$$\text{cap}_k K = \inf \{ \|p\|_K : p \in \mathcal{P}_k^1 \} \quad \text{and} \quad \|p\|_K = \sup \{ |p(z)| : z \in K \}.$$

By the main result of [5], $\text{cap } T = \text{cap } \sigma(T)$.

The local capacity of T at x can be defined analogously:

$$\text{cap}_k(T, x) = \inf \{ \|p(T)x\| : p \in \mathcal{P}_k^1 \}$$

and

$$\text{cap}(T, x) = \limsup_{k \rightarrow \infty} \text{cap}_k(T, x)^{1/k}$$

(in general the limit does not exist).

It is easy to see that $\text{cap}(T, x) \leq \text{cap } T$ for every $x \in X$.

COROLLARY 13. Let $T \in B(X)$. Then the set $\{x \in X : \text{cap}(T, x) = \text{cap } T\}$ is dense in X .

PROOF. By Theorem 12 there exists a dense subset $Y \subset X$ such that

$$\|p(T)y\| \geq \frac{C}{(\deg p + 1)^3} r_e(p(T))$$

for every polynomial p . Then

$$\begin{aligned} \text{cap}_k(T, y) &= \inf\{\|p(T)y\| : p \in \mathcal{P}_k^1\} \geq \inf_{p \in \mathcal{P}_k^1} \frac{C}{(k+1)^3} r_e(p(T)) \\ &= \inf_{p \in \mathcal{P}_k^1} \frac{C}{(n+1)^3} \sup\{|p(\lambda)| : \lambda \in \sigma_e(T)\} = \frac{C}{(n+1)^3} \text{cap}_k \sigma_e(T). \end{aligned}$$

Thus

$$\begin{aligned} \text{cap}(T, y) &= \limsup_{k \rightarrow \infty} \text{cap}_k(T, y)^{1/k} \\ &\geq \limsup_{k \rightarrow \infty} \left(\frac{C}{(k+1)^3} \right)^{1/k} (\text{cap}_k \sigma_e(T))^{1/k} = \text{cap} \sigma_e(T). \end{aligned}$$

Further $\text{cap} \sigma_e(T) = \text{cap} \sigma(T)$ as $\sigma(T) - \sigma_e(T)$ contains only countably many isolated points in the unbounded component of the complement of $\sigma_e(T)$ and $\text{cap} \sigma(T) = \text{cap} T$ by [5]. Hence $\text{cap}(T, x) = \text{cap} T$ for every $y \in Y$.

An operator $T \in B(X)$ is called *quasialgebraic* if and only if $\text{cap} T = 0$. Similarly T is called *locally quasialgebraic* if $\text{cap}(T, x) = 0$ for every $x \in X$.

It follows from Corollary 13 that these two notions are equivalent (see [8]). This gives a positive answer to a problem of Halmos [5].

THEOREM 14. *An operator is quasialgebraic if and only if it is locally quasialgebraic.*

Theorem 14 is an analogue to the well-known result of Kaplansky: an operator is algebraic (i.e. $p(T) = 0$ for some non-zero polynomial p) if and only if it is locally algebraic (i.e. for every $x \in X$ there exists a polynomial $p_x \neq 0$ such that $p_x(T)x = 0$).

III. n -tuples of commuting operators. The results of the previous section admit a generalization for n -tuples of commuting operators.

Let $T = (T_1, \dots, T_n)$ be an n -tuple of mutually commuting operators in a Banach space X . We denote by $\sigma(T) \subset \mathbb{C}^n$ the *Harte spectrum* [6] of T , i.e. $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$ does not belong to $\sigma(T)$ if and only if there exist operators $L_1, \dots, L_n, R_1, \dots, R_n \in B(X)$ such that

$$\sum_{i=1}^n L_i(T_i - \lambda_i) = I = \sum_{i=1}^n (T_i - \lambda_i)R_i.$$

Denote further by $\sigma_{\pi e}(T)$ the *essential approximate point spectrum* of the n -tuple T , i.e. $\lambda = (\lambda_1, \dots, \lambda_n) \in \sigma_{\pi e}(T)$ if and only if

$$\inf \left\{ \sum_{i=1}^n \|(T_i - \lambda_i)x\| : x \in M, \|x\| = 1 \right\} = 0$$

for every subspace M of finite codimension.

The following result is a generalization of Theorem 12 for n -tuples of commuting operators (see [14]):

THEOREM 15. *Let $T = (T_1, \dots, T_n) \in B(X)^n$ be a mutually commuting n -tuple of operators. Let $x \in X$ and $\varepsilon > 0$. Then there exists $y \in X$ and a positive constant $C = C(\varepsilon)$ such that $\|y - x\| \leq \varepsilon$ and*

$$\|p(T)y\| \geq \frac{C}{(1 + \deg p)^{2n+\varepsilon}} r_e(p(T))$$

for every polynomial p with n variables.

Every polynomial p in n complex variables with $\deg p \leq k$ can be written in the form

$$p(z) = \sum_{|\alpha| \leq k} c_\alpha(p) z^\alpha$$

where $\alpha = (\alpha_1, \dots, \alpha_n)$ is an n -tuple of non-negative integers, $|\alpha| = \alpha_1 + \dots + \alpha_n$, the coefficients $c_\alpha(p)$ are complex, $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ and $z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n}$.

The notion of capacity of commuting n -tuples of operators was introduced by Stirling [16]:

Denote by $\mathcal{P}_k^1(n)$ the set of all “monic” polynomials $p(z) = \sum_{|\mu| \leq k} c_\mu(p) z^\mu$ of degree k in n variables with $\sum_{|\mu|=k} |c_\mu(p)| = 1$.

Let $T = (T_1, \dots, T_n)$ be an n -tuple of mutually commuting operators in a Banach space X . The joint capacity of T was defined in [16] by

$$\text{cap } T = \liminf_{k \rightarrow \infty} \text{cap}_k(T)^{1/k} \quad \text{where} \quad \text{cap}_k(T) = \inf\{\|p(T)\| : p \in \mathcal{P}_k^1(n)\}$$

(in fact the \liminf in the definition of $\text{cap } T$ can be replaced by limit, see [12]).

For a compact subset $K \subset \mathbb{C}^n$ define the corresponding capacity by

$$\text{cap } K = \lim_{k \rightarrow \infty} \text{cap}_k(K)^{1/k} \quad \text{where} \quad \text{cap}_k(K) = \inf\{\|p\|_K : p \in \mathcal{P}_k^1(n)\}.$$

This capacity was studied in [16] and called the “homogeneous Chebyshev constant” of K .

By [16], $\text{cap } \sigma(T) \leq \text{cap } T \leq 2^n \text{cap } \sigma(T)$. Actually, equality holds here (see [12]).

THEOREM 16. *Let $T = (T_1, \dots, T_n)$ be an n -tuple of mutually commuting operators in a Banach space X . Then $\text{cap } T = \text{cap } \sigma(T)$.*

THEOREM 17 (see [13]). *Let T be an n -tuple of mutually commuting operators in a Banach space X . Then $\sigma(T) - \widehat{\sigma}_{\pi_e}(T)$ consist of at most countably many isolated joint eigenvalues, where $\widehat{\sigma}_{\pi_e}(T)$ denotes the polynomially convex hull of $\sigma_{\pi_e}(T)$. In particular, $\text{cap } \sigma(T) = \text{cap } \sigma_{\pi_e}(T)$ (actually all reasonable joint spectra have the same capacity).*

Let $T = (T_1, \dots, T_n)$ be a commuting n -tuple of operators in a Banach space X and let $x \in X$. We define the local capacity $\text{cap}(T, x)$ by

$$\text{cap}(T, x) = \limsup_{k \rightarrow \infty} \text{cap}_k(T, x)^{1/k}$$

where

$$\text{cap}_k(T, x) = \inf\{\|p(T)x\| : p \in \mathcal{P}_k^1(n)\}.$$

Clearly $\text{cap}(T, x) \leq \text{cap}T$ for every $x \in X$.

THEOREM 18 (see [14]). *Let T be an n -tuple of mutually commuting operators in a Banach space X . Then the set of all $y \in X$ with $\text{cap}(T, y) = \text{cap}T$ is dense in X .*

References

- [1] C. Apostol, *On the left essential spectrum and non-cyclic operators in Banach spaces*, Rev. Roumaine Math. Pures Appl. 17 (1972), 1141–1147.
- [2] B. Beauzamy, *Introduction to Operator Theory and Invariant Subspaces*, North-Holland Math. Library 42, North-Holland, Amsterdam, 1988.
- [3] J. J. Buoni, R. Harte and T. Wickstead, *Upper and lower Fredholm spectra*, Proc. Amer. Math. Soc. 66 (1977), 309–314.
- [4] M. Fekete, *Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten*, Math. Z. 17 (1923), 228–249.
- [5] P. R. Halmos, *Capacity in Banach algebras*, Indiana Univ. Math. J. 20 (1971), 855–863.
- [6] R. Harte, *Tensor products, multiplication operators and the spectral mapping theorem*, Proc. Roy. Irish Acad. Sect. A 73 (1973), 285–302.
- [7] R. Harte and T. Wickstead, *Upper and lower Fredholm spectra II*, Math. Z. 154 (1977), 253–256.
- [8] V. Müller, *On quasialgebraic operators in Banach spaces*, J. Operator Theory 17 (1987), 291–300.
- [9] —, *Local spectral radius formula for operators in Banach spaces*, Czechoslovak Math. J. 38 (1988), 726–729.
- [10] —, *Local behaviour of the polynomial calculus of operators*, J. Reine Angew. Math. 430 (1992), 61–68.
- [11] —, *On the essential approximate point spectrum of operators*, Integral Equations Operator Theory 15 (1992), 1033–1041.
- [12] —, *A note on joint capacities in Banach algebras*, Czechoslovak Math. J. 43 (1993), 367–382.
- [13] —, *On the joint essential spectrum of commuting operators*, Acta Sci. Math. (Szeged) 57 (1993), 199–205.
- [14] V. Müller and A. Sołtysiak, *On local joint capacities of operators*, Czechoslovak Math. J. 43 (1993), 743–751.
- [15] S. Rolewicz, *On orbits of elements*, Studia Math. 32 (1969), 17–22.
- [16] D. S. G. Stirling, *The joint capacity of elements of Banach algebras*, J. London Math. Soc. 10 (1975), 374–389.
- [17] P. Vrbová, *On local spectral properties of operators in Banach spaces*, Czechoslovak Math. J. 23 (1973), 483–492.
- [18] V. P. Zakharyuta, *Transfinite diameter, Chebyshev constant and a capacity of a compact set in \mathbb{C}^n* , Mat. Sb. 96 (1975), 374–389 (in Russian).