

References

- [1] H. Andréka and I. Németi, *Formulas and ultraproducts in categories*, Beitrage zur Algebra u. Geom. 8 (1979), 133–151.
- [2] B. Banaschewski and H. Herrlich, *Subcategories defined by implications*, Houston Journ. Math. 2 (1976), 149–171.
- [3] B. J. Day, *Density representations of functors*, Bull. Austral. Math. Soc. 3 (1977), 321–480.
- [4] Y. Diers, *Type de densité d'une sous-catégorie pleine*, Ann. Soc. Sci. Bruxelles, I, 90 (1976), 25–47.
- [5] P. Gabriel und F. Ulmer, *Local presentierbare Kategorien*, Lecture Notes in Mathematics, Vol. 221, Springer-Verlag, 1971.
- [6] H. Herrlich and G. E. Strecker, *Category theory*, Allyn and Bacon, 1973.
- [7] R. John, *A note on implicational subcategories*, In: *Contributions to Universal Algebra* (Proc. Coll. Szeged 1975), Colloq. Math. Soc. J. Bolyai, Vol. 17, North-Holland, Amsterdam 1977, 213–222.
- [8] S. MacLane, *Categories for the working mathematician*, Springer-Verlag, 1971.
- [9] G. Matthiessen, *Regular and strongly finitary structures over regularly finitary categories*, Canadian J. Math. 30 (1978), 250–261.
- [10] I. Németi and I. Sain, *Cone-injective subcategories and some Birkhoff type theorems*, In: *Universal Algebra* (Proc. Coll. Esztergom 1977), Colloq. Math. Soc. J. Bolyai, Vol. 29, North-Holland, Amsterdam 1982, 535–578.
- [11] M. Smyth, *Power domains*, Journ. of Computer and System Sci. 16 (1) (1978).
- [12] H. Andréka and I. Németi, *Loś lemma holds in every category*, Studia Sci. Math. Hungar. 13 (1978), 361–376.
- [13] —, —, *Injectivity in categories to represent all first order formulas*, Demonstration Mathematica 12 (1979), 717–732.
- [14] I. Sain and B. H. Hien, *In which categories are first order axiomatisable hulls characterizable by ultraproducts*, Math. Inst. Hungar. Acad. Sci. Preprint No 56/1981, to appear in Cahiers. Diff. Geom. Top.

*Presented to the Semester
 Universal Algebra and Applications
 (February 15 – June 9, 1978)*

A GENERALIZATION OF ELEMENTARY FORMAL SYSTEMS

MELVIN FITTING

Herbert H. Lehman College, Bronx, N.Y., U.S.A.

1. Introduction

In [6] Smullyan gave an elegant development of recursion theory based on elementary formal systems. These dealt directly with words over a finite alphabet, and only indirectly with numbers, via “names” for them. We generalize the notion of elementary formal system, by separating “structural properties” from “subject matter.” The result provides a natural “recursion theory” for any structure, words and numbers being particular examples.

Our notion of recursion theory over the natural numbers can be turned into hyperarithmetical theory by the addition of a simple infinitary rule (an ω -rule) [1]. We formulate the rule so that it applies to all our recursion theories, turning them into what we call ω -recursion theories. For both recursion and ω -recursion theories we define a natural generalization of enumeration operator. We investigate the structural characteristics of these operators, and prove an analog of the First Recursion Theorem for them.

2. Elementary formal systems

Let \mathcal{A} be an infinite set, and let $\mathcal{R}_1, \dots, \mathcal{R}_k$ be relations on \mathcal{A} . We call $k+1$ tuple $\langle \mathcal{A}, \mathcal{R}_1, \dots, \mathcal{R}_k \rangle$ a *structure*. We allow trivial structures $\langle \mathcal{A} \rangle$. We set up a simple logical calculus relative to a particular structure, so for the rest of this section, let $\mathfrak{A} = \langle \mathcal{A}, \mathcal{R}_1, \dots, \mathcal{R}_k \rangle$ be a fixed structure.

We suppose available an unlimited supply of n -place predicate symbols for each $n > 0$. We informally use P, Q, R , etc. to represent them. The other two symbols of our alphabet are an *arrow* and a *comma*. We will use axiom schemas, so variables are not needed in the language itself, and we need no rule of substitution.

By an *atomic formula* we mean an expression of the form Pv_1, \dots, v_n where $v_1, \dots, v_n \in \mathcal{A}$ and P is an n -place predicate symbol. For convenience we may write Pv for Pv_1, \dots, v_n . We also define a *pseudo-atomic formula* to be anything of the form Pv_1, \dots, v_n where each x_i is in \mathcal{A} or is a variable. Pseudo-atomic formulas are expressions of the metalanguage only.

The notion of *formula* is defined by the following rules:

- 1) an atomic formula is a formula,
- 2) if X_1, X_2, \dots, X_n are formulas, so is $X_1 \rightarrow X_2 \rightarrow \dots \rightarrow X_n$. Formulas are to be thought of as being associated to the right. Thus $A \rightarrow B \rightarrow C \rightarrow D$ should be read as if it were $A \rightarrow (B \rightarrow (C \rightarrow D))$ and thought of as saying A , B and C together imply D .

The metalinguistical notion of *pseudo-formula* is defined analogously, being built up from pseudo-atomic formulas. And the notion of an *instance* of a pseudo-formula (over \mathcal{A}) also has an obvious definition. Any instance of a pseudo-formula is a formula.

By the *conclusion* of a (pseudo) formula we mean the final (pseudo) atomic part of it. Thus if A is (pseudo) atomic, A is the conclusion of $X \rightarrow A$, and also of A itself.

Let R_1, \dots, R_k be distinct predicate symbols permanently assigned to the relations $\mathcal{R}_1, \dots, \mathcal{R}_k$, such that R_i is n -place if \mathcal{R}_i is an n -ary relation. Let \mathcal{R}_i^* consist of all atomic formulas of the form $R_i v$ where $\mathcal{R}_i v$ is true.

We say a pseudo-formula X is *allowable* if none of R_1, \dots, R_k occurs in the conclusion of X .

Let $\{A_1, \dots, A_m\}$ be some finite set of pseudo-formulas, each allowable. By a *derivation from* $\{A_1, \dots, A_m\}$ (over \mathfrak{U}) we mean a finite sequence of formulas, X_1, \dots, X_i such that each term of the sequence either

- 1) is a member of $\mathcal{R}_1^* \cup \dots \cup \mathcal{R}_k^*$ or
- 2) is an instance of some A_i or
- 3) comes from two earlier terms by the rule

$$\text{MP} \quad \frac{X \quad X \rightarrow Y}{Y} \quad \text{provided } X \text{ is atomic.}$$

If there is a derivation ending with X , we say X is *derivable from* $\{A_1, \dots, A_m\}$ (over \mathfrak{U}).

$\{A_1, \dots, A_m\}$ determines, relative to \mathfrak{U} , a simple deductive system, called an *elementary formal system* (over \mathfrak{U}). Each A_i is an *axiom* for that elementary formal system.

Let P be a p -place predicate symbol, and $\mathcal{P} \subseteq \mathcal{A}^p$. We say P *represents* \mathcal{P} in the elementary formal system determined by $\{A_1, \dots, A_m\}$ if $v \in \mathcal{P}$ iff Pv is derivable from $\{A_1, \dots, A_m\}$.

We say \mathcal{P} is *representable in the elementary formal system determined by* A_1, \dots, A_m , if there is some predicate symbol P which represents \mathcal{P} in that elementary formal system.

Finally, we say \mathcal{P} is *recursively enumerable (r.e.)* (over \mathfrak{U}) if \mathcal{P} is representable in some elementary formal system (over \mathfrak{U}). Also \mathcal{P} is *recursive* (over \mathfrak{U}) if both \mathcal{P} and $\mathcal{A}^p - \mathcal{P}$ are r.e. (over \mathfrak{U}).

3. Examples

EXAMPLE 1. $\mathcal{A} = \omega$. \mathcal{R} is the successor relation: $\mathcal{R}x, y$ iff $y = x+1$. Let \mathfrak{U}_N be the structure $\langle \mathcal{A}, \mathcal{R} \rangle$. It can be shown that a relation is r.e. over \mathfrak{U}_N iff it is r.e. in the usual sense.

EXAMPLE 2. \mathcal{A} is the set of signed integers. \mathcal{R} is the successor relation for signed integers. Let \mathfrak{U}_Z be the structure $\langle \mathcal{A}, \mathcal{R} \rangle$. A relation is r.e. over \mathfrak{U}_Z iff under any standard Gödel numbering of the signed integers, it corresponds to an r.e. set of natural numbers.

EXAMPLE 3. $\mathcal{A} = \omega \times \omega$. We use two relations on \mathcal{A} . $\mathcal{R}_1 x, y$ iff the first component of y is the successor of the first component of x . \mathcal{R}_2 is similarly the second component successor relation. Let \mathfrak{U}_Q be the structure $\langle \mathcal{A}, \mathcal{R}_1, \mathcal{R}_2 \rangle$. A relation is r.e. over \mathfrak{U}_Q iff the result of applying a recursive pairing function to its members produces a relation r.e. in the conventional sense.

EXAMPLE 4. $\mathcal{A} = \text{H.F.} = R_\omega$. \mathcal{R}_1 is the union relation, $\mathcal{R}_1(x, y, z)$ iff $x \cup y = z$. \mathcal{R}_2 is the unordered pair relation, $\mathcal{R}_2(x, y, z)$ iff $\{x, y\} = z$. Let $\mathfrak{U}_S = \langle \mathcal{A}, \mathcal{R}_1, \mathcal{R}_2 \rangle$. A relation is r.e. over \mathfrak{U}_S iff it is Σ over H. F. *Remark:* This example is essentially unchanged if we use the single relation $x \cup \{y\} = z$.

EXAMPLE 5. \mathcal{A} is the set of words over a fixed finite alphabet. \mathcal{R} is the concatenation relation on \mathcal{A} . Let \mathfrak{U}_P be the structure $\langle \mathcal{A}, \mathcal{R} \rangle$. Elementary formal systems over \mathfrak{U}_P are those called *pure* in [6]. *Remark:* This example too is essentially unchanged if, instead of \mathcal{R} we use one relation for each letter a of our finite alphabet, taking, say, $\mathcal{R}_a x, y$ iff y is word x with a added to the end.

EXAMPLE 6. \mathcal{A} is the set of real numbers. Take the two relations of addition and order for reals. In the resulting structure, for example, the set of integers is r.e. Also, the function $f(x) = [x]$, mapping each real x to the greatest integer $\leq x$, is recursive (as a relation).

EXAMPLE 7. Let \mathfrak{U} be the structure $\langle F \cup V; F, +_F, \times_F, V, +_V \times_S, * \rangle$ where F is a field under the operations $+_F$ and \times_F , V is an n -dimensional vector space over F , with vector addition $+_V$ and scalar multiplication

\times_S , and $*$ is an inner product on V . Call a function *partial recursive* if its graph is r.e. The Gram-Schmidt orthogonalization process can be “captured” by an elementary formal system over this structure to show: there is a partial recursive function f , taking n -tuples of vectors to n -tuples of vectors, such that if $f(x_1, \dots, x_n) = \langle y_1, \dots, y_n \rangle$ then $\{x_1, \dots, x_n\}$ and $\{y_1, \dots, y_n\}$ span the same subspace of V , and $\{y_1, \dots, y_n\}$ is an orthogonal set (some of the y_i may be 0).

4. Enumeration operators

We modify elementary formal systems so that they may accept inputs as well as produce outputs.

Some notation. Suppose \mathfrak{U} is the structure $\langle \mathcal{A}, \mathcal{R}_1, \dots, \mathcal{R}_k \rangle$ and \mathcal{R} is some relation on \mathcal{A} . We write $\langle \mathfrak{U}, \mathcal{R} \rangle$ for the structure $\langle \mathcal{A}, \mathcal{R}_1, \dots, \mathcal{R}_k, \mathcal{R} \rangle$. Suppose E is a set of axioms for an elementary formal system over \mathfrak{U} , and A is a predicate symbol. We write $E \vdash_{\mathfrak{U}} Ax$ to mean there is a derivation of Ax from E over \mathfrak{U} . Then $\{x \mid E \vdash_{\mathfrak{U}} Ax\}$ is the relation which A represents in the elementary formal system (with axioms) E over \mathfrak{U} .

Now, let \mathfrak{U} be some structure $\langle \mathcal{A}, \mathcal{R}_1, \dots, \mathcal{R}_k \rangle$, fixed for the rest of this section. Suppose \mathcal{P} is some n -place relation on \mathcal{A} . Let E be an elementary formal system over the structure $\langle \mathfrak{U}, \mathcal{P} \rangle$ in which, say, the predicate symbol P has been assigned to \mathcal{P} . Let Q be an m -place predicate symbol. Using the axioms E , Q represents a certain m -ary relation on \mathcal{A} . Now suppose we keep E fixed, but change \mathcal{P} to \mathcal{P}' , still using P to represent it. Then Q will represent a different relation on \mathcal{A} . In this way a certain operator on \mathcal{A} is created, which we may symbolize by $[E_Q^P]$. It uses the axioms E , takes whatever P represents as input, and gives whatever Q represents as output. Formally

$$[E_Q^P](\mathcal{P}) = \{x \mid E \vdash_{\langle \mathfrak{U}, \mathcal{P} \rangle} Qx\}$$

[where the predicate symbol P is assigned to \mathcal{P}].

We call the maps $[E_Q^P]$ *enumeration operators* over the structure \mathfrak{U} .

Let \mathfrak{U}_N be the structure of arithmetic as defined in Example 1 of the last section. It can be shown that the enumeration operators over \mathfrak{U}_N coincide with enumeration operators as defined in [5].

Let $[E_Q^P]$ be an enumeration operator over the structure \mathfrak{U} taking n -ary relations to m -ary relations. That is, P is n -place and Q is m -place. We call pair $\langle n, m \rangle$ the *order* of $[E_Q^P]$. Of course, $n, m > 0$.

5. The ω -rule

We add an infinite-premise rule of derivation to the machinery of elementary formal systems as defined above.

First, we modify the alphabet by adding the additional symbol ∇ .

Now an atomic formula is a string Px_1, \dots, x_n where P is n -place and each x_i either is in \mathcal{A} , or is ∇ . (We similarly modify the notion of pseudo-atomic formula, and formula.) Otherwise no syntactical changes are made; *instance* still means instance over \mathcal{A} ; for example.

Intuitively, Pv, ∇, w is to mean Pv, a, w holds for each $a \in \mathcal{A}$. Now we give rules governing the formal use of ∇ . We make the *restriction* that ∇ may not occur in the conclusion of any axiom schema. And we add one more rule of derivation.

ω -rule: If Pv, ∇, w is atomic, then

$$\frac{Pv, a, w \text{ for each } a \in \mathcal{A}}{Pv, \nabla, w}$$

The notion of an ω -elementary formal system is formulated as expected. Derivations are now well ordered (possibly) infinite sequences, allowing ω -rule applications. Call a relation $\mathcal{P} \subseteq \mathcal{A}^n$ ω -r.e. over \mathfrak{U} if it is representable in some ω -elementary formal system over \mathfrak{U} . Also \mathcal{P} is ω -recursive if both \mathcal{P} and $\mathcal{A}^n - \mathcal{P}$ are ω -r.e.

Let \mathfrak{U}_N be the structure of arithmetic (Example 1 from § 3). In [1] is a direct proof that, for \mathfrak{U}_N , ω -r.e. is the same as Π_1^1 and ω -recursive is the same as hyperarithmetical. More generally, in any structure, ω -recursive coincides with hyperelementary, as defined in [4].

The definition of enumeration operator over \mathfrak{U} may be modified in the obvious way to define the notion of ω -enumeration operator over \mathfrak{U} . We skip the details.

6. Basic structural properties

The collection of enumeration operators over a structure $\mathfrak{U} = \langle \mathcal{A}, \mathcal{R}_1, \dots, \mathcal{R}_k \rangle$ and the collection of ω -enumeration operators over \mathfrak{U} have certain common structural features, which we now develop. We note that the ω -rule plays no role in many of our proofs below, so we can treat recursion and ω -recursion theories simultaneously.

THEOREM 6.1. *Suppose I and J are (ω) enumeration operators over \mathfrak{U} , I is of order $\langle n, m \rangle$ and J is of order $\langle m, p \rangle$. Then the composition $J \circ I$ is also an (ω) enumeration operator over \mathfrak{U} , of order $\langle n, p \rangle$.*

Proof. Say $I = [E_D^A]$ and $J = [E'_D{}^C]$. We may suppose without loss of generality that E and E' contain no predicate symbols in common other than R_1, \dots, R_k . (Call E and E' *disjoint* if this is the case.) Consider the (ω) enumeration operator $[F_D^A]$ where F consists of
the axioms of E ,
the axioms of E' ,
 $Bx \rightarrow Cx$.

It is easy to see that $J \circ I = [F_D^A]$.

DEFINITION. Let I and J be (ω) enumeration operators over \mathfrak{A} , both of order $\langle n, m \rangle$. We define two maps as follows:

$$(I \cap J)(\mathcal{P}) = I(\mathcal{P}) \cap J(\mathcal{P}),$$

$$(I \cup J)(\mathcal{P}) = I(\mathcal{P}) \cup J(\mathcal{P}).$$

THEOREM 6.2. The collection of (ω) enumeration operators over \mathfrak{A} is closed under \cap and \cup .

Proof. Say $I = [E_B^A]$ and $J = [E'_D]$ where E and E' are disjoint. Then $I \cap J = [F_H^A]$ and $I \cup J = [F_K^A]$ where F consists of the axioms of E , the axioms of E' , $Gx \rightarrow Ax$, $Gx \rightarrow Cx$, $By \rightarrow Dy \rightarrow Hy$, $By \rightarrow Ky$, $Dy \rightarrow Ky$.

DEFINITION. Let I be an (ω) enumeration operator of order $\langle n, m \rangle$ and let J be an (ω) enumeration operator of order $\langle n, m' \rangle$ over \mathfrak{A} . By $I \times J$ we mean the map of order $\langle n, m + m' \rangle$ given by

$$(I \times J)(\mathcal{P}) = I(\mathcal{P}) \times J(\mathcal{P}).$$

THEOREM 6.3. The collection of (ω) enumeration operators over \mathfrak{A} is closed under \times .

DEFINITION. P^n is the projection operator of order $\langle n, n-1 \rangle$ defined by

$$P^n(\mathcal{P}) = \{ \langle x_1, \dots, x_{n-1} \rangle \mid (\exists y) \langle y, x_1, \dots, x_{n-1} \rangle \in \mathcal{P} \}.$$

D^n is the dual projection operator of order $\langle n, n-1 \rangle$ defined by

$$D^n(\mathcal{P}) = \{ \langle x_1, \dots, x_{n-1} \rangle \mid (\forall y) \langle y, x_1, \dots, x_{n-1} \rangle \in \mathcal{P} \}.$$

THEOREM 6.4. 1) P^n is both an enumeration and an ω -enumeration operator over \mathfrak{A} .

2) D^n is an ω -enumeration operator over \mathfrak{A} .

DEFINITION. Let S be a map from \mathcal{A}^n to \mathcal{A}^k . S is a sequential operator if, for every $j \leq k$, either

1) there is an $i \leq n$ such that for every $\alpha \in \mathcal{A}^n$, the j th term of $S(\alpha)$ is the i th term of α , or

2) there is some $c \in \mathcal{A}$ such that for every $\alpha \in \mathcal{A}^n$ the j th term of $S(\alpha)$ is c .

DEFINITION. Let $S: \mathcal{A}^n \rightarrow \mathcal{A}^k$ be a sequential operator. By S^{-1} : power set $(\mathcal{A}^k) \rightarrow$ power set (\mathcal{A}^n) we mean the map given by

$$S^{-1}(\mathcal{B}) = \{ \mathbf{v} \in \mathcal{A}^n \mid S(\mathbf{v}) \in \mathcal{B} \}.$$

We call such a map an explicit transformation.

THEOREM 6.5. Every explicit transformation is an (ω) enumeration operator.

THEOREM 6.6. Each (ω) enumeration operator I over \mathfrak{A} is monotone, that is, $\mathcal{A} \subseteq \mathcal{B} \Rightarrow I(\mathcal{A}) \subseteq I(\mathcal{B})$.

THEOREM 6.7. Each enumeration operator I over \mathfrak{A} is compact, that is, $\mathbf{v} \in I(\mathcal{P})$ iff for some finite $\mathcal{F} \subseteq \mathcal{P}$, $\mathbf{v} \in I(\mathcal{F})$.

7. The first recursion theorem

THEOREM 7.1. Let I be an (ω) enumeration operator of order $\langle n, n \rangle$. I has a minimal fixed point which is (ω) r.e.

Proof. I is monotone. Set

$$I^a = I\left(\bigcup_{\beta < a} I^\beta\right).$$

By standard monotone operator arguments (see [4]) $I^\infty = \bigcup_a I^a$ is the minimal fixed point of I . We show it is (ω) r.e. [Remark: If I is an enumeration operator, it is compact, in which case I^∞ is simply I^ω .]

Say $I = [E_B^A]$. (We note that, in E , A can not occur in the conclusion of any axiom.) By definition,

$$\mathbf{x} \in I(\mathcal{P}) \quad \text{iff} \quad E \upharpoonright_{\langle \mathfrak{A}, \mathcal{P} \rangle} B\mathbf{x}.$$

Now let $E' = E \cup \{B\mathbf{x} \rightarrow A\mathbf{x}\}$ (that is, E' is E + "output = input"). We claim B represents I^∞ using E' . That is,

$$I^\infty = \{ \mathbf{x} \mid E' \upharpoonright_{\mathfrak{A}} B\mathbf{x} \}.$$

1) For each a , we show

$$I^a \subseteq \{ \mathbf{x} \mid E' \upharpoonright_{\mathfrak{A}} B\mathbf{x} \}$$

by induction on a . Well, suppose

$$I^{<a} \subseteq \{ \mathbf{x} \mid E' \upharpoonright_{\mathfrak{A}} B\mathbf{x} \}$$

[where $I^{<a} = \bigcup_{\beta < a} I^\beta$ so that $I^a = I(I^{<a})$].

Also suppose $\mathbf{a} \in I^a$ then $\mathbf{a} \in I(I^{<a})$, so, by definition

$$(*) \quad E \upharpoonright_{\langle \mathfrak{A}, I^{<a} \rangle} B\mathbf{a}.$$

By induction hypothesis, for all $\mathbf{x} \in I^{<a}$, $E' \upharpoonright_{\mathfrak{A}} B\mathbf{x}$. But then, for all $\mathbf{x} \in I^{<a}$,

$$(**) \quad E' \upharpoonright_{\mathfrak{A}} A\mathbf{x}.$$

By (*) and (**), since also $E \subseteq E'$,

$$E' \upharpoonright_{\mathfrak{A}} B\mathbf{a}.$$

Hence $I^a \subseteq \{ \mathbf{x} \mid E' \upharpoonright_{\mathfrak{A}} B\mathbf{x} \}$.

2) $I^\infty \subseteq \{ \mathbf{x} \mid E' \upharpoonright_{\mathfrak{A}} B\mathbf{x} \}$.

This is immediate.

3) $\{ \mathbf{x} \mid E' \upharpoonright_{\mathfrak{A}} B\mathbf{x} \} \subseteq I^\infty$.

We show this by an induction on the length of derivations from E' in \mathfrak{U} .

Suppose: if Bx is derivable from E' in $< \alpha$ steps, then $x \in I^\infty$.

Suppose: Ba is derivable from E' in α steps. We show $a \in I^\infty$.

Let $\mathcal{R}_A = \{x \mid Ax \text{ occurs before the last line in some (fixed) derivation from } E' \text{ of } Ba\}$.

Let $\mathcal{R}_B = \{x \mid Bx \text{ occurs before the last line in this derivation}\}$.

Now, A occurs in the conclusion of only one axiom in E' , namely $Bx \rightarrow Ax$. It follows that

$$\mathcal{R}_A \subseteq \mathcal{R}_B.$$

Also, by induction hypothesis, $\mathcal{R}_B \subseteq I^\infty$. Hence

$$(*) \quad \mathcal{R}_A \subseteq I^\infty.$$

Finally, it should be clear that

$$E' \vdash_{\langle \mathcal{R}_A, \mathcal{R}_A \rangle} Ba.$$

This says

$$a \in I(\mathcal{R}_A).$$

Then by (*), since I is monotone,

$$a \in I(I^\infty) = I.$$

This concludes the proof.

References

- [1] M. Fitting, *Elementary formal systems for hyperarithmetical relations*, Zeitschrift für Mathematische Logik und der Grundlagen der Mathematik (to appear).
- [2] T. Grillo, *Dissecting abstract recursion theory*, in: *Generalized Recursion Theory* (Fenstad & Hinman, editors), North Holland, Amsterdam 1974, pp. 405-420.
- [3] S. Kripke, (abstracts), *Transfinite recursion on admissible ordinals I and II*, Journal of Symbolic Logic 29 (1964), 161-162.
- [4] Y. Moschovakis, *Elementary induction on abstract structures*, North Holland, Amsterdam 1974.
- [5] H. Rogers, *Theory of recursive functions and effective computability*, McGraw-Hill, N. Y., 1967.
- [6] R. Smullyan, *Theory of formal systems*, Revised Edition, Princeton University Press, Princeton 1961.

*Presented to the Semester
 Universal Algebra and Applications
 (February 15 - June 9, 1978)*

ON COMPATIBLE AND ORDER-PRESERVING FUNCTIONS ON LATTICES

D. DORNINGER and G. EIGENTHALER

*Institute of Algebra and Discrete Mathematics,
 Technical University, Vienna, Austria*

1

Let V be a lattice, k a positive integer and $F_k(V)$ the direct power $V^{(F^k)}$. A function $f \in F_k(V)$ is called *compatible* if for any congruence θ on V and $(a_i, b_i) \in \theta$, $i = 1, \dots, k$, $(f(a_1, \dots, a_k), f(b_1, \dots, b_k)) \in \theta$ holds, and f is called *order-preserving* if $a_i \leq b_i$, $i = 1, \dots, k$, implies $f(a_1, \dots, a_k) \leq f(b_1, \dots, b_k)$. We denote by $C_k(V)$ the set of all k -place compatible functions on V and by $OF_k(V)$ the set of all k -place order-preserving functions on V . As it immediately follows by a result of Wille [12], $OF_k(V) \subseteq C_k(V)$ iff V is simple.

In the present paper we determine all distributive lattices V with $C_k(V) \subseteq OF_k(V)$, and we give necessary conditions for an arbitrary lattice V to satisfy $C_k(V) \subseteq OF_k(V)$. Thereby we obtain necessary conditions for a lattice to be (locally) k -affine complete and (locally) k -order affine complete resp. (for these concepts of completeness cf. Schweigert [9] and Wille [12]). Furthermore, we show that every distributive lattice is locally k -order affine complete (generalizing a result of Grätzer [4]) and that 1-affine completeness implies k -affine completeness in case of a distributive lattice.

Throughout this paper we adopt the following notational conventions: join, meet, inclusion, and proper inclusion in a lattice are denoted by \cup , \cap , \leq , and $<$, resp.; k always stands for a positive integer and V always denotes a lattice.

2

First we show that it is sufficient to consider the case $k = 1$ in order to answer the question whether $OF_k(V) \subseteq C_k(V)$ and whether $C_k(V) \subseteq OF_k(V)$, resp.