

Rotundity, smoothness and duality

by

Jean-Paul Penot

Laboratoire de Mathématiques appliquées
CNRS FRE 2570
Av. de l'Université, 64000 Pau, France

Abstract: The duality between smoothness and rotundity of functions is studied in a nonlinear abstract framework. Here smoothness is enlarged to subdifferentiability properties and rotundity is formulated by means of approximation properties.

Keywords: conjugacy, convexity, coupling, duality, estimates, generalized convexity, rotundity, smoothness, subdifferential.

1. Introduction

The main interest of the many transforms known in mathematics (Fourier transform, Laplace transform, Radon transform, Fenchel transform, ...) lies in the fact that they convert a desired property into another one which may be more tractable. It has been known for several decades that Fenchel duality transforms a smoothness property into a well-posedness property and vice-versa. The first instance of such a phenomenon is probably constituted by the Smulian theorems about Fréchet differentiability of norms (Beauzamy, 1982, Deville, Godefroy and Zizler, 1993, Diestel, 1975, etc.). Asplund and Rockafellar (1969) have extended that result to general convex functions on topological vector spaces in duality. Following previous results of Vladimirov, Nesterov and Chekanov (1978) and Zalinescu (1983), Azé and the author (1995) have stressed a quantitative viewpoint in the preceding connection. Azé (1999) and Azé and Rahmouni (1994, 1995, 1996) have extended the preceding results to the case an element of the subdifferential is replaced by the whole subdifferential. Such questions are considered in the books (Azé, 1997; Zalinescu, 2002). They are linked with well-posedness properties.

It is our purpose here to consider the preceding properties in a unified way and to extend them to the case of general dualities. We introduce subdifferentiability properties which are uniform with respect to a certain set and dually, we use a uniform B-differentiability property (in the terminology of Pang, 1990,

1995; Robinson, 1991). After completing a draft of our paper, papers by S. Rolewicz came to our attention (Rolewicz, 1993, 1996). In these references, and in the monograph (Pallaschke, Rolewicz, 1997) an extension of the correspondence between rotundity and smoothness to a nonlinear framework is given, which anticipated our work. However our approach includes the subdifferential case and the uniform properties just mentioned; moreover, being set in the framework of conjugacies using coupling functions, it is more symmetric, so that it allows to interchange the roles of the spaces.

Our study underlines the interest of general dualities which have been brought to the fore in many articles (Balder, 1977; Dolecki, Kurcyusz, 1978; Fan, 1963; Flores-Bazán, Martínez-Legaz, 1998; Ioffe, 2001; Martínez-Legaz, 1988, 1990, 1991, 1995; Moreau, 1970; Penot, 1982, 1997, 2000, 2001; Penot, Volle, 1987, 1988, 1990; Singer, 1986, 1987, etc.) and treated in a systematic way in several recent monographs (Pallaschke, Rolewicz, 1997; Rubinov, 2000; Singer, 1997). However, for the sake of simplicity, we do not tackle the case of the most general dualities but we limit our study to the case of conjugacies, i.e. dualities of Fenchel-Moreau type, which are obtained by means of coupling functions. In doing so, our results remain close enough to the case of the classical Fenchel duality and they keep their intuitive character.

Let us observe that the efforts in putting duality into a general metric framework in the works quoted above are not isolated in mathematics. Parallel developments extending differential calculus and dynamical systems to metric spaces (Aubin, 1999; Pichard, 2001, etc.), differential geometry to metric spaces (Gromov, 1999; Lafontaine, Pansu, 1981), measure theory and probability theory to general topological spaces or metric spaces (Dellacherie, Meyer, 1975; Elworthy, 1975; Schwartz, 1973, 1980, etc.), criteria for estimates to metric spaces (Azé, Corvellec, Lucchetti, 2002; Ioffe, 2001), have reached remarkable achievements which give some hopes for duality questions.

After recalling some basic facts about conjugacies in the next section, we exhibit two conditions relating the coupling to the metric. They play a key role in the sequel. We also present some examples. The main estimates are obtained in Sections 3 (coercivity properties), 4 (passage from growth properties to upper estimates), and 5 (reverse passage).

2. Preliminaries and metric conjugacies

It will be convenient to denote by A the set of functions α from \mathbb{R}_+ into $\mathbb{R}_+ \cup \{+\infty\}$ satisfying $\alpha(0) = 0$ (in Azé and Penot, 1995, such functions are called Asplund functions). Any element α of A is identified with its even extension. The *reduced function* (or slope) associated with α is the function $\hat{\alpha}$ given by $\hat{\alpha}(0) = 0$, $\hat{\alpha}(t) = t^{-1}\alpha(t)$ for $t > 0$. When $\hat{\alpha}$ is nondecreasing, α is said to be *starshaped*. It is said to be *firm* if $(t_n) \rightarrow 0$ whenever $(\alpha(t_n)) \rightarrow 0$. An Asplund function γ is said to be a *gage* (or *admissible* or *forcing*) if it is firm and nondecreasing; equivalently, $\gamma \in A$ is a gage if it is nondecreasing and if

$\gamma(t) > 0$ for all $t > 0$. An Asplund function $\mu \in A$ is said to be a *modulus* if μ is nondecreasing and if $\lim_{t \searrow 0} \mu(t) = 0$. It is said to be an *hypermodulus* (or a remainder) if its reduced function $\widehat{\mu}$ is a modulus.

The following result of Asplund (Asplund, 1968), Lemma 1, see also Azé and Penot, 1995, Lemma 2.1) connects the preceding properties when taking conjugates. Recall that the (Fenchel) *conjugate* of α is the function α^* given by $\alpha^*(t) := \sup \{st - \alpha(s) : s \in \mathbb{R}\}$.

LEMMA 2.1 (a) For any starshaped element γ of the set G of gages, γ^* is an hypermodulus.

(b) For any element ω of the set Ω of hypermodulus, ω^* is a starshaped gage.

We devote the rest of the present section to some basic facts about conjugacies and we present some material we will use. We refer to Balder (1977), Dolecki, Kurcyusz (1978), Moreau (1970), Pallaschke, Rolewicz (1977), Penot (2000), Penot, Volle (1987), Rubinov (2000), Singer (1997) and to several other items of our bibliography for further information.

In the sequel, X and Y are metric spaces (or semi-metric spaces) whose metrics (or semi-metrics) are denoted by d inasmuch there is no risk of confusion. For a subset S of X , d_S stands for the distance function to S : $d_S(x) := \inf \{d(x, w) : w \in S\}$; the closed ball with center x and radius r is denoted by $B(x, r)$ or $B_X(x, r)$. A *coupling* or pairing of X with Y is a function $c : X \times Y \rightarrow \overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$. Given a function $f : X \rightarrow \overline{\mathbb{R}}$, its conjugate is the function $f^c : Y \rightarrow \overline{\mathbb{R}}$ defined by

$$f^c(y) := - \inf \{f(x) - c(x, y) : x \in X\}, \quad (1)$$

where $r - s$ mean $r + (-s)$, the addition of $\overline{\mathbb{R}}$ being the extension of the addition of \mathbb{R} given by $+\infty + s = +\infty$ for each $s \in \overline{\mathbb{R}}$, $-\infty + s = -\infty$ for each $s \in \mathbb{R} \cup \{-\infty\}$. When f (or c) takes only finite values, one gets the more familiar expression $f^c(y) = \sup \{c(x, y) - f(x) : x \in X\}$. We refer to Moreau (1970) for the subtleties of the calculus rules in $\overline{\mathbb{R}}$. For the sake of simplicity, in the following sections we assume that the coupling is real-valued.

The *subdifferential* $\partial^c f$ of $f : X \rightarrow \overline{\mathbb{R}}$ at $x_0 \in \text{dom} f := f^{-1}(\mathbb{R})$ associated with a coupling c is defined by: $y_0 \in \partial^c f(x_0)$ iff $c(x_0, y_0) \in \mathbb{R}$ and

$$\forall x \in X \quad f(x) \geq f(x_0) + c(x, y_0) - c(x_0, y_0). \quad (2)$$

Thus one sees that $y_0 \in \partial^c f(x_0)$ iff $c(x_0, y_0) \in \mathbb{R}$ and the Young-Fenchel equality holds:

$$f(x_0) + f^c(y_0) = c(x_0, y_0).$$

The symmetry between X and Y enables us to define in a similar way the subdifferential $\partial^c g$ of a function g on Y .

The following definitions relate the coupling with the metrics of the spaces.

DEFINITION 2.1 Given $\lambda > 0$, one says that the coupling c is a λ -submetric coupling at $(x_0, y_0) \in X \times Y$ if $c(x_0, y_0) \in \mathbb{R}$ and for any $(x, y) \in X \times Y$ one has

$$c(x, y) - c(x_0, y) \leq c(x, y_0) - c(x_0, y_0) + \lambda d(x, x_0)d(y, y_0). \quad (3)$$

If S and T are subsets of X and Y respectively, one says that c is a λ -submetric coupling at $S \times T$ if it is a λ -submetric coupling at (x_0, y_0) for any $(x_0, y_0) \in S \times T$. When $S = X$, $T = Y$, one simply says that c is a λ -submetric coupling.

Changing the metric d in X or Y to λd would enable to take $\lambda = 1$; in such a case we say that c is a *submetric coupling* (at $(x_0, y_0) \in X \times Y$, etc...). For a real-valued coupling, the preceding definition is symmetric in X and Y . Note that relation (3) can also be written in the following non symmetric form which allows for a clear comparison with the next notion: for any $r \in \mathbb{R}_+$, $x \in X$, one has

$$\sup_{y \in B(y_0, r)} (c(x, y) - c(x_0, y)) \leq c(x, y_0) - c(x_0, y_0) + \lambda r d(x, x_0).$$

DEFINITION 2.2 Given $\kappa > 0$, one says that the coupling c is a κ -super-metric coupling at $(x_0, y_0) \in X \times Y$ if $c(x_0, y_0) \in \mathbb{R}$ and for any $r \in \mathbb{R}_+$, $x \in X$, one has

$$\sup_{y \in B(y_0, r)} (c(x, y) - c(x_0, y)) \geq c(x, y_0) - c(x_0, y_0) + \kappa r d(x, x_0), \quad (4)$$

where $B(y_0, r)$ denotes the closed ball with center y_0 and radius r . One says that c is a κ -super-metric coupling at $(S, T) \subset (X, Y)$ if it is a κ -super-metric coupling at $(x_0, y_0) \in X \times Y$ for any $(x_0, y_0) \in S \times T$.

Observe that in this second definition the roles of X and Y are not symmetric. If the transposed coupling $c^T : Y \times X \rightarrow \overline{\mathbb{R}}$ given by $c^T(y, x) := c(x, y)$ is also a κ -super-metric coupling with respect to (y_0, x_0) , then we say that c is κ -super-metric pairing with respect to (x_0, y_0) . When $\kappa = 1$ we omit to mention it; moreover, c is said to be a *metric coupling* if it is both a submetric coupling and a super-metric coupling.

An anonymous referee raised the question of comparing the preceding notion with the concept of monotonicity property with constant $\kappa \in (0, 1)$ introduced in Rolewicz (1993, 1994). In order to answer this question let us say that c has the κ -monotonicity property at (x_0, y_0) if for each $x \in X$ there exists a sequence (r_n) of positive real numbers with limit 0 such that for each $n \in \mathbb{N}$ one has

$$\sup_{y \in B(y_0, r_n)} (c(x, y) - c(x_0, y)) \geq c(x, y_0) - c(x_0, y_0) + \kappa r_n d(x, x_0). \quad (5)$$

Then, following Rolewicz (1993, 1994), one says that c has the κ -monotonicity property if it has it at any $(x_0, y_0) \in X \times Y$. Clearly, the κ -monotonicity property

at (x_0, y_0) is a consequence of the property that c is a κ -super-metric coupling at (x_0, y_0) , hence the κ -monotonicity property is a consequence of the property that c is a κ -super-metric coupling at (X, Y) . It may happen that the κ -monotonicity property at (x_0, y_0) holds while c is not a κ -super-metric coupling at (x_0, y_0) . This fact occurs when $X = Y = \mathbb{R}$, $x_0 = y_0 = 0$, $c(x, y) = |x||y|$ for $(x, y) \in X \times [-1, 1]$, $c(x, y) = 0$ otherwise (take any $\kappa \in (0, 1)$ and any sequence (r_n) with limit 0). On the other hand, our conditions bear on a particular pair (x_0, y_0) while condition (5) has to be satisfied for any pair $(x_0, y_0) \in X \times Y$. When Y contains an isolated point \bar{y} the κ -monotonicity property at (x_0, \bar{y}) cannot hold, whatever x_0 is.

Let us give some examples. As just mentioned, they show that in general one cannot expect that the properties defined above are valid for any couple (x_0, y_0) .

EXAMPLE 1 (classical duality)

Let X and Y be normed vector spaces in metric duality. This means that there exists a continuous bilinear map $c : X \times Y \rightarrow \mathbb{R}$ such that the associated mappings $c_X : X \rightarrow Y^*$, $c_Y : Y \rightarrow X^*$ given by $c_X(x)(y) := c(x, y)$, $c_Y(y)(x) := c(x, y)$ are isometric embeddings. Then c and c^T are metric couplings. Note that the present example contains the cases $Y = X^*$ and $X = Y^*$ and other cases. This framework is often convenient. For instance, when $X = L_p(T, E)$, $Y = L_q(T, E^*)$, where T is an interval, E is a Banach space and $1 \leq p < \infty$, $q = (1 - 1/p)^{-1}$ one does not need to assume that the Radon-Nikodým property holds for E , as it would be the case if one wanted to have $Y = X^*$.

EXAMPLE 2

Let X and Y be normed vector spaces and let $c : X \times Y \rightarrow \mathbb{R}$ be an arbitrary continuous bilinear map. Then, for any $\lambda \geq \|c\|$, c is a λ -submetric coupling. If c_X open at 0 with rate κ onto its image, i.e. if for each $t > 0$ one has $B_{Y^*}(0, \kappa t) \cap c_X(X) \subset c_X(B_X(0, t))$, and if c_X is injective, then c is a κ -super-metric coupling at (x_0, y_0) for each $(x_0, y_0) \in X \times Y$. A similar result holds for c^T .

EXAMPLE 3 (Stepanov duality)

Let X and Y be metric spaces and let $c_Y : Y \rightarrow \mathbb{R}^X$ be an arbitrary map. Then, setting $c(x, y) := c_Y(y)(x)$, we get a coupling $c : X \times Y \rightarrow \mathbb{R}$. Suppose the image of c_Y is contained in the set $S(X, x_0)$ of Stepanov functions at x_0 (or stable at x_0 functions) which is the set of functions $f : X \rightarrow \mathbb{R}$ such that there exists some $k > 0$ for which $|f(x) - f(x_0)| \leq kd(x, x_0)$. Endow $S(X, x_0)$ with the semi-norm given by

$$\|f\|_S^0 := \sup_{x \in X \setminus \{x_0\}} \frac{|f(x) - f(x_0)|}{d(x, x_0)}, \quad f \in S(X, x_0)$$

or the norm given by $\|f\|_S := \|f\|_S^0 + |f(x_0)|$. If c_Y is stable at $y_0 \in Y$ in the sense that there exists some $\lambda > 0$ such that, for each $y \in Y$, one has

$\|c_Y(y) - c_Y(y_0)\|_S^0 \leq \lambda d(y, y_0)$, then c is a λ -submetric coupling at (x_0, y_0) . If c_Y is a Lipschitz mapping with rate λ , the coupling c is a λ -submetric coupling at (x_0, Y) . If c_Y is such that, for each $y \in Y$, one has $\|c_Y(y) - c_Y(y_0)\|_S^0 \geq \kappa d(y, y_0)$ then c is a κ -super-metric coupling at (x_0, y_0) .

EXAMPLE 4

In particular, if X is a metric space not reduced to x_0 , $Y = S(X, x_0)$ endowed with the norm $\|\cdot\|_S$ and if c is the evaluation mapping, c is a metric coupling at (x_0, Y) , as follows from the fact that for any $r > 0$ the function $y_0 + rd(x_0, \cdot)$ belongs to the closed ball of center y_0 and radius r in Y .

EXAMPLE 5 (Lipschitz duality, Martínez-Legaz, 1988; Pallasche, Rolewicz, 1997; Rolewicz, 1994)

Suppose the mapping c_Y of the preceding example takes its values in the subspace $L(X)$ of Lipschitzian functions on X . Given $x_0 \in X$, endow $L(X)$ with the norm

$$\|f\|_L := \sup_{w, x \in X, w \neq x} \frac{|f(w) - f(x)|}{d(w, x)} + |f(x_0)|.$$

If c_Y is stable at $y_0 \in Y$, then c is a λ -submetric coupling at (X, y_0) for some $\lambda > 0$. If c_Y is a Lipschitz mapping with rate λ , c is a λ -submetric coupling at (X, Y) .

In particular, if X is a metric space not reduced to x_0 , $Y = L(X)$ and if c is the evaluation mapping, c is a metric coupling.

EXAMPLE 6 (homogeneous duality, Rubinov, 2000, and its references)

Let X be a normed vector space, let Y be a metric space and let $c : X \times Y \rightarrow \mathbb{R}$ be such that the image of c_Y is contained in the space $H(X)$ of positively homogeneous functions on X which are bounded on the unit ball B of X . Let us endow $H(X)$ with the norm given by $\|h\| = \sup\{|h(x)| : x \in B\}$. If c_Y is stable at some $y_0 \in Y$ with rate λ , then c is a λ -submetric coupling at $(0, y_0)$.

In particular, if $Y = H(X)$, c is a metric coupling at $(0, Y)$.

A special case of interest is the choice for Y of the set of infima of finite families of continuous linear forms on X (see Rubinov, 2000, Rubinov, Shveidel, 2000).

EXAMPLE 7 (subaffine duality, Martínez-Legaz, 1988; Penot, Volle, 1987, 1988, 1990)

Let X be a normed vector space with dual X^* and let $Y = X^* \times \mathbb{R}$ endowed with the sup norm; let c be given by $c(x, y) = x^*(x) \wedge s$ for $y := (x^*, s)$, where $r \wedge s := \min(r, s) = \frac{1}{2}(r + s - |r - s|)$. As

$$\begin{aligned} |r \wedge s - r' \wedge s'| &\leq \frac{1}{2} |r - r' + s - s'| + \frac{1}{2} ||r - s| - |r' - s'| \\ &\leq \frac{1}{2} |r - r'| + \frac{1}{2} |s - s'| + \frac{1}{2} |r - s - r' + s'| \\ &\leq |r - r'| + |s - s'| \end{aligned}$$

and as

$$|x^*(x) \wedge s - x^*(x') \wedge s| \leq \|x^*\| \|x - x'\| \leq \|y\| \|x - x'\|,$$

we see that c induces a submetric coupling at (X, y_0) , where $y_0 = (0, 0)$. The Hahn-Banach theorem ensures that it is a 1-super-metric coupling at $(0, y_0)$ when X is endowed with the metric given by $d(x, x') := \|x - x'\| \wedge 1$. It is not the case for the distance associated with the norm.

EXAMPLE 8 (discrete measures)

Given a set E , let X be a subfamily of the set $\mathcal{P}_f(E)$ of finite subsets of E and let Y be a subspace of the space of bounded functions on E , endowed with the sup norm. Let us set for $x, x' \in X$, $y \in Y$,

$$d(x, x') := \#(x \Delta x'), \quad c(x, y) := \sum_{e \in x} y(e)$$

where $\#S$ denotes the number of elements of the finite set S and $x \Delta x' := (x \setminus x') \cup (x' \setminus x)$. One easily checks that d defines a metric on X and that, for $x, x' \in X$, $y, y' \in Y$, one has

$$c(x, y) - c(x', y) - c(x, y') + c(x', y') \leq \sum_{e \in x \Delta x'} |y - y'| (e) \leq d(x, x') \|y - y'\|,$$

while for $x, x' \in X$, $y \in Y$, one has

$$c(x, y) - c(x', y) - c(x, y') + c(x', y') = \sum_{e \in x \Delta x'} |y - y'| (e) \geq rd(x, x')$$

for y' given by $y'(e) := y(e)$ for $e \in S \setminus x \Delta x'$, $y'(e) := y(e) + r$ for $e \in x' \setminus x$, $y'(e) := y(e) - r$ for $e \in x \setminus x'$, so that c is a metric coupling. This example is important for discrete optimization.

EXAMPLE 9 (homotone duality)

Let (X, \leq) be an ordered metric space and let Y be a subset of the set of homotone functions on X , a function f being homotone (or isotone or nondecreasing) if for any $x, x' \in X$ with $x \leq x'$ one has $f(x) \leq f(x')$. Given $x_0 \in X$ let us set for $y, y' \in Y$ $d(y, y') := \sup\{r^{-1} |y(x) - y'(x)| : x \in B(x_0, r), r > 0\}$. Then the evaluation mapping is a submetric coupling at (x_0, Y) . The special case when $X = \mathbb{R}^d$, Y being the set of superadditive homotone functions null at 0 is considered in Wolsey (1981). The special case when X is a distributive lattice and Y is formed of modular or submodular functions is also important (see Fujishige (1984), for instance).

EXAMPLE 10 (shady conjugacy)

Let X be a n.v.s. and let Y be its dual. Let c^\vee and c^∇ be the couplings given by $c^\vee(x, y) = -\iota_{[y \leq 1]}(x)$ and $c^\nabla(x, y) = -\iota_{[y < 1]}(x)$, where $[y \leq 1] := \{x \in$

$X : y(x) \leq 1$, $[y < 1] := \{x \in X : y(x) < 1\}$ and where, for a subset S of X , ι_S is the indicator function of S given by $\iota_S(x) = 0$ for $x \in S$, $\iota_S(x) = +\infty$ for $x \in X \setminus S$. Then c^\vee and c^∇ are submetric couplings at $(0, 0)$. See Penot (1997, 2000, 2001), Thach (1991, 1993, 1994, 1995), Volle (1985).

EXAMPLE 11 (partial sublevel duality)

Let X be a n.v.s. and let $Y := X^* \times \mathbb{R}_-$. Let c^Σ be given by $c^\Sigma(x, (x^*, r)) = -\iota_{[x^* \geq r]}(x)$. Then c^∇ is a submetric coupling at $(0, (0, r))$ for each $r \in \mathbb{R}_-$. Similarly, if $Y := X^* \times (-\mathbb{P})$, where \mathbb{P} is the set of positive numbers and if $c^S(x, (x^*, r)) = -\iota_{[x^* > r]}(x)$, then c^S is a submetric coupling at $(0, (0, r))$ for each $r \in \mathbb{P}$ (see Penot, 2000).

EXAMPLE 12 (augmented duality)

Given a coupling $c : X \times Y \rightarrow \overline{\mathbb{R}}$ and a function $k : X \rightarrow \mathbb{R}$, let \widehat{c} be given by $\widehat{c}(x, y) := c(x, y) + k(x)$. Then one easily checks that \widehat{c} is a metric coupling iff c is a metric coupling. The case in which X is a Hilbert space and $k(x) := \frac{r}{2} \|x\|^2$ for some $r > 0$ is known to be of special importance (Janin, 1973; Rockafellar, 1974; Penot, Volle, 1988; Poliquin, 1992; Eberhard, Nyblom, 1998, etc.).

Other examples are given in Rubinov (2000), Rubinov, Glover (1997, 1998, 1999), Rubinov, Andramonov and Penot (1999, 2000, 2001), Rubinov, Shveidel (2000), Rubinov, Simsek (1995), Schwartz (1973, 1980), Singer (1986, 1987, 1997), for instance.

3. Coercivity

Hereafter X and Y are metric spaces, c is a coupling between X and Y . In order to illustrate the possible uses of nonlinear dualities, we intend to study the interplay between boundedness, conjugacy and coercivity. This can be done in a quantitative (and simple) way (see also Cominetti, 1994, Fougères, 1977, Penot, 1995, for the classical case).

Let us say that a function f on a metric space X is *coercive* if there exists $x_0 \in X$ such that $f(x) \rightarrow \infty$ as $d(x, x_0) \rightarrow \infty$. This property is obviously independent of the point x_0 : it amounts to the requirement that for each $r \in \mathbb{R}$ the sublevel set $[f \leq r]$ is bounded. We say that f is *super-coercive* if there exists $x_0 \in X$ such that $c_\infty(f) := \liminf_{d(x, x_0) \rightarrow \infty} f(x)/d(x, x_0) > 0$. We call $c_\infty(f)$ the *coercivity rate* of f . It does not depend on the choice of x_0 . Clearly, when $c_\infty(f)$ is finite, it is the supremum of the real numbers b such that $f(x) \geq bd(x, x_0)$ for $d(x, x_0)$ large enough. Note that, if f is bounded below on balls, then $c_\infty(f)$ is also the supremum of the real numbers b such that for some real number a one has $f(x) \geq bd(x, x_0) - a$ for each $x \in X$. Let us say that f is *hyper-coercive* if $f(x)/d(x, x_0) \rightarrow \infty$ as $d(x, x_0) \rightarrow \infty$, or, equivalently, when $c_\infty(f) = +\infty$.

In the following two statements we suppose X and Y are provided with base points x_0 and y_0 respectively and that $c(\cdot, y_0) = 0$. This last assumption

is satisfied in several of the examples displayed above. It can be obtained by replacing c by \widehat{c} given by $\widehat{c}(x, y) := c(x, y) - c(x, y_0)$.

PROPOSITION 3.1 (a) *Suppose the coupling c is a λ -submetric coupling at (x_0, y_0) and $c(\cdot, y_0) = 0$. If for some $a \in \mathbb{R}, b > 0$, a function f on X satisfies $f(x) \geq bd(x, x_0) - a$ for each $x \in X$, then $g := f^c$ is bounded above by $a + a'$ on the ball $B(y_0, \lambda^{-1}b)$ provided $c(x_0, \cdot)$ is bounded above by a' on this ball.*

(b) *Suppose the coupling c is a κ -super-metric coupling at (x_0, y_0) and $c(\cdot, y_0) = 0$. If g is a function on Y which is bounded above by a on a ball $B(y_0, b)$ for some $a \in \mathbb{R}, b > 0$, then $f := g^c$ satisfies $f(x) \geq \kappa bd(x, x_0) - a + a''$ for each $x \in X$ provided $c(x_0, \cdot)$ is bounded below by a'' on this ball. In particular f is super-coercive and its coercivity rate c_∞ is such that $c_\infty \geq \kappa b$.*

When $\kappa = \lambda = 1$ as in the classical case, we get a dual interpretation of the coercivity rate of a function bounded below on balls with center x_0 as the supremum of the radius of the balls centered at y_0 on which the conjugate function is bounded above.

Proof. (a) Let f be such that for some $a \in \mathbb{R}, b > 0$, one has $f(x) \geq bd(x, x_0) - a$ for each $x \in X$. If the coupling c is a λ -submetric coupling at (x_0, y_0) , and if $c(x_0, \cdot)$ is bounded above by a' on $B(y_0, \lambda^{-1}b)$ then for $y \in B(y_0, \lambda^{-1}b)$ one has

$$\begin{aligned} f^c(y) &\leq \sup_{x \in X} (c(x, y) - bd(x, x_0) + a) \\ &\leq \sup_{x \in X} (c(x_0, y) + \lambda d(x, x_0)d(y, y_0) - bd(x, x_0) + a) \leq a + a'. \end{aligned}$$

(b) Let $g : Y \rightarrow \overline{\mathbb{R}}$ be bounded above by a on a ball $B(y_0, b)$ for some $a \in \mathbb{R}, b > 0$. If the coupling c is a κ -super-metric coupling at (x_0, y_0) and $c(x_0, \cdot)$ is bounded below by a'' on $B(y_0, b)$, then for each $x \in X$, one has

$$\begin{aligned} g^c(x) &\geq \sup_{y \in B(y_0, b)} c(x, y) - a \\ &\geq \sup_{y \in B(y_0, b)} (c(x, y) - c(x_0, y) + a'') - a \geq \kappa bd(x, x_0) + a'' - a. \end{aligned}$$

■

REMARK 3.1 The preceding proof shows that $f^c - c(x_0, \cdot)$ is bounded above by a on the ball $B(y_0, \lambda^{-1}b)$ whenever f satisfies $f \geq bd(\cdot, x_0) - a$ and the coupling c is such that $c(x, y) - c(x_0, y) \leq \lambda d(x, x_0)d(y, y_0)$ for any $x \in X, y \in Y$. This condition is satisfied if c is a λ -submetric coupling at (x_0, y_0) such that $c(\cdot, y_0) = 0$. Similarly, $g^c \geq \kappa bd(\cdot, x_0) - a$ whenever $g - c(x_0, \cdot)$ is bounded above by a on the ball $B(y_0, b)$ and the coupling c is such that $\sup_{y \in B(y_0, b)} (c(x, y) - c(x_0, y)) \geq \kappa bd(x, x_0)$ for any $x \in X$. This condition is satisfied if c is a κ -super-metric coupling at (x_0, y_0) such that $c(\cdot, y_0) = 0$.

COROLLARY 3.1 (a) *Suppose the coupling c is a λ -submetric coupling at (x_0, y_0) , $c(\cdot, y_0) = 0$ and $c(x_0, \cdot)$ is bounded above on balls. If a function f on X is hyper-coercive and bounded below on bounded sets, then f^c is bounded above on bounded sets of Y .*

(b) *Suppose the coupling c is a κ -super-metric coupling at (x_0, y_0) , $c(\cdot, y_0) = 0$ and $c(x_0, \cdot)$ is bounded below on balls. If a function g on Y is bounded above on bounded sets of Y , then g^c is hyper-coercive.*

4. From rotundity properties to smoothness properties

In this section we suppose a function f on X satisfies a lower estimate and we get an upper estimate on its conjugate. Several of the estimates we deal with involve functions of the form

$$\sigma_{T, x_0}(x) := \sup_{v \in T} (c(x, v) - c(x_0, v))$$

for some $x_0 \in X$ and some $T \subset Y$. When X has a group structure and $c(\cdot, y_0)$ is additive for each $y_0 \in T$, this term is just the support function of T evaluated at $x - x_0$: $\sigma_{T, x_0}(x) = \sigma_T(x - x_0)$ with

$$\sigma_T(u) := \sup_{v \in T} c(u, v).$$

Furthermore, if c is the classical duality, X being a normed space, if f is convex and continuous at x_0 and if $T = \partial f(x_0) := \{y_0 \in X^* : \langle y_0, \cdot \rangle \leq f(x_0 + \cdot) - f(x_0)\}$, one has $\sigma_T(x - x_0) = f'(x_0, x - x_0)$, the lower Hadamard derivative (or contingent derivative) of f at x_0 given by

$$f'(x_0, u) := \liminf_{(t, v) \rightarrow (0+, u)} t^{-1} (f(x_0 + tv) - f(x_0))$$

for $u \in X$.

We start with a variant of Theorem 3.2 of Rolewicz (1994), an extension of the famous Asplund (1968) and Brønsted (1964) theorem; see also Azé, Penot (1995), Proposition 3.5.

PROPOSITION 4.1 *Given $x_0 \in \text{dom} f$, $y_0 \in Y$ such that c is a λ -submetric coupling at (x_0, y_0) , let $\gamma : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be such that*

$$f(x) \geq f(x_0) + c(x, y_0) - c(x_0, y_0) + \gamma(d(x, x_0)) \quad \forall x \in X. \quad (6)$$

Then one has

$$f^c(y) \leq f^c(y_0) + c(x_0, y) - c(x_0, y_0) + \gamma^*(\lambda d(y, y_0)) \quad \forall y \in Y. \quad (7)$$

When γ is a starshaped gage, relation (6) can be interpreted as a reinforced subdifferentiability property or a rotundity property; then, by Lemma 2.1, γ^* is

an hypermodulus, so that relation (7) is an approximation property. When Y is a normed vector space, $x_0 \in \partial^c f^c(y_0)$ and $y \mapsto c(x_0, y) - c(x_0, y_0)$ is linear and continuous, this last relation is a smoothness property which amounts to Fréchet super-differentiability of f^c at y_0 . Moreover, when $c(\cdot, y_0) = 0$, and $\gamma(t) > 0$ for $t > 0$, relation (6) is a conditioning property of the minimizer x_0 .

The preceding statement is a consequence of the following result which is more technical but more versatile (take $S = \{x_0\}$, $T = \{y_0\}$).

LEMMA 4.1 *Given $S \subset \text{dom} f$, $T \subset Y$ such that c is a λ -submetric coupling at (S, T) and $\gamma : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ such that*

$$f(x) \geq \inf_{x_0 \in S} \left[f(x_0) + \inf_{y_0 \in T} (c(x, y_0) - c(x_0, y_0)) + \gamma(d(x, x_0)) \right] \quad \forall x \in X \quad (8)$$

one has

$$f^c(y) \leq \sup_{y_0 \in T} \left[f^c(y_0) + \sup_{x_0 \in S} (c(x_0, y) - c(x_0, y_0)) + \gamma^*(\lambda d(y, y_0)) \right] \quad \forall y \in Y. \quad (9)$$

Proof. Let $y \in Y$. When $f^c(y) = -\infty$ inequality (9) is obvious. Let $r \in \mathbb{R}$, $r < f^c(y)$. We can find $x \in X$ such that $-r > f(x) - c(x, y)$. Then $f(x) < +\infty$, $c(x, y) > -\infty$ and $f(x) < c(x, y) - r$. By (8) we can find $(x_0, y_0) \in S \times T$ such that

$$c(x, y) - r > f(x_0) - c(x_0, y_0) + c(x, y_0) + \gamma(d(x, x_0)).$$

By the relations $f(x_0) - c(x_0, y_0) \geq -f^c(y_0)$, $\gamma(s) - st \geq -\gamma^*(t)$ we get

$$\begin{aligned} -r &> -f^c(y_0) + c(x, y_0) - c(x, y) + \gamma(d(x, x_0)) \\ &\geq -f^c(y_0) + c(x_0, y_0) - c(x_0, y) - \lambda d(x, x_0)d(y, y_0) + \gamma(d(x, x_0)) \\ &\geq -f^c(y_0) + c(x_0, y_0) - c(x_0, y) - \gamma^*(\lambda d(y, y_0)). \end{aligned}$$

Since we may assume that $f^c(y_0) < +\infty$, hence that $f^c(y_0)$ is finite, reversing signs and taking the supremum over $(x_0, y_0) \in S \times T$ and over r , we obtain inequality (9). \blacksquare

We have seen that when S and T are singletons, one obtains the preceding proposition. If only one of the subsets S or T is a singleton, one also obtains interesting consequences. When $S = \{x_0\}$, relation (8) is a kind of calmness condition. Let us give a statement when T is a singleton $\{y_0\}$.

COROLLARY 4.1 *Given $S \subset \text{dom} f$, $y_0 \in Y$ such that c is a λ -submetric coupling at (S, y_0) and $\gamma : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ such that*

$$f(x) \geq \inf_{x_0 \in S} [f(x_0) + c(x, y_0) - c(x_0, y_0) + \gamma(d(x, x_0))] \quad \forall x \in X \quad (10)$$

one has

$$f^c(y) \leq f^c(y_0) + \sup_{x_0 \in S} [c(x_0, y) - c(x_0, y_0)] + \gamma^*(\lambda d(y, y_0)) \quad \forall y \in Y. \quad (11)$$

When X and Y are normed vector spaces, c is the Fenchel coupling and γ is a starshaped gage, relation (10) is satisfied when f is rotund at some $x_0 \in S$ and then relation (11) is a conical approximation property.

Taking for S a set of minimizers, and setting $d_S(x) := \inf\{d(x, w) : w \in S\}$ one gets a consequence of a conditioning property in the sense of Penot (1998).

COROLLARY 4.2 *Suppose $c(\cdot, y_0) = 0$, γ is nondecreasing and let S be a non-empty subset of the set of minimizers of f . Let $m := \inf f(X)$. Suppose that one has*

$$f(x) \geq m + \gamma(d_S(x)) \quad \forall x \in X.$$

Then, if c is a λ -metric coupling at (S, y_0) , setting $\sigma_S(y) := \sup_{x_0 \in S} c(x_0, y)$, one has

$$f^c(y) \leq f^c(y_0) + \sigma_S(y) + \gamma^*(\lambda d(y, y_0)) \quad \forall y \in Y.$$

Now let us give a variant of Lemma 4.1 which will yield other statements generalizing results in Azé (1999), Azé, Rahmouni (1994, 1995, 1996).

LEMMA 4.2 *Suppose c is a λ -submetric coupling at (S, T) with $S \subset \text{dom} f$, $T \subset Y$. Suppose that for some nondecreasing function $\gamma : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ one has for each $x \in X$*

$$f(x) \geq \sup_{x_0 \in S} \left[f(x_0) + \sup_{y_0 \in T} (c(x, y_0) - c(x_0, y_0)) \right] + \gamma(d_S(x)). \quad (12)$$

Then, for each $y \in Y$, one has

$$f^c(y) \leq \sup_{x_0 \in S} (-f(x_0) + c(x_0, y)) + \gamma^*(\lambda d_T(y)). \quad (13)$$

Moreover, if for some subset Z of Y one has $z \in \partial^c f(x_0)$ for any $x_0 \in S$, $z \in Z$, then, for each $y \in Y$ one has

$$\begin{aligned} f^c(y) &\leq \sup_{x_0 \in S} \left(\inf_{z \in Z} (f^c(z) - c(x_0, z)) + c(x_0, y) \right) + \gamma^*(\lambda d_T(y)) \\ &\leq \inf_{z \in Z} \left(f^c(z) + \sup_{x_0 \in S} (c(x_0, y) - c(x_0, z)) \right) + \gamma^*(\lambda d_T(y)). \end{aligned}$$

Proof. Given $y \in Y$ such that $f^c(y) > -\infty$, let $r \in \mathbb{R}$, $r < f^c(y)$. We can find $x \in X$ such that $r < c(x, y) - f(x)$. Taking (12) into account, for any $x_0 \in S$, $y_0 \in T$, the property of the coupling yields

$$\begin{aligned} r &< -f(x_0) + c(x, y) - c(x, y_0) + c(x_0, y_0) - \gamma(d_S(x)) \\ r &< -f(x_0) + c(x_0, y) + \lambda d(x, x_0) d(y, y_0) - \gamma(d_S(x)). \end{aligned}$$

Taking the infimum over $y_0 \in T$, and, for a given $\varepsilon > 0$, picking some $x_0 \in S$ such that $d_S(x) \geq d(x, x_0) - \varepsilon$ we get

$$r < -f(x_0) + c(x_0, y) + \lambda d(x, x_0) d_T(y) - \gamma(d(x, x_0) - \varepsilon).$$

Setting $s := d(x, x_0) - \varepsilon$, $t := \lambda d_T(y)$ and observing that $st - \gamma(s) \leq \gamma^*(t)$, we get

$$r \leq -f(x_0) + c(x_0, y) + \gamma^*(\lambda d_T(y)) + \lambda \varepsilon d_T(y).$$

Taking the supremum over $x_0 \in S$ and the infimum over $\varepsilon > 0$ we obtain

$$r \leq \sup_{x_0 \in S} (-f(x_0) + c(x_0, y)) + \gamma^*(\lambda d_T(y)).$$

Since r is arbitrarily close to $f^c(y)$, relation (13) follows. The second assertion is a consequence of (13) and of the Fenchel equality $-f(x_0) = f^c(z) - c(x_0, z)$ for each $z \in Z$. ■

Simplified statements can be given when S , T or Z are singletons or when one takes $Z = T$. This last choice can be justified by the fact that, whenever γ takes nonnegative values, in particular when $\gamma(0) = 0$, relation (12) implies that $T \subset \partial^c f(x_0)$ for each $x_0 \in S$. Let us give a statement for this special case.

PROPOSITION 4.2 *Suppose c is a λ -submetric coupling at (S, T) with $S \subset \text{dom} f$, $T \subset Y$. Suppose that for some nondecreasing function $\gamma : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}_+$ relation (12) holds. Then, for each $y_0 \in T$, one has for any $y \in Y$*

$$f^c(y) \leq f^c(y_0) + \sup_{x_0 \in S} (c(x_0, y) - c(x_0, y_0)) + \gamma^*(\lambda d_T(y)).$$

The case S is a singleton is stated in the following corollary.

COROLLARY 4.3 *Suppose c is a λ -submetric coupling at (x_0, T) for some $x_0 \in X$, $T \subset Y$. Let $f : X \rightarrow \overline{\mathbb{R}}$ be an arbitrary function with $x_0 \in \text{dom} f$, $T, Z \subset \partial^c f(x_0)$. Suppose that for some nondecreasing function $\gamma : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ one has for each $x \in X$*

$$f(x) \geq f(x_0) + \sup_{y_0 \in T} (c(x, y_0) - c(x_0, y_0)) + \gamma(d(x, x_0)). \tag{14}$$

Then, for each $y \in Y$ one has

$$f^c(y) \leq \inf_{z \in Z} (f^c(z) - c(x_0, z) + c(x_0, y)) + \gamma^*(\lambda d_T(y)). \tag{15}$$

Relation (14) can be interpreted as a kind of firm subdifferentiability property as it can be written

$$f(x) \geq f(x_0) + \sigma_{T, x_0}(x) + \gamma(d(x, x_0)).$$

When Y is a normed space and $c(x_0, \cdot)$ is additive, relation (15) yields, for each $y_0 \in T$

$$\begin{aligned} f^c(y) &\leq f^c(y_0) + c(x_0, y - y_0) + \gamma^*(\lambda d_T(y)) \\ &\leq f^c(y_0) + c(x_0, y - y_0) + \gamma^*(\lambda \|y - y_0\|), \end{aligned}$$

which says that $f^* := f^c$ is Fréchet super-differentiable at y_0 , uniformly on T , provided γ is a starshaped gage, as then γ^* is an hyper-modulus by Lemma 2.1.

With the same choice for c , relation (12) can be interpreted as a form of a firm subdifferentiability property, uniform on S ; relation (15) yields a conical upper approximation of f around z_0 when $z_0 \in T$. Such an approximation property has been studied by a number of researchers starting from Mignot (see Dontchev, Hager, 1994, Mignot, 1976, Pang, 1990 and 1995, Penot, 1982, Robinson, 1991, at least in its two-sided version, for the one-sided version used here see Agadi, Penot, 1996).

Now let us take for Z a singleton $\{z_0\}$ in Lemma 4.2.

COROLLARY 4.4 *Suppose $c, \lambda, f, \gamma, S \subset \text{dom}f, T \subset Y$ are as in Lemma 4.2 and relation (12) holds. Suppose that $z_0 \in Y$ is such that $z_0 \in \partial^c f(x_0)$ for each $x_0 \in S$. Then, for each $y \in Y$, one has*

$$f^c(y) \leq f^c(z_0) + \sup_{x_0 \in S} (c(x_0, y) - c(x_0, z_0)) + \gamma^*(\lambda d_T(y)).$$

Taking $T = \{y_0\} = \{z_0\}$, we obtain a further simplification.

COROLLARY 4.5 *Suppose $c, \lambda, \gamma, S \subset \text{dom}f$ are as above and let $y_0 \in \partial^c f(x_0)$ for each $x_0 \in S$. Suppose c is a λ -submetric coupling at (S, y_0) and that for each $x \in X$ one has*

$$f(x) \geq \sup_{x_0 \in S} (f(x_0) + c(x, y_0) - c(x_0, y_0)) + \gamma(d_S(x)).$$

Then, for each $y \in Y$, one has

$$f^c(y) \leq f^c(y_0) + \sup_{x_0 \in S} (c(x_0, y) - c(x_0, y_0)) + \gamma^*(\lambda d(y, y_0)).$$

5. From upper differentiability properties to rotundity properties

We now study the consequences of a reverse estimate: we suppose a function g on Y satisfies an upper estimate and we get a lower estimate on its conjugate. In view of the symmetry of our framework it would be possible to perform such a study on a function f on X . However, our choice is dictated by the hope to get a converse of the results of the preceding section.

PROPOSITION 5.1 *Let g be a given function on Y and let $\omega : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be nondecreasing. Suppose that for some subset T of Y and some $x_0 \in \bigcap \{\partial^c g(y_0) : y_0 \in T\}$ the following estimate holds for any $y_0 \in T$ and any $y \in Y$*

$$g(y) \leq g(y_0) + c(x_0, y) - c(x_0, y_0) + \omega(d(y, y_0)). \tag{16}$$

If for each $y_0 \in T$ the coupling c is a κ -super-metric coupling at (x_0, y_0) , then the conjugate g^c of g satisfies

$$g^c(x) \geq g^c(x_0) + \sup_{y_0 \in T} (c(x, y_0) - c(x_0, y_0)) + \omega^*(\kappa d(x, x_0)). \tag{17}$$

We observe that when ω is an hyper-modulus, relation (16) can be interpreted as a super-differentiability property at y_0 . Then, relation (17) is a rotundity property. We also note that the preceding result can be obtained from the case $T = \{y_0\}$. We deduce it from the following statement by taking $S := \{x_0\}$.

PROPOSITION 5.2 *Let g be a given function on Y and let $\omega : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be nondecreasing. Suppose that for some subset T of Y and some $S \subset \bigcap \{\partial^c g(y_0) : y_0 \in T\}$ the following estimate holds for any $y_0 \in T$*

$$g(y) \leq g(y_0) + \sup_{x_0 \in S} (c(x_0, y) - c(x_0, y_0)) + \omega(d(y, y_0)) \quad \forall y \in Y. \tag{18}$$

If c is a κ -super-metric coupling at (S, T) , then the conjugate g^c of g satisfies

$$g^c(x) \geq \sup_{x_0 \in S} \left[g^c(x_0) + \sup_{y_0 \in T} (c(x, y_0) - c(x_0, y_0)) \right] + \omega^*(\kappa d_S(x)) \quad \forall x \in X. \tag{19}$$

Proof. Let $y_0 \in T$, $x \in X$ and let $t > g^c(x)$, $t \in \mathbb{R}$. For each $y \in Y$ we can find $s_y < g(y)$ such that $t > c(x, y) - s_y$ and $x_y \in S$ such that $s_y < g(y_0) + c(x_y, y) - c(x_y, y_0) + \omega(d(y, y_0))$, hence

$$c(x, y) - s_y \geq -g(y_0) + c(x, y) - c(x_y, y) + c(x_y, y_0) - \omega(d(y, y_0)).$$

Since c is a κ -super-metric coupling at (x_y, y_0) and since ω is nondecreasing, taking the supremum over the ball $B(y_0, r)$ for a given $r \geq 0$, we get

$$\begin{aligned} \sup_{y \in B(y_0, r)} (c(x, y) - s_y) - c(x, y_0) &\geq -g(y_0) + \kappa r d(x, x_y) - \omega(r), \\ t - c(x, y_0) &\geq -g(y_0) + \kappa r d_S(x) - \omega(r). \end{aligned}$$

Given $x_0 \in S$, using the relation $-g(y_0) = -c(x_0, y_0) + g^c(x_0)$, and taking the supremum over $r \in \mathbb{R}_+$, we get

$$t \geq g^c(x_0) - c(x_0, y_0) + c(x, y_0) + \sup_{r \geq 0} (\kappa r d_S(x)) - \omega(r).$$

Taking the supremum over $y_0 \in T$ and the infimum over $t > g^c(x)$, we obtain the result from the very definition of ω^* . ■

Note that assumption (16) in Proposition 5.1 corresponds to the conclusion of Proposition 4.1 with $g = f^c$, $\omega = \gamma^*$, $\lambda = 1$ while the conclusion (17) corresponds to assumption (6) in Proposition 4.1 when $\kappa = 1$, $f = g^c$ and $\gamma = \omega^*$. Therefore we have obtained the following characterization.

THEOREM 5.1 *Let $f : X \rightarrow \overline{\mathbb{R}}$ and let $\gamma : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be nondecreasing and such that $\gamma^{**} = \gamma$. Given $x_0 \in \text{dom} f$, $y_0 \in \partial^c f(x_0)$ such that c is a metric coupling at (x_0, y_0) , the following two relations are equivalent:*

$$f(x) \geq f(x_0) + c(x, y_0) - c(x_0, y_0) + \gamma(d(x, x_0)) \quad \forall x \in X. \quad (20)$$

$$f^c(y) \leq f^c(y_0) + c(x_0, y) - c(x_0, y_0) + \gamma^*(d(y, y_0)) \quad \forall y \in Y. \quad (21)$$

Similarly, from Proposition 5.2 one can get a characterization of well-conditioning. We first state a sufficient condition.

COROLLARY 5.1 *Let f be a given function on X such that $f = f^{cc}$ and let $\omega : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be nondecreasing. Suppose there exists some element y_0 of Y such that $c(\cdot, y_0) = 0$. Suppose that for some subset S of the set of minimizers of f the following estimate holds*

$$f^c(y) \leq f^c(y_0) + \sup_{x_0 \in S} c(x_0, y) + \omega(d(y, y_0)) \quad \forall y \in Y. \quad (22)$$

If c is a κ -super-metric coupling at (S, y_0) , then

$$f(x) \geq \inf f(X) + \omega^*(\kappa d_S(x)). \quad (23)$$

Proof. Let us observe that, setting $g := f^c$, we have $S \subset \partial^c g(y_0)$ as $g(y_0) = -\inf f(X)$ because $c(\cdot, y_0) = 0$ and as for any $x_0 \in S$, $y \in Y$ we have $g^c(x_0) = f(x_0) = \inf f(X)$. Then the conclusion follows from Proposition 5.2 with $T := \{y_0\}$. ■

Gathering this sufficient condition with the necessary condition of Corollary 4.2 we get the following characterization of well-conditioning.

THEOREM 5.2 *Let f be a given function on X such that $f = f^{cc}$ and let $\gamma : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be a gage such that $\gamma^{**} = \gamma$. Suppose there exists some element y_0 of Y such that $c(\cdot, y_0) = 0$ and c is a metric coupling at (S, y_0) where S is the set of minimizers of f . Then f satisfies the estimate*

$$f(x) \geq \inf f(X) + \gamma(d_S(x)) \quad \forall x \in X$$

iff

$$f^c(y) \leq f^c(y_0) + \sup_{x_0 \in S} c(x_0, y) + \gamma^*(d(y, y_0)) \quad \forall y \in Y. \quad (24)$$

Let us end our study with the following corollary generalizing part of Azé, Rahmouni (1995), Lemma 2. It deals with the computation of the conjugate of a function of the form $g := \varphi \circ d_T$, where T is some subset of Y .

COROLLARY 5.2 *Suppose there exists a point $x_0 \in X$ such that $c(x_0, \cdot) = 0$ and the coupling c is a metric coupling at (x_0, y_0) for each $y_0 \in T$, where T is some subset of Y . Let $\varphi : \mathbb{R}_+ \rightarrow \overline{\mathbb{R}}$ be nondecreasing and such that $\varphi(0) = 0$. Then, for σ_T and g given by $\sigma_T(x) := \sup_{y_0 \in T} c(x, y_0)$, $g(y) = \varphi(d_T(y))$, one has*

$$g^c(x) = \sigma_T(x) + \varphi^*(d(x, x_0)).$$

In particular, one has $(d_T)^c(x) = \sigma_T(x)$ for $x \in B(x_0, 1)$, else $+\infty$.

Proof. Since $c(x_0, \cdot) = 0$ and since for any $y_0 \in T$, $y \in Y$, one has $g(y_0) = \varphi(0) = 0 \leq \varphi(d_T(y)) = g(y)$, so that $x_0 \in \partial^c g(y_0)$ and $g^c(x_0) = 0$. Thus, Proposition 5.1 with $\omega = \varphi$ yields

$$g^c(x) \geq g^c(x_0) + \sup_{y_0 \in T} c(x, y_0) + \varphi^*(d(x, x_0)) = \sigma_T(x) + \varphi^*(d(x, x_0)).$$

On the other hand, by substituting Y, X, T, g, φ, x_0 , to X, Y, S, f, γ, z_0 in Corollary 4.4 and observing that

$$g(y) \geq \sup_{y_0 \in T} (g(y_0) + c(x_0, y) - c(x_0, y_0)) + \varphi(d_T(y)),$$

we get

$$g^c(x) \leq g^c(x_0) + \sup_{y_0 \in T} c(x, y_0) + \varphi^*(d(x, x_0)) = \sigma_T(x) + \varphi^*(d(x, x_0)).$$

The last assertion corresponds to the case $\varphi(t) = |t|$ for which $\varphi^* = \iota_{[0,1]}$, the indicator function of $[0, 1]$ whose value is 0 on $[0, 1]$ and $+\infty$ elsewhere. ■

It is shown in Azé, Rahmouni (1994, 1995, 1996), Penot (1995), that well-conditioning of f is linked with a continuity property of the subdifferential of the conjugate function of f and with a growth property of the subdifferential of f . These questions will not be considered here.

References

- AGADI, A. and PENOT, J.-P. (1996) *A comparative study of various notions of approximation of sets*. Preprint, Univ. of Pau.
- ASPLUND, E. (1968) Fréchet differentiability of convex functions. *Acta Math* **121**, 31–47.
- ASPLUND, E. and ROCKAFELLAR, R.T. (1969) Gradients of convex functions. *Trans. Amer. Math. Soc.* **139**, 443–467.

- ATTEIA, M. and ELQORTOBI, A. (1981) Quasi-convex duality. In: A. Auslender et al., eds., *Optimization and Optimal Control, Proc. Conference Oberwolfach March 1980*. Lecture Notes in Control and Inform. Sci. **30**, Springer-Verlag, Berlin, 3–8.
- AUBIN, J.-P. (1999) *Mutational and Morphological Analysis*. Birkhäuser, Basel.
- AZÉ, D. (1997) *Éléments d'analyse convexe et variationnelle*. Ellipses, Paris.
- AZÉ, D. (1999) On the remainder of the first order development of convex functions. *Ann. Math. Québec* **23**, 1–14.
- AZÉ, D., CORVELLEC, J.-N. and LUCCHETTI, R. (2002) Variational pairs and applications to stability in nonsmooth analysis. *Nonlinear Anal.* **49**, 643–670.
- AZÉ, D. and PENOT, J.-P. (1995) Uniformly convex and uniformly smooth convex functions. *Ann. Fac. Sci. Toulouse* **4**, 705–730.
- AZÉ, D. and RAHMOUNI, A. (1994) Lipschitz behavior of the Legendre-Fenchel transform. *Set-Valued Anal.* **2** (1-2), 35–48.
- AZÉ, D. and RAHMOUNI, A. (1995) Intrinsic bounds for Kuhn-Tucker points of perturbed convex programs. In: R. Durier and C. Michelot, eds., *Recent developments in optimization, Seventh French-German Conference on Optimization*. Lecture Notes in Economics and Math. **429**, Springer Verlag, Berlin, 17–35.
- AZÉ, D. and RAHMOUNI, A. (1996) On primal dual stability in convex optimization. *J. Convex Anal.* **3**, 309–329.
- BALDER, E.J. (1977) An extension of duality-stability relations to non-convex optimization problems. *SIAM J. Control Opt.* **15**, 329–343.
- BEAUZAMY, B. (1992) *Introduction to Banach Spaces and their Geometry*. Math. Studies **68**, North-Holland, Amsterdam.
- BRONSTEDT, A. (1964) Conjugate convex functions on topological vector spaces. *Mat-Fys. Medel Danska Vod Selsk.* **2**.
- COMINETTI, R. (1994) Some remarks on convex duality in normed spaces with and without compactness. *Control and Cybern.* **23** (1-2), 123–138.
- CROUZEIX, J.-P. (1977) Contribution à l'étude des fonctions quasi-convexes. *Thèse d'Etat, Univ. de Clermont II*.
- DELLACHERIE C. and MEYER, P.A. (1975) *Probabilités et Potentiel*. Hermann, Paris.
- DEVILLE, R., GODEFROY, G. and ZIZLER, D. (1993) *Smoothness and renorming in Banach spaces*. Longman, Harlow.
- DIESTEL, J. (1975) *Geometry of Banach Spaces. Selected Topics*. Lecture Notes in Mathematics **485**, Springer Verlag.
- DIEWERT, W.E. (1982) Duality approaches to microeconomics theory. In: K.J. Arrow and M.D. Intriligator, eds., *Handbook of Mathematical Economics*, vol. 2, North Holland, Amsterdam, 535–599.
- DOLECKI, S. and KURCYUSZ, S. (1978) On Φ -convexity in extremal problems. *SIAM J. Control Optim.* **16**, 277–300.

- DONTCHEV, A. and HAGER, A. (1994) Implicit functions, Lipschitz maps and stability in optimization. *Math. Oper. Res.* **19**, 753–768.
- EBERHARD, A. and NYBLÖM, M. (1998) Jets, generalised convexity, proximal normality and differences of functions. *Nonlinear Anal.* **34**, 319–360.
- ELQORTOBI, A. (1992) Inf-convolution quasi-convexe des fonctionnelles positives. *Rech. Oper.* **26**, 301–311.
- ELQORTOBI, A. (1993) Conjugaison quasi-convexe des fonctionnelles positives. *Annales Sci. Math. Québec* **17** (2), 155–167.
- ELSTER, K.-H. and NEHSE, R. (1974) Zur theorie der polarfunktionale. *Math. Oper. Stat.* **5**, 3–21.
- ELWORTHY, K.D. (1975) Measures on infinite dimensional manifolds. In: *Functional Integration and Applications*. Clarendon Press, Oxford, 60–68.
- EVERS, J.J.M. and VAN MAAREN, H. (1981) *Duality principles in mathematics and their relations to conjugate functions*. Preprint, Univ. of Technology of Twente.
- FAN, K. (1963) On the Krein-Milman theorem. In: *Convexity*, Proc. Pure Math. **7**, Amer. Math. Soc. Providence, 211–220.
- FLACHS, J. and POLLATSCHKE, A.M. (1979) Duality theorems for certain programs involving minimum or maximum operations. *Math. Prog.* **16**, 348–370.
- FLORES-BAZÁN, F. (1995) On a notion of subdifferentiability for non-convex functions. *Optimization* **33**, 1–8.
- FLORES-BAZÁN, F. and MARTÍNEZ-LEGAZ, J.-E. (1998) Simplified global optimality conditions in generalized conjugacy theory. In: J.-P. Crouzeix, J.-E. Martínez-Legaz and M. Volle, eds., *Generalized Convexity, Generalized Monotonicity: Recent Results*. Kluwer, Dordrecht, 305–329.
- FOUGÈRES, A. (1977) Coercivité, convexité, relaxation: une extension naturelle du théorème d’inf-équitinuité de J.-J. Moreau. *C. R. Acad. Sci.*, Paris, Sér. A **285**, 711–713.
- FOUGÈRES, A. (1977) Propriétés géométriques et minoration des intégrandes convexes normales coercives. *C. R. Acad. Sci.*, Paris, Sér. A **284**, 873–876.
- FUJISHIGE, S. (1984) Theory of submodular programs: A Fenchel-type min-max theorem and subgradients of submodular functions. *Math. Programming* **29**, 142–155.
- GROMOV, M. (1999) *Metric Structures for Riemannian and Non-Riemannian Spaces*. Birkhäuser, Boston.
- HORST, R. and TUY, H. (1990) *Global Optimization*. Springer Verlag, Berlin.
- IOFFE, A.D. (2001) Abstract convexity and non-smooth analysis. *Adv. Math. Econ.* **3**, 45–61.
- IOFFE, A.D. (2001) Towards metric theory of metric regularity. In: M. Lassonde, ed., *Approximation, Optimization and Mathematical Economics*. Physica-Verlag, Heidelberg, 165–176.

- JANIN, R. (1973) Sur la dualité en programmation dynamique. *C.R. Acad. Sci.*, Paris, Sér. A **277**, 1195–1197.
- JOHN, R. (2001) A note on Minty variational inequalities and generalized monotonicity. In: N. Hadjisavvas, et al., eds., *Generalized convexity and generalized monotonicity. Proceedings of the 6th international symposium, Samos, Greece, September 1999*. Lect. Notes Econ. Math. Syst. **502**, Springer, Berlin, 240–246.
- LAFONTAINE, J. and PANSU, P. (1981) *Structures Métriques pour les Variétés Riemanniennes*. Cedic/Fernand Nathan, Paris.
- LAU, L.J. (1970) Duality and the structure of utility functions. *J. Econ. Theory*, **1**, 374–396.
- LEMAIRE, B. (1992) Bonne position, conditionnement, et bon comportement asymptotique. *Séminaire d'Analyse Convexe, Univ. of Montpellier* **22**, 5.1–5.12.
- LEMAIRE, B. (1995) Duality in reverse convex optimization. In: M. Sofonea and J.-N. Corvellec, eds., *Proceedings of the Second Catalan Days on Applied Mathematics*. Presses Universitaires de Perpignan, Perpignan, 173–182.
- LEMAIRE, B. and VOLLE, M. (1998) Duality in d.c. programming. In: J.-P. Crouzeix, J.-E. Martínez-Legaz and M. Volle, eds., *Generalized Convexity, Generalized Monotonicity: Recent Results*. Kluwer, Dordrecht, 331–342.
- MARTINEZ-LEGAZ, J.-E. (1988) On lower subdifferentiable functions. In: K.H. Hoffmann et al. eds., *Trends in Mathematical Optimization*. Series Numer. Math. **84**, Birkhauser, Basel, 197–232.
- MARTINEZ-LEGAZ, J.-E. (1988) Quasiconvex duality theory by generalized conjugacy methods. *Optimization* **19** (5), 603–652.
- MARTINEZ-LEGAZ, J.-E. (1990) Generalized conjugacy and related topics. In: A. Cambini, E. Castagnoli, L. Martein, P. Mazzoleni, S. Schaible, eds., *Generalized convexity and fractional programming with economic applications*. Lecture Notes in Econ. and Math. Systems **345**, Springer Verlag, Berlin, 168–197.
- MARTINEZ-LEGAZ, J.-E. (1991) Duality between direct and indirect utility functions under minimal hypothesis. *J. Math. Econ.* **20**, 199–209.
- MARTINEZ-LEGAZ, J.-E. (1995) Fenchel duality and related properties in generalized conjugacy theory. *Southeast Asian Bull. Math.* **19** (2), 99–106.
- MIGNOT, F. (1976) Contrôle dans les inégalités variationnelles elliptiques. *J. Funct. Anal.* **22**, 130–185.
- MOREAU, J.-J. (1970) Inf-convolution, sous-additivité, convexité des fonctions numériques. *J. Math. Pures et Appl.* **49**, 109–154.
- OETTLI, W. and SCHLÄGER, D. (1998) Conjugate functions for convex and nonconvex duality. *J. Glob. Optim.* **13** (4), 337–347.
- PALLASCHKE, D. and ROLEWICZ, S. (1997) Foundations of Mathematical Optimization: Convex Analysis without Linearity. *Mathematics and its Applications* **388**, Kluwer, Dordrecht.

- PANG, J.-S. (1990) Newton's method for B-differentiable equations. *Math. Oper. Res.* **15**, 311–341.
- PANG, J.-S. (1995) Necessary and sufficient conditions for solution stability of parametric nonsmooth equations. In: D.Z. Du et al., eds., *Recent Advances in Nonsmooth Optimization*. World Scientific, Singapore, 261–288.
- PASSY, U. and PRISMAN, E.Z. (1985) A convexlike duality scheme for quasi-convex programs. *Math. Programming* **32**, 278–300.
- PENOT, J.-P. (1982) Regularity conditions in mathematical programming. *Math. Prog. Study* **19**, 167–199.
- PENOT, J.-P. (1985) *Modified and augmented Lagrangian theory revisited and augmented*. Unpublished lecture. Fermat Days, Toulouse.
- PENOT, J.-P. (1995) Conditioning convex and nonconvex problems. *J. Optim. Th. Appl.* **90** (3), 539–558.
- PENOT, J.-P. (1997) Duality for radiant and shady problems. *Acta Math. Vietnamica* **22** (2), 541–566.
- PENOT, J.-P. (1998) Are generalized derivatives useful for generalized convex functions? In: J.-P. Crouzeix, J.-E. Martinez-Legaz and M. Volle, eds., *Generalized Convexity, Generalized Monotonicity: Recent Results*. Kluwer, Dordrecht, 3–59.
- PENOT, J.-P. (1998) Well-behavior, well-posedness and nonsmooth analysis. *Pliska Stud. Math. Bulgar.* **12**, 141–190.
- PENOT, J.-P. (2000) What is quasiconvex analysis? *Optimization* **47**, 35–110.
- PENOT, J.-P. (2001) Duality for anticonvex programs. *J. of Global Optim.* **19**, 163–182.
- PENOT, J.-P. and VOLLE, M. (1987) Dualité de Fenchel et quasi-convexité. *C. R. Acad. Sci., Paris, Série I* **304** (13), 371–374.
- PENOT, J.-P. and VOLLE, M. (1988) Another duality scheme for quasiconvex problems. In: K.H. Hoffmann et al., eds., *Trends in Mathematical Optimization*. Int. Series Numer. Math. **84**, Birkhäuser, Basel, 259–275.
- PENOT, J.-P. and VOLLE, M. (1990) On quasi-convex duality. *Math. Oper. Research* **15**, 597–625.
- PENOT, J.-P. and VOLLE, M. (1990) On strongly convex and paraconvex dualities. In: A. Cambini, E. Castagnoli, L. Martein, P. Mazzoleni, S. Schaible, eds., *Generalized convexity and fractional programming with economic applications*. Lecture notes in Econ. and Math. Systems **345**, Springer Verlag, Berlin, 198–218.
- PICHARD, K. (2001) *Equations différentielles dans les espaces métriques. Applications à l'évolution de domaines*. Thesis, Univ. of Pau.
- PICHARD, K. and GAUTIER, S. (2000) Equations with delays in metric spaces: the mutational approach. *Numer. Funct. Anal. Optim.* **21**, 917–932.
- PINI, R. and SINGH, C. (1997) A survey of recent advances in generalized convexity with applications to duality theory and optimality conditions (1985-1995). *Optimization* **39** (4), 311–360.

- POLIQVIN, R. (1992) An extension of Attouch's Theorem and its application to second-order epi-differentiation of convexly composite functions. *Trans. Amer. Math. Soc.* **332**, 861–874.
- ROBINSON, S. (1991) Strongly regular generalized equations. *Math. Oper. Research* **16**, 292–309.
- ROCKAFELLAR, R.T. (1974) Augmented Lagrange multiplier functions and duality in nonconvex programming. *SIAM J. Control Optim.* **12**, 268–285.
- ROLEWICZ, S. (1993) Generalization of Asplund inequalities on Lipschitz functions. *Arch. Math.* **61**, 484–488.
- ROLEWICZ, S. (1994) On Mazur Theorem for Lipschitz functions. *Arch. Math.* **63**, 535–540.
- ROLEWICZ, S. (1994) Convex analysis without linearity. *Control and Cybern.* **23**, 247–256.
- ROLEWICZ, S. (1996) Duality and convex analysis in the absence of linear structure. *Math. Japonica* **44**, 165–182.
- RUBINOV, A.M. (2000) *Abstract Convexity and Global Optimization*. Kluwer, Dordrecht.
- RUBINOV, A.M. and ANDRAMONOV, M.YU. (1999) Minimizing increasing starshaped functions based on abstract convexity. *J. of Global Optim.* **15**, 19–39.
- RUBINOV, A.M. and GLOVER, B.M. (1997) On generalized quasiconvex conjugacy. *Contemporary Mathematics* **204**, 199–217.
- RUBINOV, A.M. and GLOVER, B.M. (1998) Duality for increasing positively homogeneous functions and normal sets. *Rech. Opér.* **32**, 105–123.
- RUBINOV, A.M. and GLOVER, B.M. (1998) Quasiconvexity via two step functions. In: J.-P. Crouzeix, J.-E. Martinez-Legaz and M. Volle, eds., *Generalized Convexity, Generalized Monotonicity: Recent Results*. Kluwer, Dordrecht, 159–183.
- RUBINOV, A.M., GLOVER, B.M. and YANG, X.Q. (1999) Decreasing functions with applications to optimization. *SIAM J. Optim.* **10**, 289–313.
- RUBINOV, A.M. and SHVEIDEL, A.P. (2000) Separability of star-shaped sets with respect to infinity. In: X. Yang et al., eds., *Progress in Optimization*. Kluwer, Dordrecht, 45–63.
- RUBINOV, A.M. and SIMSEK, B. (1995) Conjugate quasiconvex nonnegative functions. *Optimization* **35**, 1–22.
- RUBINOV, A.M. and SIMSEK, B. (1995) Dual problems of quasiconvex maximization. *Bull. Aust. Math. Soc.* **51**, 139–144.
- SCHWARTZ, L. (1973) *Radon measures on arbitrary topological spaces and cylindrical measures*. Lecture Notes, Tata Institute **6**, Bombay, Oxford University Press, London.
- SCHWARTZ, L. (1980) *Semi-martingales sur des variétés et martingales conformes sur des variétés réelles ou complexes*. Lecture Notes in Maths. **780**, Springer Verlag, Berlin.

- SINGER, I. (1986) Some relations between dualities, polarities, coupling functions and conjugations. *J. Math. Anal. Appl.* **115**, 1–22.
- SINGER, I. (1987) Optimization by level set methods. VI: generalizations of surrogate type reverse convex duality. *Optimization* **18** (4), 485–499.
- SINGER, I. (1997) *Abstract Convex Analysis*. J. Wiley, New York.
- TAO, P.D. and EL BERNOUSSI, S. (1988) Duality in D.C. (difference of convex functions). Optimization. Subgradient methods. In: K.H. Hoffmann et al. eds., *Trends in Mathematical Optimization*. Int. Series Numer. Math. **84**, Birkhauser, Basel, 277–293.
- TAO, P.D. and EL BERNOUSSI, S. (1989) Numerical methods for solving a class of global nonconvex optimization problems. In: J.-P. Penot, ed., *New methods in optimization and their industrial uses*. Int. Series Numer. Math. **97**, Birkhäuser, Basel, 97–132.
- THACH, P.T. (1991) Quasiconjugate of functions, duality relationships between quasiconvex minimization under a reverse convex constraint and quasiconvex maximization under a convex constraint and application. *J. Math. Anal. Appl.* **159**, 299–322.
- THACH, P.T. (1993) Global optimality criterion and a duality with a zero gap in nonconvex optimization. *SIAM J. Math. Anal.* **24** (6), 1537–1556.
- THACH, P.T. (1994) A nonconvex duality with zero gap and applications. *SIAM J. Optim.* **4** (1), 44–64.
- THACH, P.T. (1995) Diewert-Crouzeix conjugation for general quasiconvex duality and applications. *J. Optim. Th. Appl.* **86** (1), 719–743.
- TUY, H. (1995) D.C. optimization: theory, methods and algorithms. In: R. Horst and P.M. Pardalos, eds., *Handbook of Global Optimization*. Kluwer, Dordrecht, Netherlands, 149–216.
- VLADIMIROV, A.A., NESTEROV, YU. E. and CHEKANOV, YU.N. (1978) Uniformly convex functions. *Vestnik Moskov. Univ. Ser. 115, Vyschisl. Mat. Kibernet.* **3**, 12–23.
- VOLLE, M. (1985) Conjugaison par tranches. *Annali Mat. Pura Appl.* **139**, 279–312.
- VOLLE, M. (1997) Quasiconvex duality for the max of two functions. In: P. Grizzman, R. Horst, E. Sachs, R. Tichatschke, eds., *Recent Advances in Optimization*. Lecture Notes in Econ. and Math. Systems **452**, Springer Verlag, Berlin, 365–379.
- VOLLE, M. (1998) Duality for the level sum of quasiconvex functions and applications. *ESAIM: Control, Optimisation and the Calculus of Variations* art. 1, **3**, 329–343, URL: <http://www.emath.fr/cocv/>.
- WOLSEY, L.A. (1981) Integer programming duality: price functions and sensitivity analysis. *Math. Programming* **20**, 173–195.
- ZALINESCU, C. (1983) *On uniformly convex functions*. *J. Math. Anal. Appl.* **95**, 344–374.
- ZALINESCU, C. (2002) *Convex Analysis in General Vector Spaces*. World Scientific, Singapore.