

On weak sharp minima in vector optimization with
applications to parametric problems

by

Ewa Bednarczuk

Systems Research Institute of the PAS
ul. Newelska 6, 01-447 Warsaw, Poland

Abstract: In the paper we discuss the concepts of weak sharp solutions to vector optimization problems. As an application we provide sufficient conditions for stability of solutions in perturbed vector optimization problems.

Keywords: vector optimization, weak sharp solutions, stability.

1. Introduction

Let X and Y be normed spaces and let $\mathcal{K} \subset Y$ be a closed convex pointed cone in Y . We consider vector optimization problems of the form

$$(VOP) \quad \begin{array}{l} \mathcal{K} - \min f(x) \\ \text{subject to } x \in A, \end{array}$$

where $f : X \rightarrow Y$ and $A \subset X$ is a feasible set. By $E \subset Y$ we denote the set of all global efficient points to (VOP) , i.e., $\alpha \in E$ iff $(f(A) - \alpha) \cap (-\mathcal{K}) = \{0\}$ and by $S \subset X$ we denote the set of all its global solutions, $S = A \cap f^{-1}(E)$.

The role of weak sharp minima in scalar optimization in relation to stability of parametric problems and error bounds is widely recognized, see, e.g., Attouch and Wets (1993); Auslander and Crouzeix (1988); Azé and Corvellec (2002); Bonnans and Shapiro (2000); Burke and Deng (2002).

In vector optimization several definitions of global weak sharp solutions has been proposed, see, e.g., Bednarczuk (2004, 2007); Deng and Yang (2004), for the linear case.

The aim of this paper is to discuss several concepts of (global) weak sharp solutions to problem (VOP) and their applications to stability of parametric problems. In Section 2 weak sharp solutions to (VOP) are presented and their basic properties are elucidated. In Section 3 weak sharp solutions are exploited to formulate sufficient conditions for stability of parametric problems.

2. Global weak sharp solutions

By B_X and \bar{B}_X we denote open and closed unit balls in X , respectively. For any set $C \subset X$, $d(x, C) = \inf\{\|x - c\| : c \in C\}$. For any $\alpha \in Y$ put $S_\alpha := \{x \in A : f(x) = \alpha\}$.

DEFINITION 1 (see Bednarczuk, 2007) *Let $\alpha \in E$. We say that the solution set S to (VOP) is (globally) S_α -weak sharp if there exists a constant $\tau > 0$ such that*

$$f(x) - \alpha \notin \tau d(x, S_\alpha)B_Y - \mathcal{K} \quad \text{for all } x \in A \setminus S_\alpha. \quad (1)$$

Optimality conditions for S_α weak sharpness in the local setting have been recently investigated by Studniarski (2007). If $\text{int } \mathcal{K} \neq \emptyset$, a point $x_0 \in A$ is a *weak solution* to (VOP), $x_0 \in WS$, if $(f(A) - f(x_0)) \cap (-\text{int } \mathcal{K}) = \emptyset$. If there exists $\alpha \in E$ such that S is S_α -weak sharp, then $S = WS$.

Let $\alpha \in E$. We define a set-valued mapping $\mathcal{E}^\alpha : R_+ \rightrightarrows X$ as

$$\mathcal{E}^\alpha(\varepsilon) := A \cap f^{-1}(\alpha + \varepsilon B_Y - \mathcal{K}). \quad (2)$$

Clearly, $\mathcal{E}^\alpha(0) = S_\alpha$ and $\text{graph } \mathcal{E}^\alpha = \{(\varepsilon, x) \in R_+ \times A : f(x) \in \alpha + \varepsilon B_Y - \mathcal{K}\}$. There exist approaches to well-posedness of (VOP) via continuity properties of set-valued mappings similar to \mathcal{E}^α (see e.g. Bednarczuk, 2004, 2007; Miglierina and Molho, 2003, 2007; Zaffaroni, 2003).

PROPOSITION 1 *Let $\alpha \in E$ and let S be S_α -weak sharp with constant $\tau > 0$.*

(i) *If f is Lipschitz on A with constant L , then $\tau \leq L$.*

(ii) *The following condition holds:*

(C1) *there exists $\varepsilon_0 > 0$ such that for each $0 \leq \varepsilon \leq \varepsilon_0$*

$$A \cap f^{-1}(\alpha + \varepsilon B_Y - \mathcal{K}) \subset S_\alpha + \varepsilon \frac{1}{\tau} B_Y.$$

Proof. (i) If $f : X \rightarrow Y$ is Lipschitz on A with constant $L > 0$, i.e.

$$\|f(x) - f(x')\| \leq L\|x - x'\| \quad \text{for all } x, x' \in A,$$

then $\|f(x) - \alpha\| \leq L\|x - x'\|$ for any $x, x' \in A$, $f(x') = \alpha$, and, consequently, $\|f(x) - \alpha\| \leq Ld(x, S_\alpha)$ for all $x \in A$. On the other hand, $f(x) - \alpha \notin \tau d(x, S_\alpha)B_Y - \mathcal{K}$ for $x \in A \setminus S_\alpha$. In particular, $\|f(x) - \alpha\| \geq \tau d(x, S_\alpha)$ for $x \in A \setminus S_\alpha$, which gives the required inequality.

(ii) Suppose, on the contrary, that (C1) does not hold, i.e., there exist sequences $\varepsilon_n \rightarrow 0^+$ and $(x_n) \subset A$ such that

$$f(x_n) \in \alpha + \varepsilon_n B_Y - \mathcal{K} \quad \text{for } n \geq 1,$$

and $d(x_n, S_\alpha) > \varepsilon_n \frac{1}{\tau}$. Hence, for $n \geq 1$, $x_n \notin S_\alpha$, $\tau d(x_n, S_\alpha) > \varepsilon_n$ and

$$f(x_n) \in \alpha + \tau d(x_n, S_\alpha)B_Y - \mathcal{K},$$

which contradicts S_α -weak sharpness of S . ■

Condition (C1) of Proposition 1 (ii) can be rephrased by saying that the set-valued mapping \mathcal{E}^α defined by (2), is upper Lipschitz at $0 \in \text{dom } \mathcal{E}$ with constant $\frac{1}{\tau} > 0$, where a set-valued mapping $\Gamma : X \rightrightarrows Y$ is *upper Lipschitz* at $x_0 \in \text{dom } \Gamma$ with constant $L > 0$ if there exists $t > 0$ such that $\Gamma(x) \subset \Gamma(x_0) + L\|x - x_0\|B_Y$ for $x \in B(x_0, t)$.

Recall that $\alpha \in E$ is a (global) *strict efficient point to (VOP)* (Bednarczuk, 2004) if there exists a constant $\gamma > 0$ such that

$$f(x) - \alpha \notin \gamma \|f(x) - \alpha\| B_Y - \mathcal{K} \quad \text{for } x \in A \quad f(x) \neq \alpha. \tag{3}$$

As before, if f is Lipschitz on A with constant L we have $\|f(x) - \alpha\| \leq L\|x - x'\|$ for all $x \in A$ and $x' \in S_\alpha$ and consequently $\|f(x) - \alpha\| \leq Ld(x, S_\alpha)$ for all $x \in A$.

If moreover, S is S_α -weak sharp with constant $\tau > 0$ we get

$$f(x) - \alpha \notin \frac{\tau}{L} \|f(x) - \alpha\| B_Y - \mathcal{K} \quad \text{for } x \in A \setminus S_\alpha, \tag{4}$$

which means that $\alpha \in E$ is strict efficient with constant $\frac{\tau}{L}$.

In this way we proved the following proposition.

PROPOSITION 2 *Let f be Lipschitz on A with constant $L > 0$. If S is S_α -weak sharp with constant $\tau > 0$, then $\alpha \in E$ is strict efficient with constant $\frac{\tau}{L}$.*

DEFINITION 2 (see Bednarczuk, 2007) *Let $\alpha \in E$. We say that the solution set S to (VOP) is α -weak sharp if there exists a constant $\tau > 0$ such that*

$$f(x) - \alpha \notin \tau d(x, S) B_Y - \mathcal{K} \quad \text{for all } x \in A \setminus S. \tag{5}$$

If, for some $\alpha \in E$, the solution set S is S_α -weak sharp, then S is α -weak sharp.

PROPOSITION 3 *Let $\alpha \in E$. If S is α -weak sharp with constant $\tau > 0$, the following condition holds:*

(C2) *there exists $\varepsilon_0 > 0$ such that for each $0 \leq \varepsilon \leq \varepsilon_0$*

$$A \cap f^{-1}(\alpha + \varepsilon B_Y - \mathcal{K}) \subset S + \varepsilon \frac{1}{\tau} B_Y.$$

Proof. Suppose, on the contrary, that (C2) does not hold, i.e., there exist sequences $\varepsilon_n \rightarrow 0^+$ and $(x_n) \subset A$ such that $f(x_n) \in \alpha + \varepsilon_n B_Y - \mathcal{K}$ and $d(x_n, S) > \varepsilon_n \frac{1}{\tau}$ for $n \geq 1$. Hence, $x_n \notin S$, $\tau d(x_n, S) > \varepsilon_n$ and $f(x_n) \in \alpha + \tau d(x_n, S) B_Y - \mathcal{K}$, which contradicts α -weak sharpness of S . ■

Consider now linear multicriteria problems of the form

$$(LMP) \quad \begin{aligned} & R_+^m - \min Cx \\ & \text{subject to } x \in A, \end{aligned}$$

where R_+^m is a nonnegative orthant, $C : R^n \rightarrow R^m$ is a linear mapping and $A \subset R^n$ is polyhedral set. According to Deng and Yang (2004), WS is a set of *weak sharp solutions to (LMP)* if there exists a constant $\tau > 0$ such that

$$d(Cx, WE) \geq \tau d(x, WS) \quad \text{for } x \in A, \quad (6)$$

where $WE = f(WS)$. Basing ourselves on this idea we define weak sharp solutions to (VOP).

DEFINITION 3 *We say that the solution set S to (VOP) is (globally) weak sharp if there exists a constant $\tau > 0$ such that*

$$d(f(x), E) \geq \tau d(x, S) \quad \text{for all } x \in A. \quad (7)$$

PROPOSITION 4 *Let $\tau > 0$ be given. If for any $\alpha \in E$ the set S is α -weak sharp with constant τ , then the solution set S is weak sharp with constant τ .*

Proof. By assumption, for any $\alpha \in E$,

$$f(x) - \alpha \notin \tau d(x, S)B_Y - \mathcal{K} \quad \text{for } x \in A \setminus S.$$

In particular, $f(x) - \alpha \notin \tau d(x, S)B_Y$ for $x \in A \setminus S$ and any $\alpha \in E$, which gives the assertion. \blacksquare

3. Lipschitz continuities of efficient points

Consider now parametric vector optimization problems of the form

$$(VOP)_u \quad \begin{array}{l} \mathcal{K} - \min f(x) \\ \text{subject to } x \in A(u), \end{array}$$

where the parameter u belongs to a normed space U . By $E(u)$ and $S(u)$ we denote the set of efficient points and the solution set to $(VOP)_u$, respectively.

In this section we exploit weak sharpness and S_α -weak sharpness to provide sufficient conditions for Lipschitzness of $E(u)$ and $S(u)$ near a given $u_0 \in U$. For other types of convergence of efficient points see e.g. Miglierina and Molho (2007).

In what follows the restrictions on behaviour of sets $A(u)$ around a given u_0 are expressed through continuity properties of the mapping $F : U \rightrightarrows X$, $F(u) = A(u)$, $F(u_0) = A$. Recall that a set-valued mapping $\Gamma : U \rightrightarrows X$ is *lower Lipschitz* at $u_0 \in \text{dom } \Gamma$ if there exist constants $L > 0$ and $t > 0$ such that $\Gamma(u_0) \subset \Gamma(u) + L\|u - u_0\|B_Y$ for $u \in B(u_0, t)$. Γ is *Lipschitz* at $u_0 \in \text{dom } \Gamma$ if Γ is upper and lower Lipschitz at u_0 . Moreover, Γ is *Lipschitz* around $u_0 \in \text{dom } \Gamma$ if there exist constants $L > 0$ and $t > 0$ such that $\Gamma(u) \subset \Gamma(u') + L\|u - u'\|B_Y$ for $u, u' \in B(u_0, t)$. The *domination property (DP)* holds for (VOP) if for any $x \in A$ there exists $\bar{x} \in S$ such that $f(\bar{x}) \in f(x) - \mathcal{K}$. Let us note that if $f : X \rightarrow R$, (DP) is satisfied provided $S \neq \emptyset$.

THEOREM 1 Let $f : X \rightarrow Y$ be Lipschitz on X with constant $L_f > 0$. If

- (i) $F : U \rightrightarrows X$ is Lipschitz at $u_0 \in \text{dom } F$ with constants $L_c > 0, t > 0$,
- (ii) (DP) holds for all $(VOP)_u$ with $u \in B(u_0, t)$,
- (iii) there exists $\tau > 0$ such that for each $\alpha \in E$ the solution set S is S_α -weak sharp with constant τ , i.e. for each $\alpha \in E$,

$$f(x) - \alpha \notin \tau d(x, S_\alpha)B_Y - \mathcal{K} \quad \text{for } x \in A \setminus S_\alpha,$$

then

$$E \subset E(u) + (L_c L_f + \frac{2L_c L_f^2}{\tau}) \|u - u_0\| B_Y \quad \text{for } u \in B(u_0, t).$$

If moreover, for $u \in B(u_0, t) \setminus \{u_0\}$ the sets $S(u)$ are weak sharp with constant τ , then

$$S \subset S(u) + (L_c + \frac{2L_c L_f}{\tau} + \frac{2L_c L_f^2}{\tau^2}) \|u - u_0\| B_X \quad \text{for } u \in B(u_0, t).$$

Proof. By (i), (ii), $u_0 \in \text{int dom } E$. Take any $\alpha \in E$ and $u \in B(u_0, t)$. There exists $\bar{x} \in S$ such that $f(\bar{x}) = \alpha$. By (i), there exists $z \in A(u)$ such that $\|\bar{x} - z\| \leq L_c \|u - u_0\|$. If $d(f(z), E(u)) \leq 2L_c L_f \|u - u_0\|$, the conclusion follows. Otherwise, by (ii), there exists $\bar{z} \in S(u)$ such that $f(\bar{z}) \in f(z) - \mathcal{K}$ and $\|f(z) - f(\bar{z})\| > 2L_c L_f \|u - u_0\|$. By (i), there exists $x \in A$ such that $\|x - \bar{z}\| \leq L_c \|u - u_0\|$ and by the Lipschitzness of f

$$\|f(x) - f(\bar{x})\| \geq \|f(z) - f(\bar{z})\| - \|f(z) - f(\bar{x})\| - \|f(\bar{z}) - f(x)\| > 0,$$

and

$$\begin{aligned} f(x) - f(\bar{x}) &= (f(x) - f(\bar{z})) + (f(\bar{z}) - f(z)) + (f(z) - f(\bar{x})) \\ &\in 2L_f L_c \|u - u_0\| B_Y - \mathcal{K}. \end{aligned}$$

By (iii) and by Proposition 2, $f(x) - f(\bar{x}) \notin \frac{\tau}{L_f} \|f(x) - f(\bar{x})\| B_Y - \mathcal{K}$. This proves that $\tau \|f(x) - f(\bar{x})\| \leq 2L_c L_f^2 \|u - u_0\|$ and consequently

$$\begin{aligned} \|f(\bar{x}) - f(\bar{z})\| &\leq \|f(\bar{x}) - f(x)\| + \|f(x) - f(\bar{z})\| \\ &\leq (L_f L_c + \frac{2L_f^2 L_c}{\tau}) \|u - u_0\| \end{aligned}$$

which proves the first assertion.

To prove the second assertion take any $x_0 \in S$ and $u \in B(u_0, t) \setminus \{u_0\}$. By the first assertion, there exists $z_0 \in S(u)$, $f(z_0) = \eta$, such that

$$f(x_0) - \eta \in (L_c L_f + \frac{2L_c L_f^2}{\tau}) \|u - u_0\| B_Y.$$

By (i), there exists $z \in A(u)$ such that $\|x_0 - z\| \leq L_c \|u - u_0\|$. If $d(z, S(u)) \leq L_c \|u - u_0\|$, the conclusion follows. Otherwise, since $S(u)$, $u \neq u_0$, is weak sharp, $f(z) - \eta \notin \tau d(z, S(u))B_Y$. Moreover,

$$f(z) - \eta = (f(z) - f(x_0)) + (f(x_0) - \eta) \in (2L_c L_f + \frac{2L_c L_f^2}{\tau}) \|u - u_0\| B_Y.$$

Hence, $\tau d(z, S(u)) \leq (2L_c L_f + \frac{2L_c L_f^2}{\tau}) \|u - u_0\|$ and

$$d(x_0, S(u)) \leq \|x_0 - z\| + d(z, S(u)) \leq (L_c + \frac{2L_c L_f}{\tau} + \frac{2L_c L_f^2}{\tau^2}) \|u - u_0\|.$$

Let $u \in U$ and $\eta \in Y$. Put $S_\eta(u) = \{x \in A(u) : f(x) = \eta\}$. ■

THEOREM 2 Let $f : X \rightarrow Y$ be Lipschitz on X with constant $L_f > 0$. If
 (i) $F : U \rightrightarrows X$ is Lipschitz at $u_0 \in \text{dom } F$ with constants $L_c > 0$ and $t > 0$,
 (ii) (DP) holds for (VOP),
 (iii) there exists $\tau > 0$ such that for $u \in B(u_0, t)$, $u \neq u_0$, and $\eta \in E(u)$ the sets $S(u)$ are $S_\eta(u)$ -weak sharp with constant τ , i.e.

$$f(x) - \eta \notin \tau d(x, S_\eta(u))B_Y - \mathcal{K} \text{ for } x \in A(u) \setminus S_\eta(u),$$

then $E(u) \subset E + (L_f L_c + \frac{2L_c L_f^2}{\tau}) \|u - u_0\| B_Y$ for $u \in B(u_0, t)$.

If, moreover, S is weak sharp, then

$$S(u) \subset S + (L_c + \frac{2L_c L_f}{\tau} + \frac{2L_c L_f^2}{\tau}) \|u - u_0\| B_X \text{ for } u \in B(u_0, t).$$

Proof. Note that by (ii), $E \neq \emptyset$. Take any $u \in B(u_0, t)$. If $E(u) = \emptyset$, the conclusion follows. Otherwise, take any $\eta \in E(u)$. There exists $z_0 \in S(u)$, $f(z_0) = \eta$. By (i), there exists $x \in A$ such that $\|z_0 - x\| \leq L_c \|u - u_0\|$. If $d(f(x), E) \leq 2L_c L_f \|u - u_0\|$, the conclusion follows.

Otherwise, by (ii), there is $x_0 \in S$, $f(x_0) = \alpha$, such that $f(x_0) \in f(x) - \mathcal{K}$ and $\|f(x) - \alpha\| > 2L_c L_f \|u - u_0\|$. By (i), there exists $z \in A(u)$ such that $\|z - x_0\| \leq L_c \|u - u_0\|$. By the Lipschitzness of f , $f(z) - \eta = f(z) - f(x_0) + f(x_0) - f(x) + f(x) - \eta \in 2L_c L_f \|u - u_0\| B_Y - \mathcal{K}$. Since

$$\|f(z) - \eta\| \geq \|f(x) - \alpha\| - \|f(x) - \eta\| - \|f(z) - \alpha\| > 0,$$

by (iii) and by Proposition 2, $f(z) - \eta \notin \frac{\tau}{L_f} \|f(z) - \eta\| B_Y - \mathcal{K}$. Consequently, $\|f(z) - \eta\| \leq \frac{2L_c L_f^2}{\tau} \|u - u_0\|$ and

$$\|\eta - \alpha\| \leq \|f(z) - \eta\| + \|f(z) - \alpha\| \leq (L_c L_f + \frac{2L_c L_f}{\tau}) \|u - u_0\|.$$

To prove the second assertion, take any $z_0 \in S(u)$, $u \in B(u_0, t)$. By the first assertion of the theorem, there exists $x_0 \in S$, $f(x_0) = \alpha$, such that $f(z_0) - f(x_0) \in (L_c L_f + \frac{2L_c L_f}{\tau}) \|u - u_0\|$. By (i), there exists $x \in A$ such that $\|x - z_0\| \leq L_c \|u - u_0\|$. If $d(x, S) \leq 2L_c \|u - u_0\|$, the conclusion follows. Otherwise, by (ii), there exists $x_0 \in S$, $f(x_0) = \alpha$, such that $f(x_0) \in f(x) - \mathcal{K}$ and $\|x - x_0\| > 2L_c \|u - u_0\|$. Hence,

$$f(x) - f(x_0) = f(x) - f(z_0) + f(z_0) - f(x_0) \in (2L_c L_f + \frac{2L_c L_f^2}{\tau}) \|u - u_0\|.$$

Since S is weak sharp, $f(x) - f(x_0) \notin \tau d(x, S) B_Y$, which proves that $d(z_0, S) \leq (L_c + \frac{2L_c L_f}{\tau} + \frac{2L_c L_f^2}{\tau^2}) \|u - u_0\|$. ■

The above theorems immediately lead to the following result.

THEOREM 3 *Let $f : X \rightarrow Y$ be Lipschitz on X with constant $L_f > 0$. Assume that*

- (i) *the set valued mapping $F : U \rightrightarrows X$ is Lipschitz around $u_0 \in \text{dom } F$ with constants $L_c > 0$ and $t > 0$,*
- (ii) *(DP) holds for $(VOP)_u$, $u \in B(u_0, t)$,*
- (iii) *there exists $\tau > 0$ such that for $u \in B(u_0, t)$ and $\eta \in E(u)$ the solution sets $S(u)$ to $(VOP)_u$ are $S_\eta(u)$ -weak sharp with constant τ .*

Then

$$S(u) \subset S(u') + (L_c + \frac{2L_c L_f}{\tau} + \frac{2L_c L_f^2}{\tau^2}) \|u' - u\| B_X \text{ for } u, u' \in B(u_0, t/2).$$

References

- ATTOUCH, H. and WETS, R.J.-B. (1993) Quantitative stability of variational systems II. A framework for nonlinear conditioning. *SIAM J. Optim.* **3**, 359-381.
- AUSLANDER, A. and CROUZEIX, J.-P. (1988) Global regularity conditions. *Math. of O.R.* **13**, 243-253.
- AZÉ, D. and J.-N. CORVELLEC (2002) On the sensitivity analysis of Hoffmann constants for systems of linear inequalities. *SIAM J. Optim.* **12**, 913-927.
- BEDNARCZUK, E.M. (2004) Weak sharp efficiency and growth condition for vector-valued functions with applications. *Optimization* **53**, (5-6), 455-474
- BEDNARCZUK, E.M. (2007) Stability analysis for parametric vector optimization problems. *Dissertationes Mathematicae*, **442**, Warsaw.
- BONNANS, F. and SHAPIRO, A. (2000) *Perturbation Analysis of Optimization Problems*. Springer Verlag, New York.
- BURKE, J.V. and DENG, S. (2002) Weak Sharp Minima Revisited. Part I: Basic Theory. *Control and Cybernetics* **31**, 439-471.

- DENG, S. and YANG, X.Q. (2004) Weak sharp minima in multicriteria linear programming. *SIAM J. Optim.* **15**, 456-460.
- JIMÉNEZ, B. and NOVO, V. (2003) First and second order sufficient conditions for strict minimality in nonsmooth vector optimization. *JMAA* **284**, 496-510.
- MIGLIERINA, E. and MOLHO, E. (2003) Well-posedness and convexity in vector optimization. *Math. Meth. Op. Res.* **58**, 375-385.
- MIGLIERINA, E. and MOLHO, E. (2005) Convergence of minimal sets in convex vector optimization. *SIAM J. Opt.* **15**, 513-526.
- STUDNIARSKI, M. (2007) Weak sharp minima in multiobjective optimization. *preprint*
- ZAFFARONI, A. (2003) Degrees of efficiency and degrees of minimality. *SIAM J. Control and Optim.* **42**, 1071-1086.