## COLLOQUIUM MATHEMATICUM

VOL. LXIX 1995 FASC. 1

## ERRATUM TO "ON CONVOLUTION OPERATORS WITH SMALL SUPPORT WHICH ARE FAR FROM BEING CONVOLUTION BY A BOUNDED MEASURE"

(COLLOQ. MATH. 67 (1994), 33-60)

BY

EDMOND E. GRANIRER (VANCOUVER, BRITISH COLUMBIA)

On p. 35, row 31, delete "ultrathin".

On p. 42, row 14, insert after "...p. 23).": "By [LT], p. 103, A(F) contains a complemented copy of  $\ell^1$ ."

On p. 55, row 6, insert between "a" and "function algebra": "(not necessarily semisimple)".

On p. 56, row 14, replace " $v_{n_j}w_0=0$  on F" by " $v_{n_j}w_0\in J$ . Since if  $w_0=0$  on the open set  $U_0$  with  $e\in U_0$  let i be such that  $F\cap\operatorname{cl} V_{n_j}\subset F\cap U_0$  if  $j\geq i$ . Now  $D=\{x\in G:w_0(x)\neq 0\text{ and }v_{n_j}(x)\neq 0\}\subset (G\sim U_0)\cap\operatorname{cl} V_{n_j},$  a closed set in G. Thus  $F\cap\operatorname{supp}(w_0v_{n_j})\subset F\cap\operatorname{cl} D\subset (G\sim U_0)\cap\operatorname{cl} V_{n_j}=\emptyset$  if  $j\geq i$ . Since  $\operatorname{supp} w_0v_{n_j}$  is compact and disjoint from  $F,w_0v_{n_j}\in J_F\subset J$  if  $j\geq i$  (here cl denotes closure)."

DEPARTMENT OF MATHEMATICS UNIVERSITY OF BRITISH COLUMBIA VANCOUVER, BRITISH COLUMBIA CANADA V6T 1Z2

Reçu par la Rédaction le 12.12.1994