

Table des matières du tome LXXV, fascicule 3

	Pages
G. L. Itzkowitz, On the density character of compact topological groups	201-203
D. M. Hyman, A remark on Fox's paper on shape	205-208
H. H. Wicke and J. M. Worrell, Jr., Topological completeness of first countable Hausdorff spaces I	209-222
A. Mostowski, Models of second order arithmetic with definable Skolem functions	223-234
J. F. Wells, The restricted cancellation law in a Noether lattice	235-247
T. W. Ma, Non-singular set-valued compact fields in locally convex spaces	249-259
R. H. Reese, The local contractibility of the homeomorphism space of a 2-polyhedron	261-273
W.W. Comfort and S. Negrepontis, On families of large oscillation	275-290
W.J. Gilbert, Menger's Theorem for topological spaces	291-295

Les FUNDAMENTA MATHEMATICAE publient, en langues des congrès internationaux, des travaux consacrés à la *Théorie des Ensembles, Topologie, Fondements de Mathématiques, Fonctions Réelles, Algèbre Abstraite*.
Chaque volume paraît en 3 fascicules

Adresse de la Rédaction et de l'Échange:
FUNDAMENTA MATHEMATICAE, Śniadeckich 8, Warszawa 1 (Pologne)

Le prix de ce fascicule est 4.35 \$

Tous les volumes sont à obtenir par l'intermédiaire de
ARS POLONA - RUCH, Krakowskie Przedmieście 7, Warszawa 1 (Pologne)

On the density character of compact topological groups

by

Gerald L. Itzkowitz (Buffalo, N. Y.)

1. Introduction. In 1943, S. Kakutani [8] proved that, for a compact Abelian group G , if $w(G) = \mathfrak{w}$ is the least cardinal of a basis of open sets for G then the density character (see Hewitt [2]) of G is \mathfrak{n} , where \mathfrak{n} is the least cardinal number satisfying $2^{\mathfrak{n}} \geq \mathfrak{w}$. In that paper Kakutani also showed that the cardinal of G is $|G| = 2^{\mathfrak{w}}$. Thus one could state this theorem in the following form: Let \mathfrak{n} be the least cardinal number such that $\text{card } G \leq 2^{\mathfrak{n}}$; then there is a dense subgroup in G of cardinal \mathfrak{n} .

In 1958 S. Hartman and A. Hulanicki [1] gave a simpler proof of this theorem, based on a theorem of Hewitt [2] on the density character of a certain product of topological spaces. In a further paper Hulanicki [4] extended Hewitt's theorem to prove that a certain product of measurable spaces $P_{t \in T} X_t$, $w(X_t) \leq \aleph_0$, $\text{card } T \leq 2^{\mathfrak{n}}$, had a dense set of full outer measure of cardinal \mathfrak{n}^{\aleph_0} . With this result he was able to prove that if \mathfrak{n} is the least cardinal number such that $\text{card } G \leq 2^{2^{\mathfrak{n}}}$, G being a compact Abelian group, then G contains a dense subgroup of full outer measure and cardinal \mathfrak{n}^{\aleph_0} .

G. L. Itzkowitz [6], using again a theorem similar to Hewitt's, extended this result to prove that such a G contains a dense pseudo-compact subgroup of cardinal \mathfrak{n}^{\aleph_0} (hence of full outer measure in G). H. Wilcox [12], using a result of N. Th. Varopoulos, proved that this result may be extended to include all compact groups. Left unanswered was Kakutani's question in [8]: Is \mathfrak{n} the density character of G when G is a compact group and \mathfrak{n} is the least cardinal such that $\text{card } G \leq 2^{2^{\mathfrak{n}}}$? Wilcox's theorem shows that the density character of G is at most \mathfrak{n}^{\aleph_0} .

It is the purpose of this note to answer Kakutani's question in the affirmative. Our method of proof is essentially the same as the method followed by H. Wilcox. For completeness we will give a new proof, also based on Hewitt's theorem, of Kakutani's theorem for compact Abelian groups.

2. The density character in groups. Let $w(G)$ denote the least cardinal of a basis of open sets of G ($w(G)$ is also called the density character of G). If \mathcal{B} is a collection of sets, we shall say that the collection of sets C is a *weak base* for \mathcal{B} if, given $B \in \mathcal{B}$, there is a $C \in \mathcal{C}$ such that $C \subset B$.

We recall the theorem of Hewitt, [2].

2.1. THEOREM (Hewitt). Let $X = \prod_{t \in T} X_t$, where $\{(X_t, \tau_t) : t \in T\}$ is a family of topological spaces such that $w(X_t) \leq \aleph$, for each $t \in T$, and card $T \leq 2^\aleph$. Then the product space (X, τ) has a weak base \mathcal{B} for the open sets for which card $\mathcal{B} \leq \aleph$. (Note that $w(X) = 2^\aleph$.)

2.2. COROLLARY. Let $G = \prod_{t \in T} H_t$, where each H_t is a compact separable topological group, and card $T = 2^\aleph$. Then $w(G) = 2^\aleph$, and there is a weak base for the open sets of G having cardinal \aleph .

2.3. THEOREM (Vilenkin [12]). Let G be a compact Abelian group. For some cardinal number w , there is a continuous mapping of $\{-1, 1\}^w$ onto G ; w can be taken to be $\max\{s_0, r\}$, where r is the rank of the character group of G .

Note. If card $G > s_0$ then $r = \text{card } G$.

2.4. THEOREM (Kakutani). Let G be an infinite compact Abelian group satisfying $w(G) = w$, and let \aleph be the least cardinal such that $w \leq 2^\aleph$. Then G has a dense subgroup J of cardinal \aleph .

Proof. Pontryagin [10] has shown that if G is a locally compact Abelian group then $w(G) = w(\hat{G})$, where \hat{G} is the character group of G . Thus 2.3 implies that \hat{G} is the continuous image of $G' = \{-1, 1\}^w$. Evidently G' has a weak base for its open sets of cardinal \aleph , and so \hat{G} has a weak base \mathcal{B} for its open sets of cardinal \aleph . Let J be the subgroup of G generated by the set obtained by selecting one point from each element of the weak base \mathcal{B} .

Hulanicki [5], Ivanovskii [7], and Kuzminov [9] have shown that a 0-dimensional infinite topological group G that is compact is homeomorphic with the space $\{-1, 1\}^w$, where $\{-1, 1\}$ is discrete and $w = w(G)$. Thus an immediate corollary of Hewitt's Theorem 2.1 that we have cited is:

2.5. THEOREM. Let G be an infinite compact 0-dimensional topological group satisfying $w(G) = w$, and let \aleph be the least cardinal such that $w \leq 2^\aleph$. Then G has a dense subgroup J of cardinal \aleph .

Varopoulos has shown in [11] that if G is a connected compact group and Z is its centre, then G/Z has the form $\prod_{\alpha \in A} M_\alpha$ with all the M_α compact metrizable groups. Thus we may state the following theorem:

2.6. THEOREM. Let G be an infinite compact connected topological group satisfying $w(G) = w$, and let \aleph be the least cardinal such that $w \leq 2^\aleph$. Then G has a dense subgroup J of cardinal \aleph .

Proof. Let Z be the centre of G : then it is evident that $w(Z) \leq w$ and that 2.4 applies to Z . It is also clear that $G/Z = \prod_{\alpha \in A} M_\alpha$ is of the proper form for an application of 2.1. Thus we observe that both Z and G/Z have dense subsets of cardinal at most \aleph . It is now an elementary exercise (see Hewitt and Ross, 5.38(f)) to show that G has a dense subgroup of cardinal \aleph .

2.7. THEOREM. Let G be an infinite compact topological group satisfying $w(G) = w$, and let \aleph be the least cardinal such that $w \leq 2^\aleph$. Then G has a dense subgroup J of cardinal \aleph .

Proof. Let C be the component of the identity in G . Then G/C is 0-dimensional. Obviously $w(G/C) \leq w$, and $w(C) < w$, so that Theorems 2.5 and 2.6 apply to G/C and C respectively. Thus again we may conclude that G contains a dense subgroup of cardinal \aleph .

Remark. The theorem in Hewitt and Ross, 5.38(f), is concerned with a dense set of the desired size; however, it is clear that the group generated by this set also has cardinal \aleph .

References

- [1] S. Hartman and A. Hulanicki, *Sur les ensembles denses de puissance minimum dans les groupes topologiques*, Coll. Math. 6 (1958), pp. 187-191.
- [2] E. Hewitt, *A remark on density characters*, Bull. Amer. Math. Soc. 52 (1946), pp. 641-643.
- [3] — and K. A. Ross, *Abstract Harmonic Analysis I*, Berlin 1963.
- [4] A. Hulanicki, *On subsets of full outermeasure in products of measurable spaces*, Bull. Acad. Polon. Sci. 6 (1959), pp. 331-335.
- [5] — *On the topological structure of 0-dimensional topological groups*, Fund. Math. 46 (1959), pp. 317-320.
- [6] G. L. Itzkowitz, *Extensions of Haar measure for compact connected Abelian groups*, Proc. Koninkl. Nederl. Akad. van Wet.-en, Amsterdam, Ser. A, 68, No. 2 and Indag. Math. 27, No. 2 (1965), pp. 190-207.
- [7] L. N. Ivanovskii, *On an hypothesis of P. S. Aleksandrov*, Dokl. Akad. Nauk 123 (1958), pp. 785-786.
- [8] S. Kakutani, *On cardinal numbers related with a compact Abelian group*, Proc. Acad. Tokyo 19, pp. 366-372.
- [9] V. Kuzminov, *On an hypothesis of P. S. Aleksandrov in the theory of topological groups*, Dokl. Akad. Nauk 125 (1959), pp. 727-729.
- [10] L. S. Pontryagin, *Continuous groups*, 2nd Edition, Moscow 1954.
- [11] N. Th. Varopoulos, *A theorem on the continuity of homomorphisms of locally compact groups*, Proc. Camb. Phil. Soc. 60, (1964), pp. 449-463.
- [12] N. Ya. Vilenkin, *On the dyadicity of the group space of bicommutative groups*, Uspehi Mat. Nauk 13, 6 (84), (1958), pp. 79-80.
- [13] H. J. Wilcox, *Pseudocompact groups*, Pac. J. of Math. 19, No 2 (1966), pp. 365-379.

Reçu par la Rédaction le 14. 4. 1970