

Références

- [1] C. A. Akemann and G. K. Pedersen, *Ideal perturbations of elements in C^* -algebras*, Math Scand. 41 (1977), 117–139.
- [2] C. Apostol, *The reduced minimum modulus*, Michigan Math. J. 32 (1985), 279–294.
- [3] K. Astala and H.-O. Tylli, *On the bounded compact approximation property and measures of noncompactness*, J. Funct. Anal. 70 (1987), 388–401.
- [4] S. Goldberg, *Unbounded Linear Operators*, McGraw-Hill, New York, 1966.
- [5] P. R. Halmos, *A Hilbert Space Problem Book*, D. Van Nostrand, 1967.
- [6] P. de la Harpe, *Initiation à l'algèbre de Calkin*, in: Lecture Notes in Math. 725, Springer, 1978, 180–219.
- [7] R. E. Harte and M. Mbekhta, *On generalized inverses in C^* -algebras*, Studia Math. 103 (1992), 71–77.
- [8] —, —, *Generalized inverses in C^* -algebras II*, ibid. 106 (1993), 129–138.
- [9] T. Kato, *Perturbation Theory for Linear Operators*, Springer, Berlin, 1966.
- [10] J. Ph. Labrousse et M. Mbekhta, *Les opérateurs points de continuité pour la conorme et l'inverse de Moore-Penrose*, Houston J. Math. 18 (1992), 7–23.
- [11] L. E. Labuschagne, A. Ströh and J. Swart, *The uniqueness of operational quantities in von Neumann algebras*, preprint.
- [12] C. L. Olsen and J. K. Plastiras, *Quasialgebraic operators, compact perturbations, and the essential norm*, Michigan Math. J. 21 (1974), 385–397.
- [13] G. K. Pedersen, *Spectral formulas in quotient C^* -algebras*, Math. Z. 148 (1976), 299–300.
- [14] J. Zemánek, *Geometric characteristics of semi-Fredholm operators and their asymptotic behaviour*, Studia Math. 80 (1984), 219–234.
- [15] —, *The semi-Fredholm radius of a linear operator*, Bull. Polish Acad. Sci. Math. 32 (1984), 67–76.

Adresse actuelle de Mostafa Mbekhta:

UNIVERSITÉ DES SCIENCES ET
TECHNOLOGIES DE LILLE
U.R.A. D 751 CNRS "GAT"
U.F.R. DE MATHÉMATIQUES
F-59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

UNIVERSITÉ GALATASARAY
CIRAGAN CAD. NO. 102
ORTAKOY 80840
ISTANBUL, TURQUIE

Received August 10, 1995
Revised version August 18, 1995

(3450)

On the exponential Orlicz norms of stopped Brownian motion

by

GORAN PEŠKIR (Aarhus and Zagreb)

Abstract. Necessary and sufficient conditions are found for the exponential Orlicz norm (generated by $\psi_p(x) = \exp(|x|^p) - 1$ with $0 < p \leq 2$) of $\max_{0 \leq t \leq \tau} |B_t|$ or $|B_\tau|$ to be finite, where $B = (B_t)_{t \geq 0}$ is a standard Brownian motion and τ is a stopping time for B . The conditions are in terms of the moments of the stopping time τ . For instance, we find that $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$ as soon as

$$E(\tau^k) = O(C^k k^k)$$

for some constant $C > 0$ as $k \rightarrow \infty$ (or equivalently $\|\tau\|_{\psi_1} < \infty$). In particular, if $\tau \sim \text{Exp}(\lambda)$ or $|N(0, \sigma^2)|$ then the last condition is satisfied, and we obtain

$$\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq K\sqrt{E(\tau)}$$

with some universal constant $K > 0$. Moreover, this inequality remains valid for any class of stopping times τ for B satisfying $E(\tau^k) \leq C(E\tau)^k k^k$ for all $k \geq 1$ with some fixed constant $C > 0$. The method of proof relies upon Taylor expansion, Burkholder-Gundy's inequality, best constants in Doob's maximal inequality, Davis' best constants in the L^p -inequalities for stopped Brownian motion, and estimates of the smallest and largest positive zero of Hermite polynomials. The results extend to the case of any continuous local martingale (by applying the time change method of Dubins and Schwarz).

1. Introduction. The main aim of the paper is to investigate and establish necessary and sufficient conditions for the exponential integrability of the supremum of a reflecting Brownian motion taken over a random time interval (as well as of stopped Brownian motion itself).

More precisely, let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, let τ be a stopping time for B , and let $\|\cdot\|_\psi$ denote the Orlicz norm generated

1991 *Mathematics Subject Classification*: Primary 60G40, 60J65; Secondary 46E30, 60G44, 60H05.

Key words and phrases: Brownian motion (Wiener process), stopping time, exponential Young function, exponential Orlicz norm, Doob's maximal inequality for martingales, Burkholder-Gundy's inequality, Davis' best constants, Hermite polynomial, continuous (local) martingale, Ito's integral, the quadratic variation process, time change (of Brownian motion), Kahane-Khinchin's inequalities.

by the Young function ψ . Thus, if X is a random variable, then

$$(1.1) \quad \|X\|_\psi = \inf\{c > 0 \mid E(\psi(|X|/c)) \leq 1\}$$

with $\inf(\emptyset) = \infty$. In this paper we are interested in Young functions of exponential growth, and therefore choose to work with $\psi_p(x) = \exp(|x|^p) - 1$ for $0 < p \leq 2$. (To handle the small convexity problem around zero when $0 < p < 1$, we can let $\psi_p(x) = \exp(|x|^p) - 1$ for $x \geq (1/p - 1)^{1/p}$ and take $\psi_p(x)$ to be linear on $[0, (1/p - 1)^{1/p}]$.) The main problem under consideration is to find out when the quantities $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_p}$ (or $\|B_\tau\|_{\psi_p}$) are finite, and to obtain sharp estimates of these for $0 < p \leq 2$. In view of the Burkholder–Davis–Gundy (and related) inequalities, where the integrability is usually established for Young functions of a moderate growth, we think that this problem appears worthy of consideration and its solution by itself might be of theoretical and practical interest.

At the very beginning of thinking about this problem, it was not clear to us in what terms the conditions (we should look for) are to be expressed. For this reason we find it convenient here to explain our minding in this direction in more detail. In this context we were firstly motivated by some fundamental (closely related) results in the discrete parameter case. If $\varepsilon = \{\varepsilon_k\}_{k \geq 1}$ is a Rademacher sequence of random variables and $\{a_k\}_{k \geq 1}$ is a sequence of real numbers, then the *Khinchin inequality* states (see [7]) that

$$(1.2) \quad \left\| \sum_{k=1}^n a_k \varepsilon_k \right\|_p \leq A_p \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}$$

for all $0 < p < \infty$ and all $n \geq 1$, where the constant A_p depends only on p . The best value for A_p in (1.2) is known (see [6]):

$$(1.3) \quad A_p = \begin{cases} \sqrt{2}(\Gamma((p+1)/2)/\sqrt{\pi})^{1/p} & \text{if } 2 \leq p < \infty, \\ 1 & \text{if } 0 < p \leq 2. \end{cases}$$

In particular, taking $a_1 = \dots = a_n = 1$ in (1.2), we obtain

$$(1.4) \quad \left\| \sum_{k=1}^n \varepsilon_k \right\|_p \leq A_p \sqrt{n}$$

for all $0 < p < \infty$ and all $n \geq 1$.

Having (1.2) with (1.3) for $2 \leq p < \infty$, and using Taylor expansion of $x \mapsto \exp(|x|^2)$, the passage from the power Orlicz norms (generated by $\varphi_p(x) = |x|^p$) in (1.2) to the exponential Orlicz norm generated by ψ_2 is rather smooth (see [11]–[13]):

$$(1.5) \quad \left\| \sum_{k=1}^n a_k \varepsilon_k \right\|_{\psi_2} \leq \sqrt{\frac{8}{3}} \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}$$

for all $a_1, \dots, a_n \in \mathbb{R}$ with $n \geq 1$. Moreover, in this way we obtain a very precise information: the constant $\sqrt{8/3}$ in (1.5) is the best possible. By using exactly the same procedure one could deduce the inequality (see [7])

$$(1.6) \quad \left\| \sum_{k=1}^n a_k \varepsilon_k \right\|_{\psi_p} \leq B_p \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}$$

for all $0 < p \leq 2$ and all $a_1, \dots, a_n \in \mathbb{R}$ with $n \geq 1$, where the constant B_p depends only on p . In particular, taking $a_1 = \dots = a_n = 1$ in (1.6), we obtain

$$(1.7) \quad \left\| \sum_{k=1}^n \varepsilon_k \right\|_{\psi_p} \leq B_p \sqrt{n}$$

for all $0 < p \leq 2$ and all $n \geq 1$.

In view of our main problem stated above, two questions arise naturally in this setting. The first one is: What happens with the right-hand side in (1.4) when n on the left-hand side is replaced by a stopping time τ for ε ? (It would be ideal that \sqrt{n} on the right-hand side may be replaced by $\sqrt{E(\tau)}$ or $E(\sqrt{\tau})$, but this is not true in general as we shall see below.) The second one is: Provided that we have an affirmative answer to the first question, is it possible to imitate the passage from (1.2) to (1.5) + (1.6) via Taylor expansion and obtain an analogue of (1.7), with n on the right-hand side being replaced by $\sqrt{E(\tau)}$ or $E(\sqrt{\tau})$? (In view of the negative answer to the first question, this also fails in general.)

The answer to the first question is well known (see [3] and [5]):

$$(1.8) \quad \left\| \sum_{k=1}^{\tau} \varepsilon_k \right\|_p \leq C_p E(\tau^{p/2})$$

for any stopping time τ for ε and all $0 < p < \infty$, where the constant C_p depends only on p . It should also be well known that this is the best we can expect in the context of the first question. Thus, in view of the second question, this indicates that in order to pass from (1.8) to an analogue of (1.7) via Taylor expansion, one has to take care about all moments of the stopping time τ . This observation significantly clarifies what our approach should be towards solution of our main problem stated above. In other words, it becomes clear that our necessary and sufficient conditions should be expressed in terms of all moments of the stopping time τ , or rather in terms of the asymptotic behaviour of the quantity $E(\tau^k)$ as $k \rightarrow \infty$. Still, it is not clear what an analogue of (1.7) could be after the extension of (1.8) via Taylor expansion as explained above. In particular, how large is (and how can be described) the class of stopping times for which the analogue of (1.7) remains valid? We find these questions of interest and leave them unanswered. *In this paper we focus only on the continuous parameter case,*

and in this context provide answers to analogous questions. In this process the preceding discussion will serve mainly as a motivation.

The preceding conclusion about the necessity of taking into account all moments of the stopping time (in order to gain exponential integrability) becomes particularly transparent in the continuous parameter case, after recalling *Burkholder-Gundy's inequality* (see [2]):

$$(1.9) \quad G_p E(\tau^{p/2}) \leq E(\max_{0 \leq t \leq \tau} |B_t|^p) \leq H_p E(\tau^{p/2}),$$

valid for all stopping times τ for B and all $0 < p < \infty$ with some universal constants G_p and H_p depending only on p . Thus in order to deduce the exponential integrability of $\max_{0 \leq t \leq \tau} |B_t|$ via Taylor expansion one should have a precise information on the constants H_p for $0 < p < \infty$. However, the best values for H_p 's in (1.9) do not seem to be known by now. Nonetheless, we will see that we can approximate these numbers in an accurate way by using Doob's maximal inequality (with best constants) and Davis' best constants for an analogue of (1.9) with $\max_{0 \leq t \leq \tau} |B_t|$ replaced by $|B_\tau|$. This leads to the necessity of estimating the (largest and smallest positive) zero of Hermite polynomials and is the approach which is taken in this paper. The details will be presented later. Here we only indicate results obtained in this way.

The main result of the paper (see Theorems 3.4, 3.12 and 3.14) states that

$$(1.10) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_p} < \infty$$

for any stopping time τ for B and $0 < p \leq 2$, as soon as

$$(1.11) \quad E(\tau^k) = O(C^k k^{k(2-p)/p})$$

with some constant $C > 0$ as $k \rightarrow \infty$. Moreover (see Theorem 3.12),

$$(1.12) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_p} \leq \sqrt{10} \frac{p}{p-1} e^{1/p} \sqrt{D} \left(1 + \frac{(\Delta_p(D))^{p/2}}{\sqrt{2\pi}} \right)$$

for all $1 < p \leq 2$ and any $D > 0$, where we set

$$(1.13) \quad \Delta_p(D) = \sup_{k \geq 1} \left(\frac{E(\tau^k)}{D^k k^{k(2-p)/p}} \right).$$

In the case $p = 1$, we have the following estimate (see Theorem 3.4):

$$(1.14) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq 6e\sqrt{D} \left(1 + \frac{\sqrt{\Delta_1(D)}}{\sqrt{2\pi}} \right)$$

for any $D > 0$. In particular, if $\tau \sim \text{Exp}(\lambda)$ or $|N(0, \sigma^2)|$ with some $\lambda, \sigma^2 > 0$, then (1.11) is satisfied, and moreover in this case (1.14) yields (see Exam-

ples 3.8 and 3.9)

$$(1.15) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq K\sqrt{E(\tau)}$$

with some universal constant $K > 0$. We remark that the given stopping time $\tau \sim \text{Exp}(\lambda)$ or $|N(0, \sigma^2)|$ is not necessarily independent of the Brownian motion B , and in view of (1.7) we think that the inequality (1.15) is by itself of theoretical and practical interest. Moreover, this inequality remains valid for any class of stopping times τ for B satisfying $E(\tau^k) \leq C(E\tau)^k k^k$ for all $k \geq 1$ with some fixed constant $C > 0$ (see Corollary 3.7). Finally, if (1.10) is satisfied for some $0 < p \leq 2$, then (see Theorem 3.2)

$$(1.16) \quad E(\tau^k) = O(C^k k^{k(2+p)/p})$$

for some constant $C > 0$ as $k \rightarrow \infty$. We note that there is a gap between the sufficient condition (1.11) and necessary condition (1.16), but this was the optimum we could obtain by using our method here (see Problem 3.10).

To conclude the introduction let us mention that the results just indicated extend from the Brownian motion case to the case of any continuous local martingale (Ito's integral). This is achieved by applying the standard time change method of Dubins and Schwarz. The results in this context are presented in more detail in Section 4.

2. Power integrability of a stopped Brownian motion. In this section we introduce the notation and collect without proof several facts which will be used in the proofs of our main results in the next section.

We work with a fixed probability space (Ω, \mathcal{F}, P) which is large enough to support all random functions under consideration. Moreover, whenever a filtration $(\mathcal{F}_t)_{t \geq 0}$ of (Ω, \mathcal{F}, P) is considered, it is assumed to satisfy the usual conditions: \mathcal{F} is P -complete, \mathcal{F}_0 contains all P -null sets in \mathcal{F} , and $\mathcal{F}_{t+} = \mathcal{F}_t$ for all $t \geq 0$. The main object under investigation in this paper is Brownian motion. We recall that a (standard) *Brownian motion* is a process $B = (B_t)_{t \geq 0}$ defined on (Ω, \mathcal{F}, P) with $B_0 = 0$ P -a.s. for which there exists a filtration $(\mathcal{F}_t)_{t \geq 0}$ such that $B_{t+h} - B_t$ is independent of \mathcal{F}_t , and has the Gaussian distribution with expectation 0 and variance h for $t, h \geq 0$. A random variable $\tau : \Omega \rightarrow [0, \infty]$ is said to be a *stopping time* for B if $\{\tau \leq t\} \in \mathcal{F}_t$ for all $t \geq 0$. If this condition holds for a filtration $(\mathcal{F}_t)_{t \geq 0}$ of (Ω, \mathcal{F}, P) which is not necessarily linked with a Brownian motion, we will say that τ is an (\mathcal{F}_t) -*stopping time*.

Let τ be a stopping time for a Brownian motion B such that $\{B_{t \wedge \tau} | t \geq 0\}$ is uniformly integrable (which holds if $E(\sqrt{\tau}) < \infty$ for instance). Then *Doob's maximal inequality* states that

$$(2.1) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_p \leq \frac{p}{p-1} \|B_\tau\|_p$$

for all $1 < p < \infty$ with the constant $p/(p - 1)$ being the best possible (see [16]).

Let τ be a stopping time for a Brownian motion B . Then *Burkholder-Gundy's inequality* (1.9) is valid for all $0 < p < \infty$ with some universal constants G_p and H_p depending only on p . The best values for G_p and H_p are not known. However, it is known (see [8]) that

$$(2.2) \quad \frac{2-p}{4-p} E(\tau^{p/2}) \leq E(\max_{0 \leq t \leq \tau} |B_t|^p) \leq \frac{4-p}{2-p} E(\tau^{p/2})$$

whenever $0 < p < 2$.

Let τ be a stopping time for a Brownian motion B . Then *Burkholder-Davis' inequality* states that

$$(2.3) \quad E(|B_\tau|^p) \leq A_p E(\tau^{p/2}) \quad \text{if } 0 < p < \infty,$$

$$(2.4) \quad a_p E(\tau^{p/2}) \leq E(|B_\tau|^p) \quad \text{if } 1 < p < \infty \text{ and } E(\tau^{p/2}) < \infty,$$

with the best values for A_p and a_p in (2.3)+(2.4) given by (see [3])

$$(2.5) \quad A_p = \begin{cases} (z_p^*)^p & \text{if } 2 \leq p < \infty, \\ (z_p)^p & \text{if } 0 < p \leq 2, \end{cases}$$

$$(2.6) \quad a_p = \begin{cases} (z_p)^p & \text{if } 2 \leq p < \infty, \\ (z_p^*)^p & \text{if } 1 < p \leq 2, \end{cases}$$

where z_p^* denotes the largest positive zero of the *parabolic cylinder function* $D_p(x)$, while z_p denotes the smallest positive zero of the *confluent hypergeometric function* $x \mapsto M(-p/2, 1/2, x^2/2)$.

We recall that $M(a, b, z)$ denotes the *Kummer function*; it is a solution of the differential equation (see [1], p. 189):

$$(2.7) \quad zy''(z) + (b - z)y'(z) - ay(z) = 0.$$

The Kummer function is explicitly given by the expression (see [1], p. 189)

$$(2.8) \quad M(a, b, z) = 1 + \sum_{k=0}^{\infty} \frac{a(a+1)\dots(a+k)}{b(b+1)\dots(b+k)} \cdot \frac{z^k}{k!}.$$

Moreover, we have (see [1], p. 194)

$$(2.9) \quad M(-n, 1/2, x^2/2) = (-1)^n 2^n \frac{n!}{(2n)!} \text{He}_{2n}(x),$$

where $\text{He}_n(x)$ denotes the *Hermite polynomial* of degree n :

$$(2.10) \quad \text{He}_n(x) = n! \sum_{k=0}^{[n/2]} (-1)^k \frac{1}{k! 2^k (n-2k)!} x^{n-2k}.$$

We recall that the *parabolic cylinder function* $D_p(x)$ is defined by (see [1], pp. 281-282)

$$(2.11) \quad D_p(x) = Y_1(x) \cos(p\pi/2) + Y_2(x) \sin(p\pi/2),$$

where $Y_1(x)$ and $Y_2(x)$ are given by

$$(2.12) \quad Y_1(x) = (2^{p/2}/\sqrt{\pi})\Gamma((p+1)/2)y_1(x),$$

$$(2.13) \quad Y_2(x) = (2^{(p+1)/2}/\sqrt{\pi})\Gamma((p+2)/2)y_2(x),$$

where $y_1(x)$ and $y_2(x)$ are linearly independent solutions of the differential equation

$$(2.14) \quad y''(x) + (ax^2 + bx + c)y(x) = 0$$

with $a = -1/4$, $b = 0$ and $c = p + 1/2$. Actually, we have

$$(2.15) \quad y_1(x) = \exp(-x^2/4)M(-p/2, 1/2, x^2/2),$$

$$(2.16) \quad y_2(x) = x \exp(-x^2/4)M(-p/2 + 1/2, 3/2, x^2/2).$$

In view of (2.9), (2.11), (2.12) and (2.15) we see that z_{2n}^* is the largest positive zero (z_{2n} is the smallest positive zero) of the Hermite polynomial $\text{He}_{2n}(x)$. Our next aim is to state estimates of these numbers. This will be used heavily in the proof of our main result in the next section.

For this, let $L_n^{(\alpha)}(x)$ denote the *generalized Laguerre polynomial*:

$$(2.17) \quad L_n^{(\alpha)}(x) = \sum_{k=0}^n (-1)^k \binom{n+\alpha}{n-k} \frac{1}{k!} x^k$$

and let $x_m^{(n,\alpha)}$ denote the m th (positive) zero of $L_n^{(\alpha)}(x)$. Let $J_\nu(z)$ denote the *Bessel function*:

$$(2.18) \quad J_\nu(z) = \frac{(z/2)^\nu e^{-iz}}{\Gamma(\nu+1)} M(\nu+1/2, 2\nu+1, 2iz),$$

and let $j_m^{(\alpha)}$ denote the m th positive zero of $J_\alpha(x)$. Then we have (see [1], p. 346)

$$(2.19) \quad \frac{(j_m^{(\alpha)})^2}{4k_m} < x_m^{(n,\alpha)} < \frac{k_m}{k_n} (2k_m + \sqrt{4k_m^2 + 1/4 - \alpha^2}),$$

where $k_r = r + (\alpha + 1)/2$.

In order to apply this result in our context, one should note that

$$(2.20) \quad \text{He}_{2n}(x) = (-1)^n 2^n n! L_n^{(-1/2)}(x^2/2),$$

$$(2.21) \quad J_{-1/2}(x) = \sqrt{\frac{2}{\pi x}} \cos(x).$$

Thus $j_1^{(-1/2)} = \pi/2$, and from (2.19)–(2.21) we obtain

$$(2.22) \quad \frac{\pi}{\sqrt{2(4n+1)}} < z_{2n},$$

$$(2.23) \quad z_{2n}^* < \sqrt{2(4n+1)},$$

for all $n \geq 1$.

A useful application of (2.22) and (2.23) is based upon the following two facts:

$$(2.24) \quad p \mapsto z_p \text{ is (strictly) decreasing on }]0, \infty[,$$

$$(2.25) \quad p \mapsto z_p^* \text{ is (strictly) increasing on }]1, \infty[.$$

This can be easily seen by recalling some results on the square root stopping boundaries. Namely, let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let us consider the stopping times defined by

$$(2.26) \quad \tau_a = \inf\{t > 0 : |B_t| = a\sqrt{t+1}\},$$

$$(2.27) \quad \sigma_a = \inf\{t > 0 : B_t = a\sqrt{t-1}\},$$

where $a > 0$. Then we have (see [15] and [10] respectively)

$$(2.28) \quad E(\tau_a)^p < \infty \text{ if and only if } a < z_{2p} \text{ whenever } 0 < p < \infty,$$

$$(2.29) \quad E(\sigma_a)^p < \infty \text{ if and only if } a > z_{2p}^* \text{ whenever } 1/2 < p < \infty.$$

Hence (2.24) and (2.25) are easily verified.

3. Exponential integrability of a stopped Brownian motion.

In this section we present the main results of the paper. Throughout $B = (B_t)_{t \geq 0}$ denotes a standard Brownian motion defined on the probability space (Ω, \mathcal{F}, P) , and τ denotes a stopping time for B with $E(\tau^k) < \infty$ for all $k \geq 1$. Our main aim is to find necessary and sufficient conditions for the exponential Orlicz norms

$$(3.1) \quad \left\| \max_{0 \leq t \leq \tau} |B_t| \right\|_{\psi_p}$$

to be finite, and obtain estimates of these for $0 < p \leq 2$ in terms of the moments of the stopping time τ . By Taylor expansion and (2.1) it is easily seen that the quantity in (3.1) is finite if and only if $\|B_\tau\|_{\psi_p}$ is finite. Here we shall concentrate on the quantity (3.1), but the proofs and estimates obtained are easily adapted to cover the case of $\|B_\tau\|_{\psi_p}$ as well. We begin by finding necessary conditions. In this context the following lemma is shown to be useful.

LEMMA 3.1. For all $p, k \geq 1$ there exist universal constants $\varepsilon_{p,k} > 0$ such that

$$(3.2) \quad \varepsilon_{p,k} \|X\|_k \leq \|X\|_{\psi_p}$$

for every random variable X . In fact, one can take

$$(3.3) \quad \varepsilon_{p,k} = (pe/k)^{1/p}.$$

Proof. Fix $p, k \geq 1$. It is clear that the problem reduces to finding the largest constant $c > 0$ for which

$$(3.4) \quad E \left(\exp \left(\frac{|X|^p}{c^p \|X\|_k^p} \right) \right) > 2$$

whenever X is a random variable. By Taylor expansion, (3.4) can be rewritten in the form

$$(3.5) \quad \sum_{n=1}^{\infty} \frac{E|X|^{np}}{c^{np} \|X\|_k^{np} n!} = \sum_{n=1}^{\infty} \frac{1}{n!} \left(\frac{\|X\|_{np}}{c \|X\|_k} \right)^{np} > 1.$$

Let m denote the smallest $n \geq 1$ satisfying $np \geq k$. Then by Jensen's inequality we will have (3.5) as soon as

$$\frac{1}{m!} \left(\frac{1}{c} \right)^{mp} \geq 1.$$

Hence by Stirling's formula we can conclude that

$$c \leq \left(\frac{1}{m!} \right)^{1/(mp)} = \left(\frac{1}{\sqrt{2\pi m} m^m e^{-m} e^{r_m}} \right)^{1/(mp)} \leq \left(\frac{e}{m} \right)^{1/p} \leq \left(\frac{pe}{k} \right)^{1/p},$$

where $1/(12m+1) < r_m < 1/(12m)$. This completes the proof of the lemma. ■

We may now state a necessary condition for the exponential integrability of the supremum of a reflecting Brownian motion taken over a random time interval (as well as of a stopped Brownian motion itself).

THEOREM 3.2. Let τ be a stopping time for a Brownian motion $B = (B_t)_{t \geq 0}$ satisfying $E(\tau^k) < \infty$ for all $k \geq 1$ such that

$$(3.6) \quad \left\| \max_{0 \leq t \leq \tau} |B_t| \right\|_{\psi_p} < \infty$$

for some $0 < p < \infty$. Then

$$(3.7) \quad E(\tau^k) = O(C^k k^{k(2+p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$.

Proof. Put $A = \left\| \max_{0 \leq t \leq \tau} |B_t| \right\|_{\psi_p}$, where $p \geq 1$ is such that (3.6) is satisfied. Then by (3.2)+(3.3) we have

$$(3.8) \quad A \geq (pe/k)^{1/p} \left\| \max_{0 \leq t \leq \tau} |B_t| \right\|_k$$

for all $k \geq 1$. From (2.2) with $p = 1$ we get

$$(3.9) \quad E \left(\max_{0 \leq t \leq \tau} |B_t| \right) \geq \frac{1}{3} E(\sqrt{\tau}).$$

By (2.4)+(2.6), (2.22) and (2.24) we easily find

$$(3.10) \quad E(\max_{0 \leq t \leq \tau} |B_t|^k) \geq \left(\frac{\pi}{11\sqrt{k}} \right)^k E(\tau^{k/2})$$

for all $k \geq 2$. Since $\pi/11 < 1/3$, by (3.9) we see that (3.10) remains valid for $k = 1$ as well. By (3.8) and (3.10) we have

$$(3.11) \quad \begin{aligned} 1 &\geq \left(\frac{(pe)^{1/p}}{A} \cdot \frac{\pi}{11} \right)^k \frac{1}{k^{k/p+k/2}} E(\tau^{k/2}) \\ &= \left(\frac{(pe)^{1/p}}{A} \cdot \frac{\pi}{11} \cdot \frac{1}{2^{(1/2)(1+2/p)}} \right)^k \frac{E(\tau^{k/2})}{(k/2)^{(k/2)(1+2/p)}} \\ &= \left(\frac{(pe/2)^{1/p}}{A} \cdot \frac{\pi}{11\sqrt{2}} \right)^k \frac{E(\tau^{k/2})}{(k/2)^{(k/2)(1+2/p)}} \end{aligned}$$

for all $k \geq 1$. Hence we conclude that

$$(3.12) \quad \sup_{k \geq 1} \left(\frac{(pe/2)^{2/p} \pi^2}{242A^2} \right)^k \frac{E(\tau^k)}{k^{k(2+p)/p}} \leq 1.$$

The case $0 < p < 1$ is treated similarly. This completes the proof of the theorem. ■

EXAMPLE 3.3. Let $\tau \sim \log N(\mu, \sigma^2)$ be a stopping time for a Brownian motion $B = (B_t)_{t \geq 0}$ with the log-normal density function

$$f(x) = \frac{1}{x\sigma\sqrt{2\pi}} \exp\left(-\frac{(\mu - \log x)^2}{2\sigma^2}\right) 1_{]0, \infty[}(x),$$

where $\mu \in \mathbb{R}$ and $\sigma^2 > 0$. Thus, if $X \sim N(\mu, \sigma^2)$ is a Gaussian random variable with mean μ and variance σ^2 , then $\tau \sim \exp(X)$. Then we have

$$E(\tau^k) = e^{\mu k + (\sigma^2/2)k^2}$$

for $k \geq 1$. Hence by (3.7) we easily find that

$$\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_p} = \infty$$

for all $0 < p < \infty$. In other words, there is no number $\varepsilon > 0$ such that

$$E(\exp(\varepsilon(\max_{0 \leq t \leq \tau} |B_t|^p))) < \infty$$

whenever $0 < p < \infty$. ■

Next we pass to sufficient conditions. First we investigate the case $p = 1$. The main result in this context is as follows.

THEOREM 3.4. Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let τ be a stopping time for B satisfying

$$(3.13) \quad E(\tau^k) = O(C^k k^k)$$

for some $C > 0$ as $k \rightarrow \infty$. Then

$$(3.14) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty.$$

Moreover, define

$$(3.15) \quad \Delta_1(D) = \sup_{k \geq 1} \left(\frac{E(\tau^k)}{D^k k^k} \right)$$

for $D > 0$. Then

$$(3.16) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq 6e\sqrt{D} \left(1 + \frac{\sqrt{\Delta_1(D)}}{\sqrt{2\pi}} \right)$$

for all $D > 0$.

Proof. Given a stopping time τ for B satisfying (3.13), define $B_\tau^* = \max_{0 \leq t \leq \tau} |B_t|$. Then by (2.1), (2.3)+(2.5), (2.23) and (2.25) we have

$$(3.17) \quad \begin{aligned} E(B_\tau^*)^k &\leq \left(\frac{k}{k-1} \right)^k E|B_\tau|^k \leq \left(\frac{k}{k-1} \right)^k (z_k^*)^k E(\tau^{k/2}) \\ &\leq \left(\frac{k}{k-1} \right)^k (z_{2k}^*)^k E(\tau^{k/2}) \\ &\leq \left(\frac{k}{k-1} \right)^k (\sqrt{2(4k+1)})^k E(\tau^{k/2}) \end{aligned}$$

for all $k \geq 2$. Moreover, by (2.2) with $p = 1$ we get

$$(3.18) \quad E(B_\tau^*) \leq 3E(\sqrt{\tau}).$$

Since $k/(k-1) \leq 2$ and $2(4k+1) \leq 9k$ for $k \geq 2$, from (3.17) and (3.18) by Taylor expansion we easily obtain

$$(3.19) \quad \begin{aligned} E\left(\exp\left(\frac{B_\tau^*}{c}\right)\right) - 1 &= \sum_{k=1}^{\infty} \frac{E(B_\tau^*)^k}{c^k k!} \\ &\leq \frac{3E(\sqrt{\tau})}{c} + \sum_{k=2}^{\infty} \left(\frac{6}{c}\right)^k \frac{k^{k/2}}{k!} E(\tau^{k/2}) \\ &\leq \sum_{k=1}^{\infty} \left(\frac{6}{c}\right)^k \frac{k^{k/2}}{k!} E(\tau^{k/2}) \end{aligned}$$

for all $c > 0$. Further, by Stirling's formula we find

$$(3.20) \quad \frac{k^{k/2}}{k!} = \frac{k^{k/2}}{\sqrt{2\pi k} k^k e^{-k} e^{r_k}} \leq \frac{e^k}{\sqrt{2\pi} k^{(k+1)/2}},$$

where $1/(12k+1) < r_k < 1/(12k)$ for $k \geq 1$. Inserting (3.20) into (3.19),

and using Jensen's inequality, we obtain the estimate

$$\begin{aligned}
 (3.21) \quad E\left(\exp\left(\frac{B_\tau^*}{c}\right)\right) - 1 &\leq \frac{1}{\sqrt{2\pi}} \sum_{k=1}^{\infty} \left(\frac{6e}{c}\right)^k \frac{E(\tau^{k/2})}{k^{(k+1)/2}} \\
 &= \frac{1}{\sqrt{2\pi}} \sum_{k=1}^{\infty} \left(\frac{6e\sqrt{D}}{c}\right)^k \frac{E(\tau^{k/2})}{D^{k/2} k^{(k+1)/2}} \\
 &\leq \frac{\sqrt{\Delta_1(D)}}{\sqrt{2\pi}} \sum_{k=1}^{\infty} \left(\frac{6e\sqrt{D}}{c}\right)^k
 \end{aligned}$$

for all $c, D > 0$. Identifying the last term in (3.21) with 1, we easily find out that

$$c = 6e\sqrt{D} \left(1 + \frac{\sqrt{\Delta_1(D)}}{\sqrt{2\pi}}\right).$$

This proves (3.16), and hence (3.14) follows as well. The proof is complete. ■

Remark 3.5. It should be noted that condition (3.13) may be equivalently formulated as follows:

$$(3.13') \quad \|\tau\|_{\psi_1} < \infty.$$

This is easily verified by applying Stirling's formula. Thus, in short, the result of Theorem 3.4 may be summarized: If τ is a stopping time for a Brownian motion $B = (B_t)_{t \geq 0}$ such that $\|\tau\|_{\psi_1} < \infty$, then $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$ as well.

Remark 3.6. Under the hypotheses of Theorem 3.4, let us refine the definition of $\Delta_1(D)$ by setting

$$\Delta_1^*(D) = \sup_{k \geq 1} \left(\frac{E(\tau^{k/2})}{D^k k^{(k+1)/2}}\right)$$

for $D > 0$. Then the preceding proof (without application of Jensen's inequality in (3.21)) shows that the following refinement of the estimate (3.16) is valid:

$$(3.16') \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq 6eD \left(1 + \frac{\Delta_1^*(D)}{\sqrt{2\pi}}\right)$$

for all $D > 0$. Although not as fine as (3.16'), the estimate (3.16) seems to be more applicable, since $\Delta_1(D)$ is generally computed more easily. We shall illustrate this in the next corollary and two examples, which serve as important applications of Theorem 3.4.

COROLLARY 3.7. Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let τ be a stopping time for B satisfying

$$(3.22) \quad E(\tau^k) \leq C(E\tau)^k k^k$$

for all $k \geq 1$ with some constant $C > 0$. Then

$$(3.23) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq 6e \left(1 + \sqrt{\frac{C}{2\pi}}\right) \sqrt{E(\tau)}.$$

Proof. In the notation of Theorem 3.4 take $D = E(\tau)$. Then by (3.22) we get $\Delta_1(D) \leq C$. Hence (3.23) follows straightforwardly from (3.16). This completes the proof. ■

EXAMPLE 3.8. Let the stopping time $\tau \sim \text{Exp}(\lambda)$ for a Brownian motion $B = (B_t)_{t \geq 0}$ be exponentially distributed with parameter $\lambda > 0$. Then $E(\tau^k) = k!/\lambda^k$ for $k \geq 1$. Thus (3.22) is satisfied with $C = 1$, and by (3.23) we see that τ satisfies the inequality

$$(3.24) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq 6e \left(1 + \frac{1}{\sqrt{2\pi}}\right) \sqrt{E(\tau)},$$

which is valid simultaneously for all $\lambda > 0$. It should be noted here that τ is not supposed to be independent of B (as is occasionally assumed in the literature). ■

Further applications of Theorem 3.4 are possible. We will not pursue this in more detail here, but instead consider another typical example.

EXAMPLE 3.9. Let the stopping time $\tau \sim |N(0, \sigma^2)|$ for Brownian motion $B = (B_t)_{t \geq 0}$ be from reflecting Gaussian distribution with mean 0 and variance σ^2 . Thus, if $X \sim N(0, \sigma^2)$, then $\tau \sim |X|$. It is well known that

$$(3.25) \quad E(\tau^k) = \frac{2^{k/2} \sigma^k}{\sqrt{\pi}} \Gamma((k+1)/2) \leq \frac{(\sqrt{2}\sigma)^k}{\sqrt{\pi}} k^k$$

for all $k \geq 1$. In the notation of Theorem 3.4 take $D = \sqrt{2}\sigma = \sqrt{\pi} E(\tau)$. Then by (3.25) we get $\Delta_1(D) \leq 1/\sqrt{\pi}$. Thus by (3.16) we see that τ satisfies the inequality

$$(3.26) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} \leq 6e \left(\sqrt{\sqrt{\pi}} + \frac{1}{\sqrt{2\pi}}\right) \sqrt{E(\tau)}$$

which is valid simultaneously for all $\sigma^2 > 0$. Note that the constant in (3.26) is only "slightly" increased in comparison with the constant appearing in (3.24). ■

PROBLEM 3.10. It should be noted that our method above, although optimally realized in accordance with our main idea which relies upon Taylor expansion and best constants in the L^p -inequalities, leaves a gap between necessary and sufficient conditions obtained. To explain this in more detail, note that Theorem 3.2 states that a necessary condition for $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$ is $E(\tau^k) = O(C^k k^{3k})$, while by Theorem 3.4 a sufficient condition is $E(\tau^k) = O(C^k k^k)$. We want to make it clear here that no attempt will be

made in the paper to single out a common necessary and sufficient condition, due partially to the fact that our sufficient condition is relatively satisfactory (recall Examples 3.8 and 3.9) and partially to the fact that such an attempt would require a new method (which is far from being evident). Our main aim was to extract as much as possible from the method which relies upon Taylor expansion and best constants in the L^p -inequalities, and it is clear from our proof that this goal is achieved. Anyway, from the general point of view, the following problem appears worthy of consideration. Consider the family of stopping times τ for a Brownian motion $B = (B_t)_{t \geq 0}$ satisfying $E(\tau^k) < \infty$ for all $k \geq 1$, and find the number $\alpha > 0$ such that the condition $E(\tau^k) = O(C^k k^{\alpha k})$ is equivalent to $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$. Note that our results above show that such a number α belongs to the interval $[1, 3]$.

In the context of the preceding problem, the following example is of interest.

EXAMPLE 3.11. Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and consider the following stopping time for B :

$$\tau = \inf\{t \geq 1 : |B_t| = at^\beta\},$$

where $a > 0$ and $0 < \beta < 1/2$ are given. Set $\tau_n = \tau \wedge n$ for $n \geq 1$. Note that by Jensen's inequality we have

$$(3.27) \quad E|B_{\tau_n}|^{2k} = a^{2k} E(\tau_n)^{2\beta k} \leq a^{2k} (E(\tau_n)^k)^{2\beta}$$

for all $n, k \geq 1$. On the other hand, by (2.4)+(2.6) and (2.22) we have

$$(3.28) \quad \left(\frac{\pi}{\sqrt{2(4k+1)}}\right)^{2k} E(\tau_n)^k \leq E|B_{\tau_n}|^{2k}$$

for all $n, k \geq 1$. From (3.27) and (3.28) by letting $n \rightarrow \infty$, we obtain

$$(E(\tau^k))^{1-2\beta} \leq \left(\frac{10}{\pi^2} a^2\right)^k k^k$$

for all $k \geq 1$. This shows that

$$(3.29) \quad E(\tau^k) = O(C^k k^{k/(1-2\beta)})$$

for some $C > 0$ as $k \rightarrow \infty$. In view of (2.1) it is clear that $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$ if and only if $\|B_\tau\|_{\psi_1} < \infty$, which is by Stirling's formula easily verified to be equivalent to $E(\tau^k) = O(C^k k^{k/\beta})$ for some $C > 0$ as $k \rightarrow \infty$. Thus, putting $1/(1-2\beta) \leq 1/\beta$, or equivalently $\beta \leq 1/3$, by (3.29) we see that $E(\tau^k) = O(C^k k^{k/\beta})$, showing that

$$(3.30) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$$

for all $0 < \beta \leq 1/3$. (We do not know what is going on with (3.30) for $\beta \in]1/3, 1/2[.$)

It is not known to us whether the estimate (3.29) is optimal, which would show (take $\beta = 1/3$) that there exists a stopping time τ for B with $E(\tau^k) \geq C^k k^{3k}$ such that $\|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_1} < \infty$ (recall Theorem 3.2 with $p = 1$). This sort of argument would pass through in the case of any optimal improvement upon (3.29) (by taking another β as above). Note, however, that the estimate (3.29) is asymptotically efficient in the following sense. The case $\beta \downarrow 0$ corresponds to the hitting time of the point $a > 0$ given by $\tau = \inf\{t \geq 1 : |B_t| = a\}$. By the i.i.d. increments we have $P\{\tau > n + 1\} = P\{\max_{1 \leq t \leq n+1} |B_t| < a\} \leq P\{|B_2 - B_1| < 2a, |B_3 - B_2| < 2a, \dots, |B_{n+1} - B_n| < 2a\} = (P\{|B_1| < 2a\})^n$ for all $n \geq 1$. Thus $P\{\tau > n\} \leq \varepsilon^n$ for all $n \geq 1$ with some $\varepsilon > 0$, which easily implies that $E(\exp(\tau/c)) < \infty$ for some $c > 0$. This is equivalent to $\|\tau\|_{\psi_1} < \infty$, or in other words, $E(\tau^k) = O(C^k k^k)$, which agrees with (3.29) as $\beta \downarrow 0$. The case $\beta \uparrow 1/2$ corresponds to the case of the hitting time of the square root stopping boundaries (2.26), for which we know by (2.28) (and (2.24) with $z_p \downarrow 0$ as $p \rightarrow \infty$) that only finitely many moments are finite, thus again agreeing with (3.29) as $\beta \uparrow 1/2$. ■

In the next theorem we generalize and extend the result of Theorem 3.4 to the case $1 < p \leq 2$.

THEOREM 3.12. Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let τ be a stopping time for B satisfying

$$(3.31) \quad E(\tau^k) = O(C^k k^{k(2-p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$, where $1 < p \leq 2$ is fixed. Then

$$(3.32) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_p} < \infty.$$

Moreover,

$$(3.33) \quad \Delta_p(D) = \sup_{k \geq 1} \left(\frac{E(\tau^k)}{D^k k^{k(2-p)/p}}\right)$$

for $D > 0$. Then

$$(3.34) \quad \|\max_{0 \leq t \leq \tau} |B_t|\|_{\psi_p} \leq \sqrt{10} \frac{p}{p-1} e^{1/p} \sqrt{D} \left(1 + \frac{(\Delta_p(D))^{p/2}}{\sqrt{2\pi}}\right)$$

for all $D > 0$.

Proof. Given a stopping time τ for B satisfying (3.31), define $B_\tau^* = \max_{0 \leq t \leq \tau} |B_t|$. Then by (2.1), (2.3)+(2.5), (2.23) and (2.25) we have

$$(3.35) \quad E(B_\tau^*)^{kp} \leq \left(\frac{kp}{kp-1}\right)^{kp} E|B_\tau|^{kp} \leq \left(\frac{kp}{kp-1}\right)^{kp} (z_{kp}^*)^{kp} E(\tau^{kp/2}) \\ \leq \left(\frac{kp}{kp-1}\right)^{kp} (z_{2k}^*)^{kp} E(\tau^{kp/2})$$

$$\leq \left(\frac{kp}{kp-1} \right)^{kp} (\sqrt{2(4k+1)})^{kp} E(\tau^{kp/2})$$

for all $k \geq 1$. Since $kp/(kp-1) \leq p/(p-1)$ and $2(4k+1) \leq 10k$ for $k \geq 1$, from (3.35) by Taylor expansion we easily obtain

$$(3.36) \quad E \left(\exp \left(\frac{B_\tau^*}{c} \right)^p \right) - 1 = \sum_{k=1}^{\infty} \frac{E(B_\tau^*)^{kp}}{c^{kp} k!} \\ \leq \sum_{k=1}^{\infty} \left(\frac{\sqrt{10}p}{c(p-1)} \right)^{kp} \frac{k^{kp/2}}{k!} E(\tau^{kp/2})$$

for all $c > 0$. Further, by Stirling's formula we find

$$(3.37) \quad \frac{k^{kp/2}}{k!} = \frac{k^{kp/2}}{\sqrt{2\pi k} k^k e^{-k} e^{r_k}} \leq \frac{e^k}{\sqrt{2\pi} k^{k(1-p/2)+1/2}},$$

where $1/(12k+1) < r_k < 1/(12k)$ for $k \geq 1$. Inserting (3.37) into (3.36), and using Jensen's inequality, we obtain the estimate

$$(3.38) \quad E \left(\exp \left(\frac{B_\tau^*}{c} \right)^p \right) - 1 \\ \leq \frac{1}{\sqrt{2\pi}} \sum_{k=1}^{\infty} \left(\frac{\sqrt{10}pe^{1/p}}{c(p-1)} \right)^{kp} \frac{E(\tau^{kp/2})}{k^{k(2-p)/2+1/2}} \\ = \frac{1}{\sqrt{2\pi}} \sum_{k=1}^{\infty} \left(\frac{\sqrt{10}pe^{1/p}\sqrt{D}}{c(p-1)} \right)^{kp} \frac{E(\tau^{kp/2})}{D^{kp/2} k^{k((2-p)/p)(p/2)+1/2}} \\ \leq \frac{(\Delta_p(D))^{p/2}}{\sqrt{2\pi}} \sum_{k=1}^{\infty} \left(\frac{\sqrt{10}pe^{1/p}\sqrt{D}}{c(p-1)} \right)^{kp}$$

for all $c, D > 0$. Identifying the last term in (3.38) with 1, we easily find that

$$c = \sqrt{10} \frac{p}{p-1} e^{1/p} \sqrt{D} \left(1 + \frac{(\Delta_p(D))^{p/2}}{\sqrt{2\pi}} \right).$$

This proves (3.34), and hence (3.32) follows as well. The proof is complete. ■

Remark 3.13. It should be noted that the scope of Theorem 3.12 is rather limited, since $1 > (2-p)/p \downarrow 0$ as $p \uparrow 2$. It indicates that the real power of the method which is used throughout this paper lies in the cases when p is small. This is stated more explicitly in the next theorem. Finally, it is clear that the analogues of Remark 3.6 and Problem 3.10 can be formulated in the context of Theorem 3.12 as well. The details are left to the reader.

THEOREM 3.14. Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let τ be a stopping time for B satisfying

$$(3.39) \quad E(\tau^k) = O(C^k k^{k(2-p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$, where $0 < p \leq 2$ is fixed. Then

$$(3.40) \quad E(\exp(\varepsilon(\max_{0 \leq t \leq \tau} |B_t|^p))) < \infty$$

for some (small enough) $\varepsilon > 0$.

Proof. The case $p = 1$ is proved in Theorem 3.4, while the case $1 < p \leq 2$ is proved in Theorem 3.12. Consider the case $0 < p < 1$. For this, let J denote the smallest $k \geq 2$ such that $kp > 1$. Then by Taylor expansion, Jensen's inequality, and (2.2) (with $p = 1$), we have

$$(3.41) \quad E \left(\exp \left(\frac{B_\tau^*}{c} \right)^p \right) - 1 \\ = \sum_{k=1}^{J-1} \frac{E(B_\tau^*)^{kp}}{c^{kp} k!} + \sum_{k=J}^{\infty} \frac{E(B_\tau^*)^{kp}}{c^{kp} k!} \\ \leq \frac{J-1}{c^{(J-1)p}} \max_{1 \leq k \leq J-1} \frac{(3E(\sqrt{\tau}))^{kp}}{k!} + \sum_{k=J}^{\infty} \frac{E(B_\tau^*)^{kp}}{c^{kp} k!} \\ \leq \frac{J-1}{c^{(J-1)p}} ((3E(\sqrt{\tau}))^p + (3E(\sqrt{\tau}))^{(J-1)p}) + \sum_{k=J}^{\infty} \frac{E(B_\tau^*)^{kp}}{c^{kp} k!}$$

for all $c > 1$. The first term on the right-hand side in (3.41) is arbitrarily small when c becomes large, while the second term is controlled (and proved to be finite and arbitrarily small) in exactly the same way as in the proof of Theorem 3.12. This completes the proof of the theorem. ■

Remark 3.15. Note that (3.41) with (3.35)+(3.37) (as in (3.36)+(3.38)) leaves a possibility of obtaining an estimate of the quantity on the left-hand side in (3.40). The details are left to the reader.

4. Exponential integrability of continuous local martingales. The results obtained in Section 3 for a Brownian motion will be extended in this section to continuous local martingales (and Ito's integral). First we want to display the facts which make this extension possible. Recall that a process $M = (M_t)_{t \geq 0}$ is called a *local martingale* (with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$ of $(\Omega, \mathcal{F}, \mathcal{P})$) if M_0 is \mathcal{F}_0 -measurable and there exists an increasing sequence $(T_n)_{n \geq 1}$ of (\mathcal{F}_t) -stopping times with $T_n \uparrow \infty$ as $n \uparrow \infty$, such that each "stopped" process $(M_{t \wedge T_n} - M_0)_{t \geq 0}$ is a martingale (with

respect to $(\mathcal{F}_t)_{t \geq 0}$ for $n \geq 1$. If $M = (M_t)_{t \geq 0}$ is a continuous local martingale with $M_0 = 0$, then there exists a unique continuous increasing process $[M] = ([M]_t)_{t \geq 0}$ such that $M^2 - [M] = (M_t^2 - [M]_t)_{t \geq 0}$ is a continuous local martingale (see [14]). The process $[M]$ is called the *quadratic variation process* of M .

The extension mentioned above relies upon a well-known fact that every continuous local martingale is a time-changed Brownian motion. This result is due to Dubins and Schwarz [4], and more precisely may be stated as follows. Let $M = (M_t)_{t \geq 0}$ be a continuous local martingale (with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$ of $(\Omega, \mathcal{F}, \mathcal{P})$) such that $M_0 = 0$ and $[M]_t \uparrow \infty$ as $t \uparrow \infty$. Define the stopping time

$$(4.1) \quad \tau_t = \inf\{s > 0 : [M]_s = t\}$$

and set $\mathcal{G}_t = \mathcal{F}_{\tau_t}$ for $t \geq 0$. Then $B_t = M_{\tau_t}$ with $t \geq 0$ defines a standard Brownian motion B with respect to the filtration $(\mathcal{G}_t)_{t \geq 0}$. Moreover, for any given and fixed $t \geq 0$, the random variable $[M]_t$ is a (\mathcal{G}_t) -stopping time, and we have

$$(4.2) \quad M_t = B_{[M]_t}$$

for all $t \geq 0$. The result remains valid without the restriction that $[M]_t \uparrow \infty$ as $t \uparrow \infty$, but at the expense of an enlargement of the underlying probability setting $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, P)$ (for more details see [14]).

The preceding result has the following analogue for Ito's integral (see [9]). Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let $Z = (Z_t)_{t \geq 0}$ be a non-anticipating random process satisfying $\int_0^\infty Z_t^2 dt < \infty$ P -a.s. Then there exist a standard Brownian motion $\widehat{B} = (\widehat{B}_t)_{t \geq 0}$ and a stopping time τ for \widehat{B} (both depending on Z) such that

$$(4.3) \quad \text{Law}(\tau) = \text{Law}\left(\int_0^\infty Z_t^2 dt\right),$$

$$(4.4) \quad \text{Law}(\widehat{B}_\tau) = \text{Law}\left(\int_0^\infty Z_t dB_t\right).$$

In order to apply the results of Section 3 in the context of a continuous local martingale $(M_t)_{t \geq 0}$, it is enough to recall that $[M]_t$ is a stopping time for the Brownian motion $B = (B_t)_{t \geq 0}$ (relative to the filtration $(\mathcal{G}_t)_{t \geq 0}$ defined above), while by (4.2) we clearly have

$$(4.5) \quad \max_{0 \leq s \leq t} |M_s| = \max_{0 \leq s \leq t} |B_{[M]_s}| = \max_{0 \leq s \leq [M]_t} |B_s|$$

for all $t \geq 0$. These two facts clarify generalizations and extensions of Theorems 3.2, 3.4, 3.12 and 3.14, which are formulated in the following theorem.

THEOREM 4.1. *Let $M = (M_t)_{t \geq 0}$ be a continuous local martingale such that its quadratic variation process $[M] = ([M]_t)_{t \geq 0}$ satisfies*

$$(4.6) \quad E([M]_t)^k < \infty$$

for all $k \geq 1$ with some fixed $t > 0$. If

$$(4.7) \quad \left\| \max_{0 \leq s \leq t} |M_s| \right\|_{\psi_p} < \infty$$

for some $0 < p < \infty$, then

$$(4.8) \quad E([M]_t)^k = O(C^k k^{k(2+p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$. Moreover, if for some $0 < p \leq 2$,

$$(4.9) \quad E([M]_t)^k = O(C^k k^{k(2-p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$, then

$$(4.10) \quad \left\| \max_{0 \leq s \leq t} |M_s| \right\|_{\psi_p} < \infty.$$

Finally, if for $1 \leq p \leq 2$ we set

$$(4.11) \quad \Delta_p(D, t) = \sup_{k \geq 1} \left(\frac{E([M]_t)^k}{D^k k^{k(2-p)/p}} \right)$$

for $D > 0$, then

$$(4.12) \quad \left\| \max_{0 \leq s \leq t} |M_s| \right\|_{\psi_1} \leq 6e\sqrt{D} \left(1 + \frac{\sqrt{\Delta_1(D, t)}}{\sqrt{2\pi}} \right),$$

$$(4.13) \quad \left\| \max_{0 \leq s \leq t} |M_s| \right\|_{\psi_p} \leq \sqrt{10} \frac{p}{p-1} e^{1/p} \sqrt{D} \left(1 + \frac{(\Delta_p(D, t))^{p/2}}{\sqrt{2\pi}} \right),$$

for all $D > 0$ and all $1 < p \leq 2$. ■

Remark 4.2. Note that (4.9)+(4.10) with $p = 1$ states that if $M = (M_t)_{t \geq 0}$ is a continuous local martingale such that $\| [M]_t \|_{\psi_1} < \infty$ for some $t > 0$, then $\| \max_{0 \leq s \leq t} |M_s| \|_{\psi_1} < \infty$ as well. It is moreover clear that many of the facts from Section 3 (which are stated in the remarks and examples) carry over in an obvious way to the continuous local martingale case. We will not pursue this in more detail here, but instead concentrate on the Ito integral as a particular example.

The generalization and extension of the results from Section 3 to the Ito integral follow immediately by use of (4.3) and (4.4). This is formulated in the following theorem.

THEOREM 4.3. *Let $B = (B_t)_{t \geq 0}$ be a standard Brownian motion, and let $Z = (Z_t)_{t \geq 0}$ be a non-anticipating random process satisfying*

$$(4.14) \quad E\left(\int_0^\infty Z_t^2 dt\right)^k < \infty$$

for all $k \geq 1$. If

$$(4.15) \quad \left\| \int_0^\infty Z_t dB_t \right\|_{\psi_p} < \infty$$

for some $0 < p < \infty$, then

$$(4.16) \quad E \left(\int_0^\infty Z_t^2 dt \right)^k = O(C^k k^{k(2+p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$. Moreover, if for some $0 < p \leq 2$,

$$(4.17) \quad E \left(\int_0^\infty Z_t^2 dt \right)^k = O(C^k k^{k(2-p)/p})$$

for some $C > 0$ as $k \rightarrow \infty$, then

$$(4.18) \quad \left\| \int_0^\infty Z_t dB_t \right\|_{\psi_p} < \infty.$$

Finally, if for $1 \leq p \leq 2$ we set

$$(4.19) \quad \Delta_p(D, Z) = \sup_{k \geq 1} \left(\frac{E \left(\int_0^\infty Z_t^2 dt \right)^k}{D^k k^{k(2-p)/p}} \right)$$

for $D > 0$, then

$$(4.20) \quad \left\| \int_0^\infty Z_t dB_t \right\|_{\psi_1} \leq 6e\sqrt{D} \left(1 + \frac{\sqrt{\Delta_1(D, Z)}}{\sqrt{2\pi}} \right),$$

$$(4.21) \quad \left\| \int_0^\infty Z_t dB_t \right\|_{\psi_p} \leq \sqrt{10} \frac{p}{p-1} e^{1/p} \sqrt{D} \left(1 + \frac{(\Delta_p(D, Z))^{p/2}}{\sqrt{2\pi}} \right),$$

for all $D > 0$ and all $1 < p \leq 2$. ■

Remark 4.4. Note that (4.17)+(4.18) with $p = 1$ states that if $B = (B_t)_{t \geq 0}$ is a Brownian motion and $Z = (Z_t)_{t \geq 0}$ is a non-anticipating process such that $\left\| \int_0^\infty Z_t^2 dt \right\|_{\psi_1} < \infty$, then $\left\| \int_0^\infty Z_t dB_t \right\|_{\psi_1} < \infty$ as well.

Acknowledgments. Thanks are due to S. E. Graversen for valuable discussions and communicating facts used in Example 3.11, and to J. Hoffmann-Jørgensen for his constant advice and support.

References

- [1] M. Abramowitz and I. A. Stegun, *Pocketbook of Mathematical Functions* (abridged edition of *Handbook of Mathematical Functions*, National Bureau of Standards, 1964), Verlag Harri Deutsch, Thun-Frankfurt/Main, 1984.
- [2] D. L. Burkholder and R. F. Gundy, *Extrapolation and interpolation of quasi-linear operators on martingales*, Acta Math. 124 (1970), 249–304.

- [3] B. Davis, *On the L^p norms of stochastic integrals and other martingales*, Duke Math. J. 43 (1976), 697–704.
- [4] L. E. Dubins and G. Schwarz, *On continuous martingales*, Proc. Nat. Acad. Sci. U.S.A. 53 (1965), 913–916.
- [5] A. Gut, *Stopped Random Walks (Limit Theorems and Applications)*, Springer, 1988.
- [6] U. Haagerup, *The best constants in the Khintchine inequality*, Studia Math. 70 (1982), 231–283.
- [7] M. Ledoux and M. Talagrand, *Probability in Banach Spaces (Isoperimetry and Processes)*, Springer, Berlin, 1991.
- [8] R. Sh. Liptser and A. N. Shiriyayev, *Theory of Martingales*, Kluwer Academic Publ., 1989.
- [9] H. P. McKean, *Stochastic Integrals*, Wiley, 1969.
- [10] A. A. Novikov, *On stopping times for a Wiener process*, Theory Probab. Appl. 16 (1971), 449–456.
- [11] G. Peškir, *Best constants in Kahane–Khintchine inequalities in Orlicz spaces*, J. Multivariate Anal. 45 (1993), 183–216.
- [12] —, *Maximal inequalities of Kahane–Khintchine's type in Orlicz spaces*, Math. Proc. Cambridge Philos. Soc. 115 (1994), 175–190.
- [13] —, *Best constants in Kahane–Khintchine inequalities for complex Steinhaus variables*, Proc. Amer. Math. Soc., to appear.
- [14] L. C. G. Rogers and D. Williams, *Diffusions, Markov Processes, and Martingales*, Vol. 2, *Ito's Calculus*, Wiley, 1987.
- [15] L. A. Shepp, *A first passage problem for the Wiener process*, Ann. Math. Statist. 38 (1967), 1912–1914.
- [16] G. Wang, *Sharp maximal inequalities for conditionally symmetric martingales and Brownian motion*, Proc. Amer. Math. Soc. 112 (1991), 579–586.

INSTITUTE OF MATHEMATICS
UNIVERSITY OF AARHUS
NY MUNKEGADE
8000 AARHUS, DENMARK
E-mail: GORAN@MIAU.DK

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ZAGREB
BIJENIČKA 30
41000 ZAGREB, CROATIA

Received May 4, 1995

(3459)