

- [45] G. Pisier, *De nouvelles caractérisations des ensembles de Sidon*, in: Mathematical Analysis and Applications, Part B, Adv. in Math. Suppl. Stud. 7B, Academic Press, 1981, 685–726.
- [46] —, *Conditions d'entropie et caractérisations arithmétiques des ensembles de Sidon*, in: Topics in Modern Harmonic Analysis II (Torino/Milano, 1982), L. De Michele and F. Ricci (eds.), Ist. Naz. Alta Mat. Francesco Severi, 1983, 911–944.
- [47] A. I. Plotkin, *Continuation of L^p -isometries*, J. Soviet Math. 2 (1974), 143–165.
- [48] S. Ramanujan, *Notebooks*, Tata Inst. Fundam. Research, 1957.
- [49] S. K. Rao, *On sums of sixth powers*, J. London Math. Soc. 9 (1934), 172–173.
- [50] M. Rosenblatt, *A central limit theorem and a strong mixing condition*, Proc. Nat. Acad. Sci. U.S.A. 42 (1956), 43–47.
- [51] H. P. Rosenthal, *Sous-espaces de L^1 . Cours de troisième cycle*, Université Paris 6, 1979 (unpublished).
- [52] W. Rudin, *Trigonometric series with gaps*, J. Math. Mech. 9 (1960), 203–228.
- [53] —, *L^p -isometries and equimeasurability*, Indiana Univ. Math. J. 25 (1976), 215–228.
- [54] I. Z. Ruzsa, *On difference sets*, Studia Sci. Math. Hungar. 13 (1978), 319–326.
- [55] I. Singer, *Bases in Banach Spaces II*, Springer, 1981.
- [56] C. M. Skinner and T. D. Wooley, *On equal sums of two powers*, J. Reine Angew. Math. 462 (1995), 57–68.
- [57] C. L. Stewart and R. Tijdeman, *On infinite-difference sets*, Canad. J. Math. 31 (1979), 897–910.

Équipe d'Analyse
 Université Paris VI
 Boîte 186
 4, place Jussieu
 F-75252 Paris Cedex 05, France
 E-mail: neuwirth@ccr.jussieu.fr

Received August 4, 1997
 Revised version January 12, 1998

(3931)

On operators satisfying the Rockland condition

by

WALDEMAR HEBISCH (Wrocław)

Abstract. Let G be a homogeneous Lie group. We prove that for every closed, homogeneous subset Γ of G^* which is invariant under the coadjoint action, there exists a regular kernel P such that P goes to 0 in any representation from Γ and P satisfies the Rockland condition outside Γ . We prove a subelliptic estimate as an application.

Introduction. The purpose of this paper is to construct operators which satisfy the Rockland condition in a given set Γ of representations, and are equal to 0 outside Γ . Rockland operators satisfy remarkable subelliptic estimates ([11], [7], [9], [10], [14]; see also [15]) making them a good substitute for elliptic operators on homogeneous groups. Christ *et al.* [2] gave a calculus for pseudodifferential operators on homogeneous groups: the formulas for products and adjoints and criteria for existence of left or right parametrices (generalizing results of [8]). However, one should note that the great flexibility of the classical calculus of pseudodifferential operators is in large part due to the ease of constructing scalar functions (cutoffs and partitions of unity). In the homogeneous group case we want to pre-specify operators in a set of representations and still have regular kernels; this is not straightforward, in fact not always possible. Our kernels may serve as cutoffs on spectral side (for the spatial cutoffs one simply uses multiplications with smooth functions). The conditions we impose seem to be necessary. We present also a simple application in which we derive some L^p estimates.

Acknowledgements. I would like to thank Jean Nourrigat for valuable suggestions.

Preliminaries. We consider a homogeneous group G , that is, a nilpotent Lie group equipped with a family of automorphisms (*dilations*) $\{\delta_t\}_{t>0}$ such that $\delta_t\delta_s = \delta_{ts}$ and for all $x \in G$ we have $\delta_t x \rightarrow e$ as $t \rightarrow 0$. The reader may wish to consult [6] (our definition is a bit more general). We identify G with

1991 *Mathematics Subject Classification*: Primary 22E30.
 This work is part of the research project 2 P301 052 07.

its Lie algebra via the exponential map, and write 0 instead of e . With our identification all δ_t become linear maps.

As $\det(\delta_t)$ must be a power of t there exists a number $Q > 0$ such that for all bounded measurable $A \subset G$,

$$|\delta_t A| = t^Q |A|;$$

this Q is called the *homogeneous dimension* of G . More generally, one can take t to be discrete, that is, consider a dilation operator D such that $D^{-k}x \rightarrow e$ as $k \rightarrow \infty$.

A distribution T on G is said to be a *kernel of order* $r \in \mathbb{C}$ if T coincides with a locally finite measure away from the origin, and is homogeneous of degree $-r - Q$, that is,

$$(f \circ \delta_t, T) = t^r (f, T)$$

for all $f \in C_c^\infty(G)$ and $t > 0$. We extend the action of dilations to distributions by the formula

$$(f, \delta_t T) = (f \circ \delta_t, T).$$

Then T is a kernel of order r iff for all $t > 0$,

$$\delta_t T = t^r T.$$

A kernel is called *regular* if it coincides with a smooth function away from the origin.

In the sequel we will identify right-invariant vector fields on G with distributions supported in $\{0\}$. More precisely, there is one-to-one correspondence between right-invariant differential operators and distributions supported in $\{0\}$. To get the identification we write

$$(X, f) = Xf(0).$$

Then $\tilde{X}f = X * f$, and for the left-invariant field \tilde{X} corresponding to X we have $\tilde{X}f = f * X$. We also note that dilating a vector field as an element of the Lie algebra and as a distribution gives the same result.

For a unitary representation π of G on a Hilbert space H and a kernel T of order r with $\Re(r) > 0$, the operator $\pi(T)$ is defined on the space $C^\infty(\pi)$ of smooth vectors for π by

$$(g, \pi(T)f) = (\phi_{f,g}, T)$$

where $\phi_{f,g}(x) = (g, \pi(x)f)$. An equivalent definition is

$$\pi(T)f = T * \psi_f(e)$$

where $\psi_f(x) = \pi(x)f$. This definition also makes sense for uniformly bounded representations on Banach spaces. If $\Re(r) \leq 0$ the situation is

more tricky. For a regular kernel T one may find $h \in C_c^\infty(G)$ such that

$$Tf = \sum_k 2^{-rk} \delta_{2^k}(h) * f.$$

For $r = 0$ one must have $\int h = 0$ (otherwise T would not be a distribution), and the Cotlar–Stein lemma shows that the sum defining T is strongly convergent in any unitary representation of G to a bounded operator.

If $\Re(r) < 0$ then T defines an unbounded operator on $L^2(G)$ (to see that T is densely defined cf. [2]). The following lemma shows that the problem is caused by the trivial representation.

LEMMA 1. *Let G be a homogeneous nilpotent Lie group, π be a nontrivial irreducible unitary representation of G and h be a Schwartz class function on G . Then for any m there is a continuous seminorm $C_m(\cdot)$ such that $\|\pi(\delta_t(h))\| \leq C_m(h)(1+t)^{-m}$.*

Proof. We fix a scalar product (so also a norm $|\cdot|$) on the Lie algebra of G . Note that there exists $\alpha > 0$ such that if X is an element of the Lie algebra of G then $|\delta_{t^{-1}}(X)| \leq C(1+t)^{-\alpha}$ for $t > 1$. Let X_j span the Lie algebra of G . Put $L = \sum X_j^2$. It is known that L is invertible in π . So

$$\text{id}_\pi = \sum \pi(X_j)(\pi(X_j)\pi(L)^{-1}) = \sum \pi(X_j)E_j$$

where $E_j = \pi(X_j)\pi(L)^{-1}$ are bounded operators in π . Next, we write

$$\pi(\delta_t(h)) = \pi(\delta_t(h)) \sum \pi(X_j)E_j = \sum \pi(\delta_t(h * \delta_{t^{-1}}(X_j)))E_j.$$

Inductively, for any natural l ,

$$\pi(\delta_t(h)) = \sum \pi(\delta_t(h * \delta_{t^{-1}}(X_{j_1}) * \dots * \delta_{t^{-1}}(X_{j_l})))E_{j_1} \dots E_{j_l}$$

so

$$\begin{aligned} \|\pi(\delta_t(h))\| &\leq C_l \max \|\pi(\delta_t(h * \delta_{t^{-1}}(X_{j_1}) * \dots * \delta_{t^{-1}}(X_{j_l})))\| \\ &\leq C_l \max \|h * \delta_{t^{-1}}(X_{j_1}) * \dots * \delta_{t^{-1}}(X_{j_l})\|_{L^1} \\ &\leq C_l \max C_{h,l} |\delta_{t^{-1}}(X_{j_1})| \dots |\delta_{t^{-1}}(X_{j_l})| \leq C'(h, l)(1+t)^{-\alpha l}, \end{aligned}$$

which gives the claim.

If we put

$$V_s = \{f \in L^1_{\text{loc}}(G) \cap C^\infty(G - \{0\}) : \forall \phi \in C_c^\infty(G - \{0\}) \lim_{t \rightarrow 0} t^s \phi \delta_t f = 0$$

uniformly with all derivatives}

then V_s is a locally convex metrizable vector space and the Schwartz class functions are dense in V_s . For $0 > \Re(r) > -s$ regular kernels of order r are in V_s and (1.1) shows that π has a (unique) extension from the Schwartz class to V_s .

Main results. Let π_l for $l \in G^*$ be the representation associated with l according to the Kirillov theory (cf. [13]).

THEOREM 2. *Let G be a homogeneous Lie group with dilations $\{\delta_t\}_{t>0}$, and Γ be a closed subset of G^* such that $\text{Ad}^*(G)\Gamma \subset \Gamma$ and $\delta_t\Gamma \subset \Gamma$ for all $t > 0$. For every $\alpha \geq 0$ there exists a regular kernel P of order α such that for all $l \in \Gamma$ we have $\pi_l(P) = 0$ and for all $l \notin \Gamma$ the operator $\pi_l(P)$ is positive definite and injective on its domain. For every $0 > \alpha > -Q$ there exists a kernel satisfying the above conditions, except for $l = 0$. Moreover, there is a Schwartz class function H on G such that for all $l \in \Gamma$ we have $\pi_l(H) = 0$ and for all $l \notin \Gamma$ the operator $\pi_l(H)$ is positive definite and injective.*

Proof. It is enough to prove the theorem only for $-Q < \alpha \leq 0$ and small $\alpha > 0$. Indeed, taking a sufficiently high power of P we get α as large as we wish, without destroying other properties of P . Moreover, we only need to prove the last claim, that is, to construct a Schwartz class function H such that for all $l \in \Gamma$ we have $\pi_l(H) = 0$ and for all $l \notin \Gamma$ the operator $\pi_l(H)$ is positive definite and injective. If we have such a function and α is small enough, then

$$P = \int_0^\infty t^{-\alpha} \delta_t H \frac{dt}{t}$$

gives us a regular kernel of order α having the required properties. Indeed, 0 is in Γ so $\int H = 0$. Moreover, there is $\alpha_0 > 0$ such that for all $0 < t < 1$ we have $|\delta_t x| \leq t^{\alpha_0} |x|$. Fix a $\phi \in C_c^\infty(G)$. For $0 < t < 1$ we have

$$|(\phi, \delta_t H)| = |(\phi - \phi(0), \delta_t H)| \leq Ct^{\alpha_0}$$

so if $\alpha < \alpha_0$, then

$$\int_0^\infty t^{-\alpha} (\phi, \delta_t H) \frac{dt}{t}$$

is absolutely convergent. Changing variables in the integral above, one easily checks that P is homogeneous of degree $-\alpha - Q$. Smoothness of P off $\{0\}$ is clear. Also, if $\alpha > 0$, it is easy to check that

$$\pi(P) \subset \int_0^\infty t^{-\alpha} \pi(\delta_t H) \frac{dt}{t}$$

and that the right hand side defines a closed injective operator (we compute the integral applying the function under the integral to a vector; the domain is the set of all vectors for which the integral is convergent). For $\alpha \leq 0$ we get the same conclusion using Lemma 1 (condition $\alpha > -Q$ is used to prove that P is a regular kernel).

We are going to build H . Let us recall that by [1] the set $\tilde{\Gamma} = \{\pi_l : l \in \Gamma\}$ is closed in the Fell topology of the space of representations. Fix $p \notin \Gamma$. Then

$\text{Ad}^*(G)\Gamma \subset \Gamma$ implies $\pi_p \notin \tilde{\Gamma}$. By the definition of the Fell topology and density of $C_c^\infty(G)$ in $C^*(G)$, there exists a function $F \in C_c^\infty(G)$ such that

$$\|\pi_p(F)\| = 1 \quad \text{and} \quad \|\pi_l(F)\| < 1/10 \quad \text{for all } l \in \Gamma.$$

Replacing F by $F^* * F$ we may assume that F is positive definite. Choose $\phi \in C_c^\infty(\mathbb{R})$ such that $\phi(1) = 1$, $\phi \geq 0$ and $\text{supp}(\phi) \subset [1/10, 2]$. By the spectral theorem we have

$$\phi(\pi_p(F)) \neq 0 \quad \text{while} \quad \phi(\pi_l(F)) = 0 \quad \text{for all } l \in \Gamma.$$

Using functional calculus, as for example in [12], we show that there exists a Schwartz class function R on G such that (as a convolution operator on $L^2(G)$) $R = \phi(F)$. Approximating ϕ by polynomials we see that $\phi(\pi_l(F)) = \pi_l(R)$ for all l . We also note that the set of π such that $\pi(R) \neq 0$ is open (by definition).

To summarize, we constructed R such that $\pi(R) = 0$ for all $\pi \in \tilde{\Gamma}$, the set $U_R = \{\pi(R) \neq 0\}$ is open and $\pi_p \in U_R$. Since the Fell topology has a countable base, there exists a sequence $\{R_i\}_{i \in \mathbb{N}}$ such that the complement of $\tilde{\Gamma}$ is the union of U_{R_i} . Therefore, putting $S = \sum a_i R_i$ where a_i are positive and small enough for S to be a Schwartz class function, we see that $\pi(S) = 0$ for all π in $\tilde{\Gamma}$, and $\pi(S) \neq 0$ on the complement of $\tilde{\Gamma}$.

To finish the proof we need the following lemma:

LEMMA 3. *If π is an irreducible unitary representation of G , the sequence $\{g_j\}_{j \in \mathbb{N}}$ is dense in G , $f \in L^1(G)$, $\pi(f) \neq 0$, $\pi(f) \geq 0$, and the sequence $\{c_j\}_{j \in \mathbb{N}}$ is positive and summable, then*

$$A = \pi \left(\sum c_j \delta_{g_j^{-1}} f \delta_{g_j} \right)$$

(where δ_{g_j} is the convolution operator with unit mass at g_j) is injective.

Proof. Suppose that, on the contrary, A is not injective. Then there exists a nonzero v such that

$$(Av, v) = \sum c_j (\pi(\delta_{g_j^{-1}}) \pi(f) \pi(\delta_{g_j}) v, v) = 0,$$

hence for each j ,

$$(\pi(f)^{1/2} \pi(\delta_{g_j}) v, \pi(f)^{1/2} \pi(\delta_{g_j}) v) = 0,$$

or simply

$$\pi(f)^{1/2} \pi(\delta_{g_j}) v = 0.$$

Since π is irreducible, the closed linear span of $\pi(\delta_{g_j}) v$ gives the whole space, so $\pi(f)^{1/2} = 0$. As $\pi(f)$ is nonzero this gives a contradiction.

Choosing $c_j = \exp(-j - |g_j|)$ and applying Lemma 3 we conclude that

$$H = \sum c_j \delta_{g_j^{-1}} S \delta_{g_j}$$

is a Schwartz class function such that the operator $\pi(H)$ is injective for π in the complement of $\tilde{\Gamma}$, and $\pi(H) = 0$ for π in $\tilde{\Gamma}$, which ends the proof of Theorem 2.

REMARK. If P is a regular kernel of order α and $\Gamma = \{l : \pi_l(P) = 0\}$, then $\text{Ad}^*(G)\Gamma \subset \Gamma$, $\delta_t\Gamma \subset \Gamma$ for all $t > 0$ and Γ is closed. More precisely, if $\Re(\alpha) < 0$, then $\Gamma - \{0\}$ is closed in $G^* - \{0\}$. The first condition is clear. Γ is invariant under dilations because P is homogeneous. To see that Γ is closed assume first that $\Re(\alpha) > 0$. Let ϕ_n be an approximate unit in L^1 consisting of C_c^∞ functions. Then $\phi_n * P \in L^1$ so $\Gamma_n = \{l : \pi_l(\phi_n * P) = 0\}$ is closed. As $\Gamma = \bigcap_n \Gamma_n$ we see that Γ is closed. If $\Re(\alpha) \leq 0$, then we compose P with a kernel R such that $\pi_l(R)$ is injective on smooth vectors for all $l \neq 0$ and $P * R$ has order with positive real part.

An application. As an application of our construction we give an extension to L^p of a theorem by J. Nourrigat ([14], Théorème 1.3). We need some setup to state the theorem. Let Ω be a measure space with measure μ . Assume G acts on Ω preserving the measure. Let $\phi : G \times \Omega \rightarrow \mathbb{C}$ be a (measurable) cocycle for this action, that is, $|\phi| = 1$ and for all $g_1, g_2 \in G$ and all $x \in \Omega$,

$$\phi(g_1 g_2, x) = \phi(g_1, x) \phi(g_2, g_1^{-1} x).$$

Then the formula

$$\pi(g)f(x) = \phi(g, x)f(g^{-1}x)$$

gives a continuous representation of G which acts by isometries on $L^p(\Omega)$, $1 \leq p < \infty$ (on L^∞ we get isometries, but the action is only weak-* continuous). We say that π is a *cocycle representation*. The set $C^\infty(\pi)$ of smooth vectors is defined as usual (of course it may depend on p). Note that $\pi(C_c^\infty(G))(L^1 \cap L^\infty)$ is dense in $C^\infty(\pi)$ so we may do all the calculations on the common core. We also note that the usual construction of π_l gives a cocycle representation so we may consider π_l as representations on L^p . Let us also sketch the proof of the following well-known lemma:

LEMMA 4. *If W is a regular kernel of order α , $\Re(\alpha) = 0$ and W gives a bounded operator on $L^2(G)$ then W gives a bounded operator on $L^p(G)$, $1 < p < \infty$.*

REMARK. In fact, a regular kernel W of order α with $\Re(\alpha) = 0$ is always bounded on L^2 .

PROOF (of Lemma 4). This follows from [3], Chapitre III, Théorème (2.4). G equipped with a homogeneous norm is a space of homogeneous type. One may easily verify that in [3] the assumption that the kernel K is in L^2 is only used to prove that the operator T is *associated* with the kernel, that is, $Tf(x) = \int K(x, y)f(y) dy$ for x not in the support of f .

An alternative approach is to regularize W . We fix $\phi \in C_c^\infty(G)$ such that $\int \phi = 1$ and we write $\phi_t = \delta_t \phi$. One may check that for the regularized kernels $W_t = (\phi_t - \phi_{t-1}) * W$ the assumptions of [3], Chapitre III, Théorème (2.4), hold with bounds independent of the regularization (and $\lim_{t \rightarrow \infty} W_t f = Wf$ for f in $L^p(G)$, $1 < p < \infty$). This approach shows that the transference principle is applicable to W .

THEOREM 5. *Let G be a homogeneous Lie group with dilations $\{\delta_t\}_{t>0}$, and Γ be a closed subset of G^* such that $\text{Ad}^*(G)\Gamma \subset \Gamma$ and $\delta_t\Gamma \subset \Gamma$ for all $t > 0$. Let π be a cocycle representation of G such that all irreducible components of π are of the form π_l with $l \in \Gamma$. Let R be a regular kernel of order α with $\Re(\alpha) > 0$ such that for all $l \in \Gamma$, $l \neq 0$, the operator $\pi_l(R)$ is injective on the C^∞ vectors of π_l . Then for every $1 < p < \infty$, every positive integer k and every kernel A of order β , where $0 \leq \Re(\beta) \leq k\Re(\alpha)$, there exists $C_{p,k,A}$ such that*

$$\forall f \in C^\infty(\pi) \quad \|\pi(A)f\|_{L^p} \leq C_{p,k,A} (\|f\|_{L^p} + \|\pi(R)^k f\|_{L^p}).$$

If R is of order α and $0 \leq \alpha < Q$, then there exists a regular kernel B of order $-\alpha$ such that

$$\forall l \in \Gamma - \{0\} \forall f \in C^\infty(\pi_l) \quad \pi_l(B)\pi_l(R)f = f.$$

PROOF. First, assume that $k = 1$ and $\beta = \alpha$. Let S be a regular kernel of order $2\Re(\alpha)$ given by Theorem 2. We put

$$T = S + R^*R.$$

Then T is a regular kernel of order $2\Re(\alpha)$ and the image of T in any non-trivial representation of G is injective on the smooth vectors. We are going to construct the inverse of T . There exists an injective positive definite operator P on $L^2(G)$ such that for $s > -Q$ the operator P^s is given by a regular kernel of order s , and for small $s > 0$, P^s generates a semigroup of symmetric probability measures (see [2], Theorem 6.1, and [7]). Choose $s_0 < Q$ and $m \in \mathbb{N}$ such that $s_0 m = 2\Re(\alpha)$. Put $V = P^{s_0}$ and $U = V^{-m}T$. One easily checks that U is given by a regular kernel of order 0 and that the image of U in any nontrivial irreducible unitary representation of G is injective on the smooth vectors, so (by [2], Theorem 6.2) U is left-invertible on $L^2(G)$ and the inverse is given by a regular kernel U^{-1} .

Now the last claim follows if we notice that $U^{-1}V^{-m}R^*$ is a regular kernel of order $-\alpha$ for $\Re(\alpha) < Q$ and that

$$\begin{aligned} \pi(U^{-1}V^{-m}R^*)\pi(R)f &= \pi(U^{-1})\pi(V^{-1})^m\pi(R^*)\pi(R)f \\ &= \pi(U^{-1})\pi(V^{-1})^m\pi(T)f = \pi(U^{-1}V^{-m}T)f = f. \end{aligned}$$

Also the operator $AT^{-1}R^*$ is given by the regular kernel $W = AU^{-1}V^{-m}R^*$ of order 0. The operator obtained from W is bounded on $L^p(G)$ (by Lem-

ma 4) so by the transference principle [4] the image of W in π is bounded on $L^p(\pi)$. Hence

$$\begin{aligned}\pi(A)f &= \pi(A)\pi(U^{-1}V^{-m}T)f = \pi(A)\pi(U^{-1})\pi(V^{-1})^m\pi(S + R^*R)f \\ &= \pi(A)\pi(U^{-1})\pi(V^{-1})^m\pi(R^*)\pi(R)f \\ &= \pi(AU^{-1}V^{-m}R^*)\pi(R)f = \pi(W)\pi(R)f\end{aligned}$$

and

$$\|\pi(A)f\| \leq \|\pi(W)\| \|\pi(R)f\|.$$

If $\Re(\beta) = \Re(\alpha)$, then

$$\|\pi(A)f\| \leq C\|\pi(P^\beta)f\| \leq C\|P^{\beta-\alpha}\|_{L^p(G), L^p(G)}\|\pi(P^\alpha)f\| \leq C'\|\pi(R)f\|.$$

Here $\|P^{\beta-\alpha}\|_{L^p(G), L^p(G)}$ is finite by Lemma 4.

If $\beta < \alpha$ then we note that $A(1 + P^\alpha)^{-1}$ is convolution with an L^1 function (this follows from estimates in [5]) so

$$\|Af\| \leq C\|1 + P^\alpha f\| \leq C'(\|f\| + \|Rf\|).$$

If $k > 1$ we simply replace R by R^k .

References

- [1] I. D. Brown, *Dual topology of a nilpotent Lie group*, Ann. Sci. École Norm. Sup. 6 (1973), 407–411.
- [2] M. Christ, D. Geller, P. Głowacki and L. Pollin, *Pseudodifferential operators on groups with dilations*, Duke Math. J. 68 (1992), 31–65.
- [3] R. Coifman et G. Weiss, *Analyse Harmonique Non-Commutative sur Certains Espaces Homogènes*, Lecture Notes in Math. 242, Springer, Berlin, 1971.
- [4] —, —, *Transference Methods in Analysis*, CBMS Regional Conf. Ser. in Math. 31, Amer. Math. Soc., Providence, 1977.
- [5] J. Dziubański, *A remark on a Marcinkiewicz–Hörmander multiplier theorem for some non-differential convolution operators*, Colloq. Math. 58 (1989), 77–83.
- [6] G. B. Folland and E. M. Stein, *Hardy Spaces on Homogeneous Groups*, Princeton Univ. Press, 1982.
- [7] P. Głowacki, *Stable semi-groups of measures as commutative approximate identities on non-graded homogeneous groups*, Invent. Math. 83 (1986), 557–582.
- [8] —, *The inversion problem for singular integral operators on homogeneous groups*, Studia Math. 87 (1987), 53–69.
- [9] —, *The Rockland condition for nondifferential convolution operators*, Duke Math. J. 58 (1989), 371–395.
- [10] —, *The Rockland condition for nondifferential convolution operators II*, Studia Math. 98 (1991), 99–114.
- [11] B. Helffer et J. Nourrigat, *Caractérisation des opérateurs hypoelliptiques homogènes invariants à gauche sur un groupe gradué*, Comm. Partial Differential Equations 3 (1978), 889–958.
- [12] A. Hulanicki and J. W. Jenkins, *Nilpotent Lie groups and summability of eigenfunction expansions of Schrödinger operators*, Studia Math. 80 (1984), 235–244.

- [13] A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Uspekhi Mat. Nauk 17 (4) (1962), 57–110 (in Russian).
- [14] J. Nourrigat, *Inégalités L^2 et représentations de groupes nilpotents*, J. Funct. Anal. 74 (1987), 300–327.
- [15] —, *L^2 inequalities and representations of nilpotent groups*, C.I.M.P.A. School of Harmonic Analysis, Wuhan, to appear.

Institute of Mathematics

Wrocław University

Pl. Grunwaldzki 2/4

50-384 Wrocław, Poland

E-mail: hebisch@math.uni.wroc.pl

Web: <http://www.math.uni.wroc.pl/~hebisch/Title.html>

Institute of Mathematics

Polish Academy of Sciences

Śniadeckich 8

00-950 Warszawa, Poland

Received September 26, 1997

(3967)