

CORRECTIONS (received July 2, 1990):

“Almost everywhere summability of eigenfunction expansions associated to elliptic operators” by Waldemar Hebisch, *Studia Math.* 96 (3) (1990), 263–275

P. 265, l. 1:  $\mathcal{L}_{p,\text{loc}}^l(\mathbf{R}) := \{f: \varphi f \in \mathcal{L}_p^l(\mathbf{R}) \text{ for all } \varphi \in C_c^\infty(\mathbf{R})\}$ .

P. 265, l. 10 from the bottom: add the condition  $a, b \geq 0$ .

Theorem (3.1): add  $b > 0$  in the assumptions and  $c > 0$  in the proof.

P. 267, l. 8:  $\|K_n\|$  should have subscript  $B((1+R^n d)^a)$ ; l. 2 from the bottom: for “that” read “that if  $b = 1$ ”.

P. 268, l. 13: for “ $\alpha' = \alpha - \varepsilon$ ” read “if  $\alpha' = \alpha - q\varepsilon > 0$ ”; l. 2 from the bottom should read: “The same argument but using (2.3') instead of (2.3) also yields”.

P. 270, bottom line: a factor  $C$  on the RHS is missing.

P. 271, l. 9 from the bottom: for “ $t > 0$ ” read “ $s, t > 0$ ”.

P. 272, l. 16: the operator is assumed to have  $C^\infty$  coefficients.

Lemma (8.4): replace  $\partial^\alpha(\varrho \circ \varphi_i)$  by  $\partial^\alpha(\varrho \circ \varphi_i^{-1})$  and add the condition  $\int u = 1$  in the proof.

P. 273, † 14: for “scalar operator” read “operator in  $\mathbf{R}^n$ ”; l. 21: for  $\gamma$  read  $|\gamma|$ ; add “ $C_1 > 0$ ” in the second line of (8.6).

P. 274, l. 15: for  $\lambda^{q/2}$  read  $\lambda^{-q/2}$ ; l. 16: for  $s^m$  read  $s^{2km}$ ; l. 18: for  $s^m$  read  $s^{km}$  and for  $\|\dots f\|_0^{1/2}$  read  $\|\dots f\|_0$ ; l. 20: for  $ts^m$  read  $(ts^m)^k$ .